


Maharashtra Board Solutions Class 11-Chemistry: Chapter 12- Chemical Equilibrium

Class 11 - Chapter 12 Chemical Equilibrium



For any clarifications or questions you can write to info@indcareer.com

Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888, **Website:** <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-12-chemical-equilibrium/>



Maharashtra Board Solutions Class 11-Chemistry: Chapter 12- Chemical Equilibrium

Class 11: Chemistry Chapter 12 solutions. Complete Class 11 Chemistry Chapter 12 Notes.

Maharashtra Board Solutions Class 11-Chemistry: Chapter 12- Chemical Equilibrium

Maharashtra Board 11th Chemistry Chapter 12, Class 11 Chemistry Chapter 12 solutions

1. Choose the correct option

Question A.

The equilibrium, $\text{H}_2\text{O}_{(l)} \rightleftharpoons \text{H}^+_{(aq)} + \text{OH}^-_{(aq)}$ is

- a. dynamic
- b. static
- c. physical
- d. mechanical

Answer:

- a. dynamic

Question B.

For the equilibrium, $\text{A} \rightleftharpoons 2\text{B} + \text{Heat}$, the number of 'A' molecules increases if

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

- a. volume is increased
- b. temperature is increased
- c. catalyst is added
- d. concentration of B is decreased

Answer:

- b. temperature is increased

Question C.

For the equilibrium $\text{Cl}_{2(g)} + 2\text{NO}_{(g)} \rightleftharpoons 2\text{NOCl}_{(g)}$ the concentration of NOCl will increase if the equilibrium is disturbed by

- a. adding Cl_2
- b. removing NO
- c. adding NOCl
- d. removal of Cl_2

Answer:

- a. adding Cl_2

Question D.

The relation between K_c and K_p for the reaction $\text{A}_{(g)} + \text{B}_{(g)} \rightleftharpoons 2\text{C}_{(g)} + \text{D}_{(g)}$ is

- a. $K_c = K_p/RT$
- b. $K_p = K_c^2$
- c. $K_c = 1K_p\sqrt{}$
- d. $K_p/K_c = 1$

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

a. $K_c = K_p/RT$

Question E.

When volume of the equilibrium reaction $C_{(s)} + H_2O_{(g)} \rightleftharpoons CO_{(g)} + H_{2(g)}$ is increased at constant temperature the equilibrium will

- a. shift from left to right
- b. shift from right to left
- c. be unaltered
- d. can not be predicted

Answer:

- a. shift from left to right

2. Answer the following

Question A.

State Law of Mass action.

Answer:

Law of mass action: The law of mass action states that the rate of a chemical reaction at each instant is proportional to the product of concentrations of all the reactants.

Question B.

Write an expression for equilibrium constant with respect to concentration.

Answer:

For a reversible chemical reaction at equilibrium, $aA + bB \rightleftharpoons cC + dD$

Equilibrium constant (K_c) = $\frac{[C]^c[D]^d}{[A]^a[B]^b}$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>


Question C.

Derive mathematically value of K_p for $A_{(g)} + B_{(g)} \rightleftharpoons C_{(g)} + D_{(g)}$.

Answer:

When the concentrations of reactants and products in gaseous reactions are expressed in terms of their partial pressure, then the equilibrium constant is represented as K_p .

$\therefore$ For the reaction,





the equilibrium constant (K_c) can be expressed using partial pressure as: $K_p = \frac{P_C \times P_D}{P_A \times P_B}$


Where P_A , P_B , P_C and P_D are equilibrium partial pressures of A, B, C and D respectively.

Question D.


Write expressions of K_c for following chemical reactions



Answer:



$$K_c = \frac{[SO_3]^2}{[SO_2]^2 [O_2]}$$



$$K_c = \frac{[NO_2]^2}{[N_2O_4]}$$

Question E.

Mention various applications of equilibrium constant.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Various applications of equilibrium constant:

- Prediction of the direction of the reaction
- To know the extent of the reaction
- To calculate equilibrium concentrations
- Link between chemical equilibrium and chemical kinetics

Question F.

How does the change of pressure affect the value of equilibrium constant ?

Answer:

The change of pressure does not affect the value of equilibrium constant.

Question J.

Differentiate irreversible and reversible reaction.

Answer:

Irreversible reaction:

1. Products are not converted back to reactants.
2. Reaction stops completely and almost goes to completion.
3. It can be carried out in an open or closed vessel.
4. It takes place only in one direction. It is represented by $\rightarrow$
5. e.g. $\text{C}_{(s)} + \text{O}_{2(g)} \rightarrow \text{CO}_{2(g)}$

Reversible reaction:

1. Products are converted back to reactants.
2. Reaction appears to have stopped but does not undergo completion.
3. It is generally carried out in a closed vessel.
4. It takes place in both directions. It is represented by $\rightleftharpoons$
5. e.g. $\text{N}_{2(g)} + \text{O}_{2(g)} \rightleftharpoons 2\text{NO}_{(g)}$

Question K.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Write suitable conditions of concentration, temperature and pressure used during manufacture of ammonia by Haber process.

Answer:

- i. Concentration: Addition of H_2 or N_2 both favours forward reaction. This increases the yield of NH_3 .
- ii. Temperature: The formation NH_3 is exothermic. Hence, low temperature should favour the formation of NH_3 . However, at low temperatures, the rate of reaction is small. At high temperatures, the reaction occurs rapidly but decomposition of NH_3 occurs. Hence, optimum temperature of about 773 K is used.
- iii. Pressure: The forward reaction is favoured with high pressure as it proceeds with decrease in number of moles. At high pressure, the catalyst becomes inefficient. Therefore, optimum pressure needs to be used. The optimum pressure is about 250 atm.

Question L.

Relate the terms reversible reactions and dynamic equilibrium.


Answer:

- Reversible reactions are the reactions which do not go to completion and occur in both the directions simultaneously.
- If such a reaction is allowed to take place for a long time, so that the concentrations of the reactants and products do not vary with time, then the reaction will attain equilibrium.
- Since, both the forward and backward reactions continue to take place in opposite directions in the same speed, the equilibrium achieved is dynamic in nature.

Thus, if the reaction is not reversible then it cannot attain dynamic equilibrium.

Question M.

For the equilibrium.



<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

state the effect of

a. Addition of Ba^{2+} ion.

b. Removal of SO_4^{2-} ion

c. Addition of $\text{BaSO}_{4(s)}$

on the equilibrium.

Answer:

a. Addition of Ba^{2+} ion will favour the reverse reaction, (that is, equilibrium shifts from right to left). This increases the amount of BaSO_4 .

b. Removal of SO_4^{2-} ion will favour the forward reaction, (that is, equilibrium shifts from left to right). This decreases the amount of BaSO_4 .

c. Addition of $\text{BaSO}_{4(s)}$ will not affect the equilibrium as the equilibrium constant expression does not include pure solids.

3. Explain :

Question A.

Dynamic nature of chemical equilibrium with suitable example.

Answer:

Dynamic nature of chemical equilibrium:

i. Consider a chemical reaction: $\text{A} \rightleftharpoons \text{B}$.


$$K_c = [\text{B}]/[\text{A}]$$

At equilibrium, the ratio of concentration of the product to that of the concentration of the reactant is constant and this is equal to K_c .

ii. At this stage reaction takes place in both the directions with same speed although the reaction appears to have stopped. Thus, the chemical equilibrium is dynamic in nature. Dynamic means moving and at a microscopic level, the system is in motion.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

iii. For example, in the reaction between H_2 and I_2 to form HI, the colour of the reaction mixture becomes constant because the concentrations of H_2 , I_2 and HI become constant at equilibrium.




Thus, when equilibrium is reached, the reaction appears to have stopped. However, this is not the case. The reaction is still going on in the forward and backward direction but the rate of forward reaction is equal to the rate of backward reaction. Hence, chemical equilibrium is dynamic in nature and not static.

Question B.

Relation between K_c and K_p .

Answer:

Consider a general reversible reaction:



The equilibrium constant (K_p) in terms of partial pressure is given by equation:

$$K_p = \frac{(P_C)^c (P_D)^d}{(P_A)^a (P_B)^b} \dots\dots\dots(1)$$

For a mixture of ideal gases, the partial pressure of each component is directly proportional to its concentration at constant temperature.

For component A,

$$P_A V = n_A RT$$

$$P_A = n_A V \times RT$$

$n_A V$ is molar concentration of A in mol dm^{-3}

$$\therefore P_A = [A]RT \text{ where, } [A] = n_A V$$

Similarly, for other components, $P_B = [B]RT$, $P_C = [C]RT$, $P_D = [D]RT$

Now substituting equations for P_A , P_B , P_C , P_D in equation (1), we get

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

$$K_p = \frac{[C]^c (RT)^c [D]^d (RT)^d}{[A]^a (RT)^a [B]^b (RT)^b}$$
$$\therefore K_p = \frac{[C]^c [D]^d (RT)^{c+d}}{[A]^a [B]^b (RT)^{a+b}}$$
$$\therefore K_p = \frac{[C]^c [D]^d}{[A]^a [B]^b} \times (RT)^{(c+d)-(a+b)}$$
$$\therefore K_p = \frac{[C]^c [D]^d}{[A]^a [B]^b} \times (RT)^{\Delta n}$$
$$\text{But, } K_c = \frac{[C]^c [D]^d}{[A]^a [B]^b}$$
$$\therefore K_p = K_c (RT)^{\Delta n}$$

where Δn = (number of moles of gaseous products) – (number of moles of gaseous reactants) in the balanced chemical equation.

$$R = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

[Note: While calculating the value of K_p , pressure should be expressed in bar, because standard state of pressure is 1 bar. 1 pascal (Pa) = 1 N m⁻² and 1 bar = 10⁵ Pa]

Question C.

State and explain Le Chatelier's principle with reference to

1. change in temperature
2. change in concentration.

Answer:

Statement: When a system at equilibrium is subjected to a change in any of the factors determining the equilibrium conditions of a system, system will respond in such a way as to minimize the effect of change.

1. Change in temperature:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

- Consider the equilibrium reaction,
 $\text{PCl}_{5(g)} \rightleftharpoons \text{PCl}_{3(g)} + \text{Cl}_{2(g)} + 92.5 \text{ kJ}$
- The forward reaction is exothermic. According to Le Chatelier's principle an increase in temperature shifts the position of equilibrium to the left.
- The reverse reaction is endothermic. An endothermic reaction consumes heat. Therefore, the equilibrium must shift in the reverse direction to use up the added heat (heat energy converted to chemical energy).
- Thus, an increase in temperature favours formation of PCl_5 while a decrease in temperature favours decomposition of PCl_5 .

2. Change in concentration:

- Consider reversible reaction representing production of ammonia (NH_3).
 $\text{N}_{2(g)} + 3\text{H}_{2(g)} \rightleftharpoons 2\text{NH}_{3(g)} + \text{Heat}$
- According to Le Chatelier's principle, when H_2 or N_2 is added to equilibrium, the effect of addition of H_2 or N_2 is reduced by shifting the equilibrium from left to right so that the added N_2 or H_2 is consumed.
- The forward reaction occurs to a large extent than the reverse reaction until the new equilibrium is established. As a result, the yield of NH_3 is increases.
- In general, if the concentration of one of the species in equilibrium mixture is increased, the position of equilibrium shifts in the opposite so as to reduce the concentration of this species. However, the equilibrium constant remains unchanged.

Question D.

a. Reversible reaction

b. Rate of reaction

Answer:

a. Reversible reaction:



i. Reactions which do not go to completion and occur in both the directions simultaneously are called reversible reactions.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

ii. Reversible reactions proceed in both directions. The direction from reactants to products is the forward reaction, whereas the opposite reaction from products to reactants is called the reverse or backward reaction.

iii. A reversible reaction is denoted by drawing in between the reactants and product a double arrow, one pointing in the forward direction and other in the reverse direction ($\rightleftharpoons$ or $\rightleftharpoons$).

ii. At high temperature in an open container, the CO_2 gas formed will escape away. Therefore, it is not possible to obtain back



b. Rate of reaction:

Rate of a chemical reaction:

i. The rate of a chemical reaction can be determined by measuring the extent to which the concentration of a reactant decreases in the given time interval, or extent to which the concentration of a product increases in the given time interval.

ii. Mathematically, the rate of reaction is expressed as:

$$\text{Rate} = -d[\text{Reactant}]/dT = d[\text{Product}]/dT$$

where, $d[\text{reactant}]$ and $d[\text{product}]$ are the small decrease or increase in concentration during the small time interval dT .

Question E.

What is the effect of adding chloride on the position of the equilibrium ?



Answer:

Addition of Cl^- ion will favour the reverse reaction, (that is, equilibrium shift from right to left) This increases the amount of AgCl .

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Can you recall? (Textbook Page No. 174)

Question 1.

What are the types of the following changes?

Natural waterfall, spreading of smoke from burning incense stick, diffusion of fragrance of flowers.

Answer: Natural waterfall, spreading of smoke from burning incense stick and diffusion of fragrance of flowers are irreversible physical changes.

Try this. (Textbook Page No. 174)

Question 1.



Dissolve 4 g cobalt chloride in 40 mL water. It forms a reddish pink solution. Add 60 mL concentrated HCl to this. It will turn blue. Take 5 mL of this solution in a test tube and place it in a beaker containing ice water mixture. The colour of solution will become pink. Place the same test tube in a beaker containing water at 90 °C. The colour of the solution turns blue.

Answer:

Inference: The colour change of the solution from pink to blue is caused by the chemical reaction. On changing the temperature, the direction of the reaction reverses. This indicates that the chemical reaction is reversible. This activity is an example of a reversible chemical reaction.

The reaction can be written as:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>



Can you tell? (Textbook Page No. 174)

Question 1.

What does violet colour of the solution in the activity mentioned in Q.2 indicate?

Answer:

In the reaction, the reactant $\text{Co}(\text{H}_2\text{O})_6^{2+}$ is pink in colour and the product CoCl_4^{2-} is blue in colour. When the solution contains both the reactant and product, the resulting solution will appear violet. This indicates that the reaction has attained equilibrium (that is, the reaction proceeds in both the direction with equal rates and is a reversible reaction).

(Textbook Page No. 174)

Question 1.

Calcium carbonate when heated strongly, decomposes to form calcium oxide and carbon dioxide.


i. If this reaction is carried out in a closed container, what will we observe?

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

ii. Consider this reaction occurring in an open system or container, what will happen? Can we obtain back calcium carbonate?

Answer:

At high temperature in a closed container, we will find that after certain time, some calcium carbonate is present. If we continue the experiment over a longer period of time at the same temperature, the concentrations of calcium carbonate, calcium oxide and carbon dioxide remain unchanged. The reaction thus appears to have stopped and the system has attained the equilibrium. Actually, the reaction does not stop but proceeds in both the directions with equal rates. In other words, calcium carbonate decomposes to give calcium oxide and carbon dioxide at a particular rate. Exactly at the same rate the calcium oxide and carbon dioxide recombine and form calcium carbonate. Thus, in closed container, reversible reaction occurs.



ii. At high temperature in an open container, the CO_2 gas formed will escape away. Therefore, it is not possible to obtain back calcium carbonate. Thus, in an open container, irreversible reaction occurs.



Internet my friend (Textbook Page No. 175)

Question 1.

i. Equilibrium existing in the formation of oxyhaemoglobin in human body

ii. Refrigeration system in equilibrium


Answer:

i. Equilibrium existing in the formation of oxyhaemoglobin in human body:

Oxygen is transported in the body with the assistance of red blood cells. The red blood cells contain a pigment called haemoglobin. Each haemoglobin molecule binds four oxygen molecules to form oxyhaemoglobin. Thus, the oxygen molecules are carried to individual cells in the body tissue where they are released.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

The binding of oxygen to haemoglobin is a reversible reaction.



When the oxygen concentration is high (in the lungs), haemoglobin and oxygen combine to form oxyhaemoglobin and the reaction achieves equilibrium. But, when the oxygen concentration is low (in the body tissue), the reverse reaction occurs, that is, oxyhaemoglobin dissociates to haemoglobin and oxygen.


Thus, an equilibrium exists in the formation of oxyhaemoglobin in the human body.

ii. Refrigeration system in equilibrium:

a. Refrigeration system works on the principle of thermal equilibrium i.e., when a cold body comes in contact with a hot body then the heat flows from hot body to cold body until both the bodies attain the same temperature.

b. In the same way, a liquid (called as refrigerant) passes through the various compartments in the refrigerator and eventually lowers the temperature inside the refrigerator. This cycle is briefly described below:

Refrigerant flows through the compressor, which raises the pressure of the refrigerant. Next, the refrigerant flows through the condenser, where it condenses from vapor form to liquid form, giving off heat in the process. The heat given off is what makes the condenser “hot to the touch.” After the condenser, the refrigerant goes through the expansion valve, where it experiences a pressure drop. Finally, the refrigerant goes to the evaporator. The refrigerant draws heat from the evaporator which causes the refrigerant to vaporize. The evaporator draws heat from the region that is to be cooled. The vaporized refrigerant goes back to the compressor to restart the cycle. In each of the heat transfer process, equilibrium is achieved (that is, heat given off is equivalent to the cooling achieved.)



[Note: Students are expected to collect additional information about equilibrium existing in the formation of oxyhaemoglobin in human body' and 'refrigeration system in equilibrium on their own.]

Try this. (Textbook Page No. 176)

Question 1.


- i. Place some iodine crystals in a closed vessel. Observe the change in colour intensity in it.**
- ii. What do you see in the flask after some time?**

Answer:

- i. The vessel gets slowly filled up with violet coloured vapour of iodine. After a certain time, the intensity of violet colour becomes stable.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

ii. After sometime, both solid iodine and iodine vapour are present in the closed vessel. Iodine crystals will be seen deposited near the mouth of the flask and violet coloured vapour will be filled in the entire flask. It means solid iodine sublimes to give iodine vapour and the iodine vapour condenses to form solid iodine. The stable intensity of the colour indicates a state of equilibrium between solid and vapour iodine.



Try this. (Textbook Page No. 176)

Question 1.

- i. Dissolve a given amount of sugar in minimum amount of water at room temperature.**
- ii. Increase the temperature and dissolve more amount of sugar in the same amount of water to make a thick sugar syrup solution.**
- iii. Cool the syrup to the room temperature.**


Answer:

Observation: Sugar crystals separate out.

Inference: The sugar syrup solution prepared is a saturated solution. Therefore, additional amount of sugar cannot be dissolved in it at room temperature.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

In a saturated solution, there exists dynamic equilibrium between the solute molecules in the solid state and in dissolved state.



The rate of dissolution of sugar = The rate of crystallization of sugar.

However, when it is heated, additional amount of sugar can be dissolved in it. But when such a thick sugar syrup is cooled again to room temperature, sugar crystals separate out.

Do you know? (Textbook Page No. 177)

Question 1.

What is a saturated solution?


Answer:

A saturated solution is the solution when additional solute cannot be dissolved in it at the given temperature. The concentration of solute in a saturated solution depends on temperature.

Observe and discuss. (Textbook Page No. 177)

Question 1.

Colourless N_2O_4 taken in a closed flask is converted to NO_2 (a reddish brown gas).




Answer:

Observation: Initially, the colourless gas (N₂O₄) turns to reddish brown (NO₂) gas. After sometime, the colour becomes lighter indicating the formation of N₂O₄ from NO₂.

Inference: This indicates that the reaction is reversible. In such reaction, the reactants combine to form the products and the products combine to give the reactants. As soon as the forward reaction produces any NO₂, the reverse reaction begins and NO₂, starts combining back to N₂O₄. At equilibrium, the concentrations of N₂O₄ and NO₂ remain unchanged and do not vary with time, because the rate of formation of NO₂ is equal to the rate of formation of N₂O₄.


<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>



[Note: For any reversible reaction in a closed system whenever the opposing reactions (forward and reaction) are occurring at different rates, the forward reaction will gradually become slower and the reverse reaction will become faster. Finally, the rates become equal and equilibrium is established.]


Discuss (Textbook Page No. 177)

i. Consider the following dissociation reaction:



The reaction is carried out in a closed vessel starting with hydrogen iodide.

ii. Now, let us start with hydrogen and iodine vapour in a closed container at a certain temperature.



Answer:

i. Starting with hydrogen iodide:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Observations:

- a. At first, there is an increase in the intensity of violet colour.
- b. After certain time, the increase in the intensity of violet colour stops.
- c. When contents in a closed vessel are analyzed at this stage, it is observed that reaction mixture contains the hydrogen iodide, hydrogen and iodine with their concentrations being constant over time.

Inference:

The rate of decomposition of HI becomes equal to the rate of combination of H_2 and I_2 . At equilibrium, no net change is observed and both reactions continue to occur at equal rates.

Thus, the reaction represents chemical equilibrium.

ii. Starting with hydrogen and iodine:

Observations:

- a. At first, there is a decrease in the intensity of violet colour.
- b. After certain time, the decrease in the intensity of violet colour stops.
- c. When contents in a closed vessel are analyzed at this stage, it is observed that reaction mixture contains hydrogen, iodine and hydrogen iodide with their concentrations being constant over time.

Inference:

The rate of combination of H_2 and I_2 becomes equal to the rate of decomposition of HI. The reaction attains chemical equilibrium.

Can you recall? (Textbook Page No. 180)

Question 1.

Write ideal gas equation with significance of each term involved in it.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Answer:

Ideal gas equation is $PV = nRT$.

where, P = Pressure of the gas

V = Volume of the gas

n = Number of moles of the gas

R = universal gas constant

T = Absolute temperature of the gas

Just think. (Textbook page no. 181)


Question 1.

Two processes, which are taking place in opposite directions are in equilibrium. How to write equilibrium constant expressions for heterogeneous equilibrium?

Answer:

Equilibrium in a system having more than one phase is called heterogeneous equilibrium.

If ethanol is placed in a conical flask, liquid-vapour equilibrium is established.



For a given temperature,

$$K_c = [C_2H_5OH_{(g)}][C_2H_5OH_{(l)}]$$

$$\text{But } [C_2H_5OH_{(l)}] = 1$$

$$\therefore K_c = [C_2H_5OH_{(g)}]$$

Thus, at any given temperature, density is constant irrespective of the amount of liquid, and the term in the denominator is also constant.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

ii. similarly, consider $I_{2(g)} \rightleftharpoons I_{2(g)}$

$$K_c = [I_{2(g)}]$$

iii. Thus, the expression for equilibrium constant does not contain the concentration of pure solids and pure liquids. That is because for any pure liquid and solid, the concentration is simply its density and this will not change no matter how much solid or liquid is used. Hence, the expression for heterogeneous equilibrium only uses the concentration of gases and dissolved substances (aq.). Solids are pure substances with unchanging concentrations and thus equilibria including solids are simplified.

Can you tell? (Textbook Page No. 183)

Question 1.

Comment on the extent to which the forward reaction will proceed, from the magnitude of the equilibrium constant for the following reactions:

i. $H_{2(g)} + I_{2(g)} \rightleftharpoons 2HI_{(g)}$, $K_c = 20$ at 550 K

ii. $H_{2(g)} + Cl_{2(g)} \rightleftharpoons 2HCl_{(g)}$, $K_c = 10^{18}$ at 550 K

Answer:

i. For the reaction, $K_c = 20$ at 550 K

If the value of K_c is the range of 10^{-3} to 10^3 , the forward and reverse proceed to equal extents.

Hence, the given reaction will form appreciable concentrations of both reactants and the product at equilibrium.

ii. For the reaction, $K_c = 10^{18}$ at 550 K

If the value of $K_c \gg 10^3$, forward reaction is favoured.

Hence, the given reaction will proceed in the forward direction and will nearly go to completion.

Use your brain power (Textbook Page No. 183)

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Question 1.

The value of K_c for the dissociation reaction:

$H_{2(g)} \rightleftharpoons 2H_{(g)}$ is 1.2×10^{-42} at 500 K.

Does the equilibrium mixture contain mainly hydrogen molecules or hydrogen atoms?

Answer:

When the value of K_c is very low (that is, $K_c < 10^{-3}$), then at equilibrium, only a small fraction of the reactants is converted into products.

For the given reaction, $K_c \ll 10^3$ at 500 K.

Hence, the equilibrium mixture contains mainly hydrogen molecules.

Internet my friend (Textbook Page No. 183)

Question 1.

Collect information about chemical equilibrium.

Answer:

<https://www.chemguide.co.uk/physical/equilibria/introduction.html>

[Note: Students can use the above link as reference and collect information about chemical equilibrium.]

Can you tell? (Textbook Page No. 188)

- i. If NH_3 is added to the equilibrium system (Haber process), in which direction will the equilibrium shift to consume added NH_3 to reduce the effect of stress?
- ii. In this process, out of the reactions (reverse and forward reaction), which reaction will occur to a greater extent?
- iii. What will be the effect on yield of NH_3 ?

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

Answer:

- i. If NH_3 is added to the equilibrium system, the equilibrium will shift from right to left to consume added NH_3 to reduce the effect of stress.
- ii. If NH_3 is added to the equilibrium system, then reverse reaction will occur to greater extent.
- iii. If NH_3 is added to the equilibrium system, the equilibrium will shift in reverse direction and the yield of NH_3 will decrease.

Internet my friend (Textbook Page No. 188)

- i. Collect information about Haber process in chemical equilibrium.
- ii. Youtube.Freescienceslessons: The Haber process

Answer:

- i. <https://www.chemguide.co.uk/physical/equilibria/haber.html>

[Note: Students can use the above link as reference and collect information about chemical equilibrium involved in Haber process.]

- ii. Students are expected to refer 'The Haber process' on YouTube channel 'Freescienceslessons'



Maharashtra Board Solutions

Class 11 Chemistry

- Chapter 1- Some Basic Concepts of Chemistry
- Chapter 2- Introduction to Analytical Chemistry
- Chapter 3- Basic Analytical Techniques
- Chapter 4- Structure of Atom
- Chapter 5- Chemical Bonding
- Chapter 6- Redox Reactions
- Chapter 7- Modern Periodic Table
- Chapter 8- Elements of Group 1 and 2
- Chapter 9- Elements of Group 13, 14 and 15
- Chapter 10- States of Matter
- Chapter 11- Adsorption and Colloids
- Chapter 12- Chemical Equilibrium
- Chapter 13- Nuclear Chemistry and Radioactivity
- Chapter 14- Basic Principles of Organic Chemistry
- Chapter 15- Hydrocarbons
- Chapter 16- Chemistry in Everyday Life

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

About About Maharashtra State Board (MSBSHSE)

The Maharashtra State Board of Secondary and Higher Secondary Education or MSBSHSE (Marathi: महाराष्ट्र राज्य माध्यमिक आणि उच्च माध्यमिक शिक्षण मंडळ), is an **autonomous and statutory body established in 1965**. The board was amended in the year 1977 under the provisions of the Maharashtra Act No. 41 of 1965.

The Maharashtra State Board of Secondary & Higher Secondary Education (MSBSHSE), Pune is an independent body of the Maharashtra Government. There are more than 1.4 million students that appear in the examination every year. The Maha State Board conducts the board examination twice a year. This board conducts the examination for SSC and HSC.

The Maharashtra government established the Maharashtra State Bureau of Textbook Production and Curriculum Research, also commonly referred to as Ebalbharati, in 1967 to take up the responsibility of providing quality textbooks to students from all classes studying under the Maharashtra State Board. MSBHSE prepares and updates the curriculum to provide holistic development for students. It is designed to tackle the difficulty in understanding the concepts with simple language with simple illustrations. Every year around 10 lakh students are enrolled in schools that are affiliated with the Maharashtra State Board.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

FAQs

Where do I get the Maharashtra State Board Books PDF For free download?

You can download the Maharashtra State Board Books from the eBalbharti official website, i.e. cart.ebalbharati.in or from this article.

Add image

How to Download Maharashtra State Board Books?

Students can get the Maharashtra Books for primary, secondary, and senior secondary classes from here. You can view or download the Maharashtra State Board Books from this page or from the official website for free of cost. Students can follow the detailed steps below to visit the official website and download the e-books for all subjects or a specific subject in different mediums.

Step 1: Visit the official website ebalbharati.in

Step 2: On the top of the screen, select "Download PDF textbooks"

Step 3: From the "Classes" section, select your class.

Step 4: From "Medium", select the medium suitable to you.

Step 5: All Maharashtra board books for class 11th will now be displayed on the right side.

Step 6: Click on the "Download" option to download the PDF book.

Who developed the Maharashtra State board books?

As of now, the MSCERT and Balbharti are responsible for the syllabus and textbooks of Classes 1 to 8, while Classes 9 and 10 are under the Maharashtra State Board of Secondary and Higher Secondary Education (MSBSHSE).

How many state boards are there in Maharashtra?

The Maharashtra State Board of Secondary & Higher Secondary Education, conducts the HSC and SSC Examinations in the state of Maharashtra through its nine Divisional Boards located at Pune, Mumbai, Aurangabad, Nasik, Kolhapur, Amravati, Latur, Nagpur and Ratnagiri.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>

About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to **info@indcareer.com**

Postal Address

IndCareer.com
52, Shilpa Nagar,
Somalwada
Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888

Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-11-adsorption-and-colloids/>