


ML Aggarwal Solutions for Class 9 Maths

Chapter 18- Trigonometric Ratios and Standard Angles



www.indcareer.com/schools



ML Agrawal Solutions

CLASS 9 MATHS

Chapter 18: Trigonometric
Ratios and Standard
Angles

For any clarifications or questions you can write to info@indcareer.com

Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888, **Website:** <https://www.indcareer.com> Class 8: Biology
Chapter 6 solutions. Complete Class 8 Biology Chapter 6 Notes.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

ML Aggarwal Solutions for Class 9 Maths

Chapter 18- Trigonometric Ratios and Standard Angles

Class 9: Maths Chapter 18 solutions. Complete Class 9 Maths Chapter 18 Notes.

ML Aggarwal Solutions for Class 9 Maths Chapter 18- Trigonometric Ratios and Standard Angles

ML Aggarwal 9th Maths Chapter 18, Class 9 Maths Chapter 18 solutions

Exercise 18.1

1. Find the values of

(i) $7 \sin 30^\circ \cos 60^\circ$

(ii) $3 \sin^2 45^\circ + 2 \cos^2 60^\circ$

(iii) $\cos^2 45^\circ + \sin^2 60^\circ + \sin^2 30^\circ$

(iv) $\cos 90^\circ + \cos^2 45^\circ \sin 30^\circ \tan 45^\circ$.

Solution:

(i) $7 \sin 30^\circ \cos 60^\circ$

Substituting the values

$$= 7 \times \frac{1}{2} \times \frac{1}{2}$$

$$= (7 \times 1 \times 1)/(2 \times 2)$$

$$= 7/4$$

(ii) $3 \sin^2 45^\circ + 2 \cos^2 60^\circ$

Substituting the values

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 3 \times (1/\sqrt{2})^2 + 2 \times (1/2)^2$$

By further calculation

$$= 3 \times \frac{1}{2} + 2 \times \frac{1}{4}$$

$$= \frac{3}{2} + \frac{1}{2}$$

So we get

$$= (3 + 1)/2$$

$$= 4/2$$

$$= 2$$

$$(iii) \cos^2 45^\circ + \sin^2 60^\circ + \sin^2 30^\circ$$

Substituting the values

$$= (1/\sqrt{2})^2 + (\sqrt{3}/2)^2 + (1/2)^2$$

By further calculation

$$= \frac{1}{2} + \frac{3}{4} + \frac{1}{4}$$

Taking LCM

$$= (2 + 3 + 1)/4$$

$$= 6/4$$

$$= 3/2$$

$$(iv) \cos 90^\circ + \cos^2 45^\circ \sin 30^\circ \tan 45^\circ$$

Substituting the values

$$= 0 + (1/\sqrt{2})^2 \times \frac{1}{2} \times 1$$

By further calculation

$$= \frac{1}{2} \times \frac{1}{2} \times 1$$

$$= \frac{1}{4}$$

2. Find the values of

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(i) \frac{\sin^2 45^\circ + \cos^2 45^\circ}{\tan^2 60^\circ}$$

$$(ii) \frac{\sin 30^\circ - \sin 90^\circ + 2\cos 0^\circ}{\tan 30^\circ \times \tan 60^\circ}$$

$$(iii) \frac{4}{3} \tan^2 30^\circ + \sin^2 60^\circ - 3\cos^2 60^\circ + \frac{3}{4} \tan^2 60^\circ - 2\tan^2 45^\circ.$$

Solution:

$$(ii) \frac{\sin 30^\circ - \sin 90^\circ + 2\cos 0^\circ}{\tan 30^\circ \times \tan 60^\circ}$$

Substituting the values

$$= \frac{\frac{1}{2} - 1 + 2 \times 1}{\frac{1}{\sqrt{3}} \times \sqrt{3}}$$

$$(i) \frac{\sin^2 45^\circ + \cos^2 45^\circ}{\tan^2 60^\circ} = \frac{\frac{1}{2} - 1 + 2}{1}$$

Substituting the values So we get

$$= \frac{(\frac{1}{\sqrt{2}})^2 + (\frac{1}{\sqrt{2}})^2}{\sqrt{3}^2} = \frac{1}{2} - 1 + 2$$

$$\text{By further calculation} = \frac{1}{2} + 1$$

$$= \frac{\frac{1}{2} + \frac{1}{2}}{3} = \frac{1 + 2}{2}$$

$$= \frac{1}{3} = \frac{3}{2}$$

$$(iii) \frac{4}{3} \tan^2 30^\circ + \sin^2 60^\circ - 3 \cos^2 60^\circ + \frac{3}{4} \tan^2 60^\circ - 2 \tan^2 45^\circ$$

Substituting the values

$$= \frac{4}{3} (1/\sqrt{3})^2 + (\sqrt{3}/2)^2 - 3 (1/2)^2 + \frac{3}{4} \times (\sqrt{3})^2 - 2 \times 1^2$$

By further calculation

$$= \frac{4}{3} \times \frac{1}{3} + \frac{3}{4} - 3 \times \frac{1}{4} + \frac{3}{4} \times 3 - 2 \times 1$$

$$= \frac{4}{9} + \frac{3}{4} - \frac{3}{4} + \frac{9}{4} - 2$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

So we get

$$= 4/9 + 9/4 - 2$$

Taking LCM

$$= (16 + 81 - 72)/36$$

$$= (97 - 72)/36$$

$$= 25/36$$

3. Find the values of

$$(i) \frac{\sin 60^\circ}{\cos^2 45^\circ} - 3 \tan 30^\circ + 5 \cos 90^\circ$$

$$(ii) 2\sqrt{2} \cos 45^\circ \cos 60^\circ + 2\sqrt{3} \sin 30^\circ \tan 60^\circ - \cos 0^\circ$$

$$(iii) \frac{4}{5} \tan^2 60^\circ - \frac{2}{\sin^2 30^\circ} - \frac{3}{4} \tan^2 30^\circ.$$

Solution:

$$(i) \frac{\sin 60^\circ}{\cos^2 45^\circ} - 3 \tan 30^\circ + 5 \cos 90^\circ$$

Substituting the values

$$\begin{aligned} &= \frac{\left(\frac{\sqrt{3}}{2}\right)}{\left(\frac{1}{\sqrt{2}}\right)^2} - 3 \times \frac{1}{\sqrt{3}} + 5 \times 0 \\ &= \frac{\frac{\sqrt{3}}{2}}{\frac{1}{2}} - \sqrt{3} - 0 \end{aligned}$$

So we get

$$= \frac{\sqrt{3}}{2} \times \frac{2}{1} - \sqrt{3}$$

$$= \sqrt{3} - \sqrt{3}$$

$$= 0$$

$$(ii) 2\sqrt{2} \cos 45^\circ \cos 60^\circ + 2\sqrt{3} \sin 30^\circ \tan 60^\circ - \cos 0^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

Substituting the values

$$= 2\sqrt{2} \times 1/\sqrt{2} \times \frac{1}{2} + 2\sqrt{3} \times \frac{1}{2} \times \sqrt{3} - 1$$

By further calculation

$$= 2 \times 1/1 \times 1/2 + 2 \times 3 \times \frac{1}{2} - 1$$

$$= 1 + 3 - 1$$

$$= 3$$

$$(iii) \frac{4}{5} \tan^2 60^\circ + \frac{2}{\sin^2 30^\circ} - \frac{3}{4} \tan^2 30^\circ$$

Substituting the values

$$= \frac{4}{5} \times (\sqrt{3})^2 - \frac{2}{(\frac{1}{2})^2} - \frac{3}{4} \left(\frac{1}{\sqrt{3}}\right)^2$$

By further calculation

$$= \frac{4}{5} \times 3 - \frac{2}{\frac{1}{4}} - \frac{3}{4} \times \frac{1}{3}$$

So we get

$$= \frac{12}{5} - \frac{2 \times 4}{1} - \frac{1}{4}$$

$$= \frac{12}{5} - \frac{8}{1} - \frac{1}{4}$$

Taking LCM

$$= \frac{48 - 160 - 5}{20}$$

$$= \frac{43 - 160}{20}$$

$$= \frac{-117}{20}$$

$$= -5\frac{17}{20}$$

4. Prove that

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(i) $\cos^2 30^\circ + \sin 30^\circ + \tan^2 45^\circ = 2 \frac{1}{4}$

(ii) $4 (\sin^4 30^\circ + \cos^4 60^\circ) - 3 (\cos^2 45^\circ - \sin^2 90^\circ) = 2$

(iii) $\cos 60^\circ = \cos^2 30^\circ - \sin^2 30^\circ$.

Solution:

(i) $\cos^2 30^\circ + \sin 30^\circ + \tan^2 45^\circ = 2 \frac{1}{4}$

Consider

$$\text{LHS} = \cos^2 30^\circ + \sin 30^\circ + \tan^2 45^\circ$$

Substituting the values

$$= (\sqrt{3}/2)^2 + \frac{1}{2} + 1^2$$

By further calculation

$$= \frac{3}{4} + \frac{1}{2} + 1$$

Taking LCM

$$= (3 + 2 + 4)/4$$

$$= 9/4$$

$$= 2 \frac{1}{4}$$

$$= \text{RHS}$$

Therefore, LHS = RHS.

(ii) $4 (\sin^4 30^\circ + \cos^4 60^\circ) - 3 (\cos^2 45^\circ - \sin^2 90^\circ) = 2$

Consider

$$\text{LHS} = 4 (\sin^4 30^\circ + \cos^4 60^\circ) - 3 (\cos^2 45^\circ - \sin^2 90^\circ)$$

Substituting the values

$$= 4[(\frac{1}{2})^4 + (\frac{1}{2})^4] - 3 [(1/\sqrt{2})^2 - 1^2]$$

It can be written as

$$= 4[\frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} + \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2}] - 3 [\frac{1}{2} - 1]$$

By further calculation

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 4 [1/16 + 1/16] - 3 (- 1/2)$$

$$= 4[(1 + 1)/16] + 3/2$$

So we get

$$= (4 \times 3)/16 + 3/2$$

$$= 8/16 + 3/2$$

$$= 1/2 + 3/2$$

$$= (1 + 3)/2$$

$$= 4/2$$

$$= 2$$

$$= \text{RHS}$$

Therefore, LHS = RHS.

$$\text{(iii) } \cos 60^\circ = \cos^2 30^\circ - \sin^2 30^\circ$$

Consider

$$\text{LHS} = \cos 60^\circ = 1/2$$

$$\text{RHS} = \cos^2 30^\circ - \sin^2 30^\circ$$

Substituting the values

$$= (\sqrt{3}/2)^2 + (1/2)^2$$

By further calculation

$$= 3/4 - 1/4$$

$$= (3 - 1)/4$$

$$= 2/4$$

$$= 1/2$$

$$= \text{RHS}$$

Therefore, LHS = RHS.

5. (i) If $x = 30^\circ$, verify that $\tan 2x = 2\tan x / (1 - \tan^2 x)$.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(ii) If $x = 15^\circ$, verify that $4 \sin 2x \cos 4x \sin 6x = 1$.

Solution:

(i) It is given that

$$x = 30^\circ$$

Consider LHS = $\tan 2x$

Substituting the value of x

$$= \tan 60^\circ$$

$$= \sqrt{3}$$

$$RHS = \frac{2 \tan x}{1 - \tan^2 x}$$

Substituting the value of x

$$= \frac{2 \tan 30^\circ}{1 - \tan^2 30^\circ}$$

By further calculation

$$= \frac{2 \times \frac{1}{\sqrt{3}}}{1 - \left(\frac{1}{\sqrt{3}}\right)^2}$$

So we get

$$\begin{aligned} &= \frac{\frac{2}{\sqrt{3}}}{1 - \frac{1}{3}} \\ &= \frac{\frac{2}{\sqrt{3}}}{\frac{3-1}{3}} \\ &= \frac{2}{\sqrt{3}} \times \frac{3}{2} \\ &= \frac{3}{\sqrt{3}} \\ &= \sqrt{3} \end{aligned}$$

Therefore, LHS = RHS.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(ii) It is given that

$$x = 15^{\circ}$$

$$2x = 15 \times 2 = 30^{\circ}$$

$$4x = 15 \times 4 = 60^{\circ}$$

$$6x = 15 \times 6 = 90^{\circ}$$

Here

$$\text{LHS} = 4 \sin 2x \cos 4x \sin 6x$$

It can be written as

$$= 4 \sin 30^{\circ} \cos 60^{\circ} \sin 90^{\circ}$$

So we get

$$= 4 \times \frac{1}{2} \times \frac{1}{2} \times 1$$

$$= 1$$

$$= \text{RHS}$$

Therefore, LHS = RHS.

6. Find the values of

$$(i) \sqrt{\frac{1 - \cos^2 30^{\circ}}{1 - \sin^2 30^{\circ}}}$$

$$(ii) \frac{\sin 45^{\circ} \cos 45^{\circ} \cos 60^{\circ}}{\sin 60^{\circ} \cos 30^{\circ} \tan 45^{\circ}}.$$

Solution:

$$(i) \sqrt{\frac{1 - \cos^2 30^\circ}{1 - \sin^2 30^\circ}}$$

Substituting the values

$$= \sqrt{\frac{1 - (\frac{\sqrt{3}}{2})^2}{1 - (\frac{1}{2})^2}}$$

By further calculation

$$= \sqrt{\frac{1 - \frac{3}{4}}{1 - \frac{1}{4}}}$$

Taking LCM

$$= \sqrt{\frac{\frac{4-3}{4}}{\frac{4-1}{4}}}$$

$$= \sqrt{\frac{\frac{1}{4}}{\frac{3}{4}}} = \sqrt{\frac{1}{4} \times \frac{4}{3}}$$

So we get

$$= \sqrt{\frac{1}{3}}$$

$$= \frac{1}{\sqrt{3}}$$

$$(ii) \frac{\sin 45^\circ \cos 45^\circ \cos 60^\circ}{\sin 60^\circ \cos 30^\circ \tan 45^\circ}$$

Substituting the values

$$= \frac{\frac{1}{\sqrt{2}} \times \frac{1}{\sqrt{2}} \times \frac{1}{2}}{\frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2} \times 1}$$

By further calculation

$$= \frac{\frac{1}{2} \times \frac{1}{2}}{\frac{3}{4} \times 1}$$

$$= \frac{\frac{1}{4}}{\frac{3}{4}}$$

$$= \frac{1}{3}$$

7. If $\theta = 30^\circ$, verify that

(i) $\sin 2\theta = 2 \sin \theta \cos \theta$

(ii) $\cos 2\theta = 2 \cos^2 \theta - 1$

(iii) $\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$

(iv) $\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$.

Solution:

It is given that $\theta = 30^\circ$

(i) $\sin 2\theta = 2 \sin \theta \cos \theta$

Consider

LHS = $\sin 2\theta$

Substituting the value of θ

= $\sin 2 \times 30^\circ$

= $\sin 60^\circ$

= $\sqrt{3}/2$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\text{RHS} = 2 \sin \theta \cos \theta$$

Substituting the value of θ

$$= 2 \sin 30^\circ \cos 30^\circ$$

So we get

$$= 2 \times \frac{1}{2} \times \frac{\sqrt{3}}{2}$$

$$= 1 \times \frac{\sqrt{3}}{2}$$

$$= \frac{\sqrt{3}}{2}$$

Therefore, LHS = RHS.

$$(ii) \cos 2\theta = 2 \cos^2 \theta - 1$$

Consider

$$\text{LHS} = \cos 2\theta$$

Substituting the value of θ

$$= \cos 2 \times 30^\circ$$

$$= \cos 60^\circ$$

$$= \frac{1}{2}$$

$$\text{RHS} = 2 \cos^2 \theta - 1$$

Substituting the value of θ

$$= 2 \cos^2 30^\circ - 1$$

So we get

$$= 2 \left(\frac{\sqrt{3}}{2} \right)^2 - 1$$

$$= 2 \times \frac{3}{4} - 1$$

$$= \frac{3}{2} - 1$$

$$= \frac{(3 - 2)}{2}$$

$$= \frac{1}{2}$$

Therefore, LHS = RHS.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(iii) \sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$$

Consider

$$\text{LHS} = \sin 3\theta$$

Substituting the value of θ

$$= \sin 3 \times 30^\circ$$

$$= \sin 90^\circ$$

$$= 1$$

$$\text{RHS} = 3 \sin \theta - 4 \sin^3 \theta$$

Substituting the value of θ

$$= 3 \sin 30^\circ - 4 \sin^3 30^\circ$$

So we get

$$= 3 \times \frac{1}{2} - 4 \times \left(\frac{1}{2}\right)^3$$

$$= \frac{3}{2} - 4 \times \frac{1}{8}$$

$$= \frac{3}{2} - \frac{1}{2}$$

Taking LCM

$$= \frac{(3 - 1)}{2}$$

$$= \frac{2}{2}$$

$$= 1$$

Therefore, LHS = RHS.

$$(iv) \cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$$

Consider

$$\text{LHS} = \cos 3\theta$$

Substituting the value of θ

$$= \cos 3 \times 30^\circ$$

$$= \cos 90^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 0$$

$$\text{RHS} = 4 \cos^3 \theta - 3 \cos \theta$$

Substituting the value of θ

$$= 4 \cos^3 30^\circ - 3 \cos 30^\circ$$

So we get

$$= 4 \times (\sqrt{3}/2)^3 - 3 \times (\sqrt{3}/2)$$

By further calculation

$$= 4 \times 3\sqrt{3}/8 - 3\sqrt{3}/2$$

$$= 3\sqrt{3}/2 - 3\sqrt{3}/2$$

$$= 0$$

Therefore, LHS = RHS.

8. If $\theta = 30^\circ$, find the ratio $2 \sin \theta : \sin 2\theta$.

Solution:

It is given that $\theta = 30^\circ$

We know that

$$2 \sin \theta : \sin 2\theta = 2 \sin 30^\circ : \sin 2 \times 30^\circ$$

So we get

$$= 2 \sin 30^\circ : \sin 60^\circ$$

$$= 2 \sin 30^\circ / \sin 60^\circ$$

Substituting the values

$$= \frac{2 \times \frac{1}{2}}{\frac{\sqrt{3}}{2}}$$

By further simplification

$$\begin{aligned} &= \frac{1}{\frac{\sqrt{3}}{2}} \\ &= \frac{1 \times 2}{\sqrt{3}} \\ &= \frac{2}{\sqrt{3}} \end{aligned}$$

$$= 2 : \sqrt{3}$$

Therefore, $2 \sin \theta : \sin 2\theta = 2 : \sqrt{3}$.

9. By means of an example, show that $\sin (A + B) \neq \sin A + \sin B$.

Solution:

Consider $A = 30^\circ$ and $B = 60^\circ$

$$\text{LHS} = \sin (A + B)$$

Substituting the values of A and B

$$= \sin (30^\circ + 60^\circ)$$

$$= \sin 90^\circ$$

$$= 1$$

$$\text{RHS} = \sin A + \sin B$$

Substituting the values

$$= \sin 30^\circ + \sin 60^\circ$$

So we get

$$= \frac{1}{2} + \frac{\sqrt{3}}{2}$$

$$= \frac{(1 + \sqrt{3})}{2}$$

Therefore, $\text{LHS} \neq \text{RHS}$ i.e. $\sin (A + B) \neq \sin A + \sin B$.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

10. If $A = 60^\circ$ and $B = 30^\circ$, verify that

(i) $\sin (A + B) = \sin A \cos B + \cos A \sin B$

(ii) $\cos (A + B) = \cos A \cos B - \sin A \sin B$

(iii) $\sin (A - B) = \sin A \cos B - \cos A \sin B$

(iv) $\tan (A - B) = (\tan A - \tan B) / (1 + \tan A \tan B)$.

Solution:

It is given that $A = 60^\circ$ and $B = 30^\circ$

(i) $\sin (A + B) = \sin A \cos B + \cos A \sin B$

Here

$$\text{LHS} = \sin (A + B)$$

Substituting the values of A and B

$$= \sin (60^\circ + 30^\circ)$$

$$= \sin 90^\circ$$

$$= 1$$

$$\text{RHS} = \sin A \cos B + \cos A \sin B$$

Substituting the values of A and B

$$= \sin 60^\circ \cos 30^\circ + \cos 60^\circ \sin 30^\circ$$

So we get

$$= \sqrt{3}/2 \times \sqrt{3}/2 + 1/2 \times 1/2$$

By further calculation

$$= 3/4 + 1/4$$

$$= 4/4$$

$$= 1$$

Therefore, LHS = RHS.

(ii) $\cos (A + B) = \cos A \cos B - \sin A \sin B$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

Here

$$\text{LHS} = \cos (A + B)$$

Substituting the value of A and B

$$= \cos (60^{\circ} + 30^{\circ})$$

$$= \cos 90^{\circ}$$

$$= 0$$

$$\text{RHS} = \cos A \cos B - \sin A \sin B$$

Substituting the value of A and B

$$= \cos 60^{\circ} \cos 30^{\circ} - \sin 60^{\circ} \sin 30^{\circ}$$

So we get

$$= \frac{1}{2} \times \frac{\sqrt{3}}{2} - \frac{\sqrt{3}}{2} \times \frac{1}{2}$$

$$= \frac{\sqrt{3}}{4} - \frac{\sqrt{3}}{4}$$

$$= 0$$

Therefore, LHS = RHS.

$$\text{(iii) } \sin (A - B) = \sin A \cos B - \cos A \sin B$$

Here

$$\text{LHS} = \sin (A - B)$$

Substituting the values of A and B

$$= \sin (60^{\circ} - 30^{\circ})$$

$$= \sin 30^{\circ}$$

$$= \frac{1}{2}$$

$$\text{RHS} = \sin A \cos B - \cos A \sin B$$

Substituting the values of A and B

$$= \sin 60^{\circ} \cos 30^{\circ} - \cos 60^{\circ} \sin 30^{\circ}$$

So we get

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \sqrt{3}/2 \times \sqrt{3}/2 - 1/2 \times 1/2$$

$$= 3/4 - 1/4$$

$$= (3 - 1)/4$$

$$= 2/4$$

$$= 1/2$$

Therefore, LHS = RHS.

$$(iv) \tan (A - B) = (\tan A - \tan B) / (1 + \tan A \tan B)$$

Here

$$\text{LHS} = \tan (A - B)$$

Substituting the values of A and B

$$= \tan (60^\circ - 30^\circ)$$

$$= \tan 30^\circ$$

$$= 1/\sqrt{3}$$

$$\text{RHS} = (\tan A - \tan B) / (1 + \tan A \tan B)$$

Substituting the values of A and B

$$= (\tan 60^\circ - \tan 30^\circ) / (1 + \tan 60^\circ \tan 30^\circ)$$

So we get

$$= \frac{\sqrt{3} - \frac{1}{\sqrt{3}}}{1 + \sqrt{3} \times \frac{1}{\sqrt{3}}}$$

By further simplification

$$= \frac{3 - 1}{\frac{\sqrt{3}}{1+1}}$$

$$= \frac{2}{\frac{\sqrt{3}}{2}}$$

We get

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 2/\sqrt{3} \times \frac{1}{2}$$

$$= 1/\sqrt{3}$$

Therefore, LHS = RHS.

11. (i) If 2θ is an acute angle and $2 \sin 2\theta = \sqrt{3}$, find the value of θ .

(ii) If $20^\circ + x$ is an acute angle and $\cos (20^\circ + x) = \sin 60^\circ$, then find the value of x .

(iii) If $3 \sin^2 \theta = 2 \frac{1}{4}$ and θ is less than 90° , find the value of θ .

Solution:

(i) It is given that

2θ is an acute angle

$$2 \sin 2\theta = \sqrt{3}$$

It can be written as

$$\sin 2\theta = \sqrt{3}/2 = \sin 60^\circ$$

By comparing

$$2\theta = 60^\circ$$

So we get

$$\theta = 60^\circ/2 = 30^\circ$$

Therefore, $\theta = 30^\circ$.

(ii) It is given that

$20^\circ + x$ is an acute angle

$$\cos (20^\circ + x) = \sin 60^\circ$$

It can be written as

$$\cos (20^\circ + x) = \sin 60^\circ = \cos (90^\circ - 60^\circ)$$

$$= \cos 30^\circ$$

By comparing

$$20^\circ + x = 30^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$x = 30^\circ - 20^\circ = 10^\circ$$

Therefore, $x = 10^\circ$.

(iii) It is given that

$$3 \sin^2 \theta = 2 \frac{1}{4}$$

θ is less than 90°

We can write it as

$$\sin^2 \theta = 9/(4 \times 3) = \frac{3}{4}$$

So we get

$$\sin \theta = \sqrt{3}/2 = \sin 60^\circ$$

By comparing

$$\theta = 60^\circ$$

Therefore, $\theta = 60^\circ$.

12. If θ is an acute angle and $\sin \theta = \cos \theta$, find the value of θ and hence, find the value of $2 \tan^2 \theta + \sin^2 \theta - 1$.

Solution:

It is given that

$$\sin \theta = \cos \theta$$

We can write it as

$$\sin \theta / \cos \theta = 1$$

$$\tan \theta = 1$$

We know that $\tan 45^\circ = 1$

$$\tan \theta = \tan 45^\circ$$

So we get

$$\theta = 45^\circ$$

We know that

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$2 \tan^2 \theta + \sin^2 \theta - 1 = 2 \tan^2 45^\circ + \sin^2 45^\circ - 1$$

Substituting the values

$$= 2 (1)^2 + (1/\sqrt{2})^2 - 1$$

By further calculation

$$= 2 \times 1 \times 1 + \frac{1}{2} - 1$$

$$= 2 + \frac{1}{2} - 1$$

$$= \frac{5}{2} - 1$$

Taking LCM

$$= (5 - 2)/2$$

$$= 3/2$$

Therefore, $2 \tan^2 \theta + \sin^2 \theta - 1 = 3/2$.


13. From the adjoining figure, find

(i) $\tan x^\circ$

(ii) x

(iii) $\cos x^\circ$

(iv) use $\sin x^\circ$ to find y .



Solution:

(i) $\tan x^\circ = \text{perpendicular/base}$

It can be written as

$$= AB/BC$$

$$= \sqrt{3}/1$$

$$= \sqrt{3}$$

(ii) $\tan x^\circ = \sqrt{3}$

We know that $\tan 60^\circ = \sqrt{3}$

$$\tan x^\circ = \tan 60^\circ$$

$$x = 60$$

(iii) We know that

$$\cos x^\circ = \cos 60^\circ$$

So we get

$$\cos x^\circ = \frac{1}{2}$$

(iv) $\sin x^\circ = \text{perpendicular/hypotenuse} = AB/AC$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

Substitute $x = 60$ from (ii)

$$\sin 60^\circ = \sqrt{3}/y$$

We know that $\sin 60^\circ = \sqrt{3}/2$

$$\sqrt{3}/2 = \sqrt{3}/y$$

By further calculation

$$y = (\sqrt{3} \times 2) / \sqrt{3}$$

$$y = (2 \times 1) / 1 = 2$$

Therefore, $y = 2$.

14. If 3θ is an acute angle, solve the following equations for θ :

(i) $2 \sin 3\theta = \sqrt{3}$

(ii) $\tan 3\theta = 1$.

Solution:

(i) $2 \sin 3\theta = \sqrt{3}$

It can be written as

$$\sin 3\theta = \sqrt{3}/2$$

We know that $\sin 60^\circ = \sqrt{3}/2$

$$\sin 3\theta = \sin 60^\circ$$

$$3\theta = 60^\circ$$

So we get

$$\theta = 60/3 = 20^\circ$$

(ii) $\tan 3\theta = 1$

We know that $\tan 45^\circ = 1$

$$\tan 3\theta = \tan 45^\circ$$

So we get

$$3\theta = 45^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\theta = 15^\circ$$

15. If $\tan 3x = \sin 45^\circ \cos 45^\circ + \sin 30^\circ$, find the value of x.

Solution:

We know that

$$\tan 3x = \sin 45^\circ \cos 45^\circ + \sin 30^\circ$$

Substituting the values

$$= 1/\sqrt{2} \times 1/\sqrt{2} + \frac{1}{2}$$

$$= \frac{1}{2} + \frac{1}{2}$$

$$= 1$$

We know that

$$\tan 3x = \tan 45^\circ$$

By comparing

$$3x = 45^\circ$$

$$x = 45/3 = 15^\circ$$

Therefore, the value of x is 15° .

16. If $4 \cos^2 x^\circ - 1 = 0$ and $0 \leq x \leq 90$, find

(i) x

(ii) $\sin^2 x^\circ + \cos^2 x^\circ$

(iii) $\cos^2 x^\circ - \sin^2 x^\circ$.

Solution:

It is given that

$$4 \cos^2 x^\circ - 1 = 0$$

$$4 \cos^2 x^\circ = 1$$

It can be written as

$$\cos^2 x^\circ = \frac{1}{4}$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\cos x^\circ = \pm \sqrt{1/4}$$

$$\cos x^\circ = + \sqrt{1/4} \text{ [} 0 \leq x \leq 90^\circ, \text{ then } \cos x^\circ \text{ is positive]}$$

$$\cos x^\circ = 1/2$$

$$\text{We know that } \cos 60^\circ = 1/2$$

$$\cos x^\circ = \cos 60^\circ$$

By comparing

$$x = 60$$

$$\text{(ii) } \sin^2 x^\circ + \cos^2 x^\circ = \sin^2 60^\circ + \cos^2 60^\circ$$

Substituting the values

$$= (\sqrt{3}/2)^2 + (1/2)^2$$

By further calculation

$$= 3/4 + 1/4$$

$$= (3 + 1)/4$$

$$= 4/4$$

$$= 1$$

$$\text{Therefore, } \sin^2 x^\circ + \cos^2 x^\circ = 1.$$

$$\text{(iii) } \cos^2 x^\circ - \sin^2 x^\circ = \cos^2 60^\circ - \sin^2 60^\circ$$

Substituting the values

$$= (1/2)^2 - (\sqrt{3}/2)^2$$

By further calculation

$$= 1/4 - \sqrt{3}/2 \times \sqrt{3}/2$$

$$= 1/4 - 3/4$$

$$= (1 - 3)/4$$

$$= -2/4$$

$$= -1/2$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

Therefore, $\cos^2 x^0 - \sin^2 x^0 = -\frac{1}{2}$.

17. (i) If $\sec \theta = \operatorname{cosec} \theta$ and $0^\circ \leq \theta \leq 90^\circ$, find the value of θ .

(ii) If $\tan \theta = \cot \theta$ and $0^\circ \leq \theta \leq 90^\circ$, find the value of θ

Solution:

(i) It is given that

$$\sec \theta = \operatorname{cosec} \theta$$

We know that

$$\sec \theta = 1 / \cos \theta$$

$$\operatorname{cosec} \theta = 1 / \sin \theta$$

So we get

$$1 / \cos \theta = 1 / \sin \theta$$

$$\sin \theta / \cos \theta = 1$$

$$\tan \theta = 1$$

$$\text{Here } \tan 45^\circ = 1$$

$$\tan \theta = \tan 45^\circ$$

$$\theta = 45^\circ$$

(ii) It is given that

$$\tan \theta = \cot \theta$$

We know that $\cot \theta = 1 / \tan \theta$

$$\tan \theta = 1 / \tan \theta$$

So we get

$$\tan^2 \theta = 1$$

$$\tan \theta = \pm \sqrt{1}$$

$$\tan \theta = +1 \quad [0 \leq \theta \leq 90^\circ, \tan \theta \text{ is positive}]$$

$$\tan \theta = \tan 45^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

By comparing

$$\theta = 45^\circ$$

18. If $\sin 3x = 1$ and $0^\circ \leq 3x \leq 90^\circ$, find the values of

(i) $\sin x$

(ii) $\cos 2x$

(iii) $\tan^2 x - \sec^2 x$.

Solution:

It is given that

$$\sin 3x = 1$$

We know that $\sin 90^\circ = 1$

$$\sin 3x = \sin 90^\circ$$

By comparing

$$3x = 90$$

$$x = 90/3$$

$$x = 30^\circ$$

$$(i) \sin x = \sin 30^\circ = 1/2$$

$$(ii) \cos 2x = \cos 2 \times 30 = \cos 60^\circ = 1/2$$

$$(iii) \tan^2 x - \sec^2 x = \tan^2 30^\circ - \sec^2 30^\circ$$

Substituting the values

$$= (1/\sqrt{3})^2 - (2/\sqrt{3})^2$$

By further calculation

$$= 1/3 - 4/3$$

$$= (1 - 4)/3$$

$$= -3/3$$

$$= -1$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

Therefore, $\tan^2 x - \sec^2 x = -1$.

19. If $3 \tan^2 \theta - 1 = 0$, find $\cos 2\theta$, given that θ is acute.

Solution:

It is given that

$$3 \tan^2 \theta - 1 = 0$$

We can write it as

$$3 \tan^2 \theta = 1$$

$$\tan^2 \theta = 1/3$$

$$\tan \theta = 1/\sqrt{3} \text{ } [\theta \text{ is acute so } \tan \theta \text{ is positive}]$$

$$\theta = 30^\circ$$

So we get

$$\cos 2\theta = \cos 2 \times 30^\circ = \cos 60^\circ = \frac{1}{2}$$

20. If $\sin x + \cos y = 1$, $x = 30^\circ$ and y is acute angle, find the value of y .

Solution:

It is given that

$$\sin x + \cos y = 1$$

$$x = 30^\circ$$

Substituting the values

$$\sin 30^\circ + \cos y = 1$$

$$\frac{1}{2} + \cos y = 1$$

It can be written as

$$\cos y = 1 - \frac{1}{2}$$

Taking LCM

$$\cos y = (2 - 1)/2 = \frac{1}{2}$$

We know that $\cos 60^\circ = \frac{1}{2}$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\cos y = \cos 60^\circ$$

So we get

$$y = 60^\circ$$

21. If $\sin (A + B) = \sqrt{3}/2 = \cos (A - B)$, $0^\circ < A + B \leq 90^\circ$ ($A > B$), find the values of A and B.

Solution:

It is given that

$$\sin (A + B) = \sqrt{3}/2 = \cos (A - B)$$

Consider

$$\sin (A + B) = \sqrt{3}/2$$

$$\text{We know that } \sin 60^\circ = \sqrt{3}/2$$

$$\sin (A + B) = \sin 60^\circ$$

$$A + B = 60^\circ \dots\dots (1)$$

Similarly

$$\cos (A - B) = \sqrt{3}/2$$

$$\text{We know that } \cos 30^\circ = \sqrt{3}/2$$

$$\cos (A - B) = \cos 30^\circ$$

$$A - B = 30^\circ \dots\dots (2)$$

By adding both the equations

$$A + B + A - B = 60^\circ + 30^\circ$$

So we get

$$2A = 90^\circ$$

$$A = 90^\circ/2 = 45^\circ$$

Now substitute the value of A in equation (1)

$$45^\circ + B = 60^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

By further calculation

$$B = 60^\circ - 45^\circ = 15^\circ$$

Therefore, $A = 45^\circ$ and $B = 15^\circ$.


22. If the length of each side of a rhombus is 8 cm and its one angle is 60° , then find the lengths of the diagonals of the rhombus.

Solution:

It is given that

Each side of a rhombus = 8 cm

One angle = 60°



We know that the diagonals bisect the opposite angles

$$\angle OAB = 60^\circ/2 = 30^\circ$$

In right $\angle AOB$

$$\sin 30^\circ = OB/AB$$

So we get

$$\frac{1}{2} = OB/8$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

By further calculation

$$OB = 8/2 = 4 \text{ cm}$$

$$BD = 2OB = 2 \times 4 = 8 \text{ cm}$$

$$\cos 30^\circ = AO/AB$$

Substituting the values

$$\sqrt{3}/2 = AO/8$$

By further calculation

$$AO = 8\sqrt{3}/2 = 4\sqrt{3}$$

Here

$$AC = 4\sqrt{3} \times 2 = 8\sqrt{3} \text{ cm}$$

Therefore, the length of the diagonals of the rhombus are 8 cm and $8\sqrt{3}$ cm.

23. In the right-angled triangle ABC, $\angle C = 90^\circ$ and $\angle B = 60^\circ$. If AC = 6 cm, find the lengths of the sides BC and AB.

Solution:

In the right-angled triangle ABC, $\angle C = 90^\circ$ and $\angle B = 60^\circ$

$$AC = 6 \text{ cm}$$

We know that

$$\tan B = AC/BC$$

Substituting the values

$$\tan 60^\circ = 6/BC$$

So we get

$$\sqrt{3} = 6/BC$$

$$BC = 6/\sqrt{3}$$

It can be written as

$$= 6\sqrt{3}/(\sqrt{3} + \sqrt{3})$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 6\sqrt{3}/3$$

$$= 2\sqrt{3} \text{ cm}$$

$$\sin 60^\circ = AC/AB$$

Substituting the values

$$\sqrt{3}/2 = 6/AB$$

By further calculation

$$AB = (6 \times 2)/\sqrt{3}$$

So we get


$$AB = (12 \times \sqrt{3})/(\sqrt{3} \times \sqrt{3})$$

$$= 12\sqrt{3}/3$$

$$= 4\sqrt{3} \text{ cm}$$

Therefore, the lengths of the sides $BC = 2\sqrt{3} \text{ cm}$ and $AB = 4\sqrt{3} \text{ cm}$.

24. In the adjoining figure, AP is a man of height 1.8 m and BQ is a building 13.8 m high. If the man sees the top of the building by focusing his binoculars at an angle of 30° to the horizontal, find the distance of the man from the building.



Solution:

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

It is given that

Height of man AP = 1.8 m

Height of building BQ = 13.8 m

Angle of elevation from the top of the building to the man = 30°

Consider AB as the distance of the man from the building

AB = x then PC = x

QC = 13.8 – 1.8 = 12 m

In right $\triangle PQC$

$\tan \theta = QC/PC$

Substituting the values

$\tan 30^\circ = 12/x$

By further calculation


$1/\sqrt{3} = 12/x$

$x = 12\sqrt{3}$ m

Therefore, the distance of the man from the building is $12\sqrt{3}$ m.

25. In the adjoining figure, ABC is a triangle in which $\angle B = 45^\circ$ and $\angle C = 60^\circ$. If $AD \perp BC$ and BC = 8 m, find the length of the altitude AD.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

**Solution:**

In triangle ABC

$$\angle B = 45^\circ \text{ and } \angle C = 60^\circ$$

$AD \perp BC$ and $BC = 8 \text{ m}$

In right $\triangle ABD$

$$\tan 45^\circ = AD/BD$$

So we get

$$1 = AD/BD$$

$$AD = BD$$

In right $\triangle ACD$

$$\tan 60^\circ = AD/DC$$

So we get

$$\sqrt{3} = AD/DC$$

$$DC = AD/\sqrt{3}$$

Here

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$BD + DC = AD + AD/\sqrt{3}$$

Taking LCM

$$BC = (\sqrt{3}AD + AD)/\sqrt{3}$$

$$8 = [AD(\sqrt{3} + 1)]/\sqrt{3}$$

By further calculation

$$AD = 8\sqrt{3}/(\sqrt{3} + 1)$$

It can be written as

$$= \frac{8\sqrt{3}(\sqrt{3} - 1)}{(\sqrt{3} + 1)(\sqrt{3} - 1)}$$

So we get

$$= \frac{8(3 - \sqrt{3})}{3 - 1}$$

$$= \frac{8(3 - \sqrt{3})}{2}$$

$$= 4(3 - \sqrt{3}) \text{ m}$$

Therefore, the length of the altitude AD is $4(3 - \sqrt{3})$ m.

Exercise 18.2

Without using trigonometric tables, evaluate the following (1 to 5):

1. (i) $\cos 18^\circ/\sin 72^\circ$

(ii) $\tan 41^\circ/\cot 49^\circ$

(iii) $\operatorname{cosec} 17^\circ 30'/\sec 72^\circ 30'$

Solution:

(i) $\cos 18^\circ/\sin 72^\circ$

It can be written as

$$= \cos 18^\circ/\sin (90^\circ - 18^\circ)$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

So we get

$$= \cos 18^\circ / \cos 18^\circ$$

$$= 1$$

$$(ii) \tan 41^\circ / \cot 49^\circ$$

It can be written as

$$= \tan 41^\circ / \cot (90^\circ - 41^\circ)$$

So we get

$$= \tan 41^\circ / \tan 41^\circ$$

$$= 1$$

$$(iii) \operatorname{cosec} 17^\circ 30' / \sec 72^\circ 30'$$

It can be written as

$$= \operatorname{cosec} 17^\circ 30' / \sec (90^\circ - 17^\circ 30')$$

So we get

$$= \operatorname{cosec} 17^\circ 30' / \operatorname{cosec} 17^\circ 30'$$

$$= 1$$

$$2. (i) \cot 40^\circ / \tan 50^\circ - \frac{1}{2} (\cos 35^\circ / \sin 55^\circ)$$

$$(ii) (\sin 49^\circ / \cos 41^\circ)^2 + (\cos 41^\circ / \sin 49^\circ)^2$$

$$(iii) \sin 72^\circ / \cos 18^\circ - \sec 32^\circ / \operatorname{cosec} 58^\circ$$

$$(iv) \cos 75^\circ / \sin 15^\circ + \sin 12^\circ / \cos 78^\circ - \cos 18^\circ / \sin 72^\circ$$

$$(v) \sin 25^\circ / \sec 65^\circ + \cos 25^\circ / \operatorname{cosec} 65^\circ.$$

Solution:

$$(i) \cot 40^\circ / \tan 50^\circ - \frac{1}{2} (\cos 35^\circ / \sin 55^\circ)$$

It can be written as

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \frac{\cot 40^\circ}{\tan(90^\circ - 40^\circ)} - \frac{1}{2} \frac{\cos 35^\circ}{\sin(90^\circ - 35^\circ)}$$

By further calculation

$$\begin{aligned} &= \frac{\cot 40^\circ}{\cot 40^\circ} - \frac{1}{2} \frac{\cos 35^\circ}{\cos 35^\circ} \\ &= 1 - \frac{1}{2} \\ &= \frac{1}{2} \end{aligned}$$

(ii) $(\sin 49^\circ / \cos 41^\circ)^2 + (\cos 41^\circ / \sin 49^\circ)^2$

It can be written as

$$= \left[\frac{\sin 49^\circ}{\cos(90^\circ - 49^\circ)} \right]^2 + \left[\frac{\cos 41^\circ}{\sin(90^\circ - 41^\circ)} \right]^2$$

By further calculation

$$= \left(\frac{\sin 49^\circ}{\sin 49^\circ} \right)^2 + \left(\frac{\cos 41^\circ}{\cos 41^\circ} \right)^2$$

So we get

$$= 1^2 + 1^2$$

$$= 1 + 1$$

$$= 2$$

(iii) $\sin 72^\circ / \cos 18^\circ - \sec 32^\circ / \operatorname{cosec} 58^\circ$

It can be written as

$$= \frac{\sin 72^\circ}{\cos(90^\circ - 72^\circ)} - \frac{\sec 32^\circ}{\operatorname{cosec}(90^\circ - 32^\circ)}$$

By further calculation

$$= \frac{\sin 72^\circ}{\sin 72^\circ} - \frac{\sec 32^\circ}{\sec 32^\circ}$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 1 - 1$$

$$= 0$$

$$(iv) \cos 75^\circ / \sin 15^\circ + \sin 12^\circ / \cos 78^\circ - \cos 18^\circ / \sin 72^\circ$$

It can be written as

$$= \frac{\cos 75^\circ}{\sin(90^\circ - 75^\circ)} + \frac{\sin 12^\circ}{\cos(90^\circ - 12^\circ)} - \frac{\cos 18^\circ}{\sin(90^\circ - 18^\circ)}$$

By further calculation

$$= \frac{\cos 75^\circ}{\cos 75^\circ} + \frac{\sin 12^\circ}{\sin 12^\circ} - \frac{\cos 18^\circ}{\cos 18^\circ}$$

So we get

$$= 1 + 1 - 1$$

$$= 1$$

$$(v) \sin 25^\circ / \sec 65^\circ + \cos 25^\circ / \operatorname{cosec} 65^\circ$$

It can be written as

$$= (\sin 25^\circ \times \cos 65^\circ) + (\cos 25^\circ \times \sin 65^\circ)$$

By further calculation

$$= [\sin 25^\circ \times \cos(90^\circ - 25^\circ)] + [\cos 25^\circ \times \sin(90^\circ - 25^\circ)]$$

So we get

$$= (\sin 25^\circ \times \sin 25^\circ) + (\cos 25^\circ \times \cos 25^\circ)$$

$$= \sin^2 25^\circ + \cos^2 25^\circ$$

$$= 1$$

$$3. (i) \sin 62^\circ - \cos 28^\circ$$

$$(ii) \operatorname{cosec} 35^\circ - \sec 55^\circ.$$

Solution:

$$(i) \sin 62^\circ - \cos 28^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

It can be written as

$$= \sin (90^\circ - 28^\circ) - \cos 28^\circ$$

So we get

$$= \cos 28^\circ - \cos 28^\circ$$

$$= 0$$

$$(ii) \operatorname{cosec} 35^\circ - \sec 55^\circ$$

It can be written as

$$= \operatorname{cosec} 35^\circ - \sec (90^\circ - 35^\circ)$$

So we get

$$= \operatorname{cosec} 35^\circ - \operatorname{cosec} 35^\circ$$

$$= 0$$

$$4. (i) \cos^2 26^\circ + \cos 64^\circ \sin 26^\circ + \tan 36^\circ / \cot 54^\circ$$

$$(ii) \sec 17^\circ / \operatorname{cosec} 73^\circ + \tan 68^\circ / \cot 22^\circ + \cos^2 44^\circ + \cos^2 46^\circ.$$

Solution:

$$(i) \cos^2 26^\circ + \cos 64^\circ \sin 26^\circ + \tan 36^\circ / \cot 54^\circ$$

It can be written as

$$= \cos^2 26^\circ + \cos (90^\circ - 26^\circ) \sin 26^\circ + \tan 36^\circ / \cot (90^\circ - 36^\circ)$$

We know that

$$\cos (90^\circ - \theta) = \sin \theta$$

$$\cot (90^\circ - \theta) = \tan \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

So we get

$$= \cos^2 26^\circ + \sin^2 26^\circ + \tan 36^\circ / \tan 36^\circ$$

$$= 1 + 1$$

$$= 2$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(ii) \sec 17^\circ / \operatorname{cosec} 73^\circ + \tan 68^\circ / \cot 22^\circ + \cos^2 44^\circ + \cos^2 46^\circ$$

It can be written as

$$= \sec (90^\circ - 73^\circ) / \operatorname{cosec} 73^\circ + \tan (90^\circ - 22^\circ) / \cot 22^\circ + \cos^2 (90^\circ - 46^\circ) + \cos^2 46^\circ$$

By further calculation

$$= \operatorname{cosec} 73^\circ / \operatorname{cosec} 73^\circ + \cot 22^\circ / \cot 22^\circ + \sin^2 46^\circ + \cos^2 46^\circ$$

We know that $\sin^2 \theta + \cos^2 \theta = 1$

So we get

$$= 1 + 1 + 1$$

$$= 3$$

$$5. (i) \cos 65^\circ / \sin 25^\circ + \cos 32^\circ / \sin 58^\circ - \sin 28^\circ \sec 62^\circ + \operatorname{cosec}^2 30^\circ$$

$$(ii) \sec 29^\circ / \operatorname{cosec} 61^\circ + 2 \cot 8^\circ \cot 17^\circ \cot 45^\circ \cot 73^\circ \cot 82^\circ - 3 (\sin^2 38^\circ + \sin^2 52^\circ).$$

Solution:

$$(i) \cos 65^\circ / \sin 25^\circ + \cos 32^\circ / \sin 58^\circ - \sin 28^\circ \sec 62^\circ + \operatorname{cosec}^2 30^\circ$$

It can be written as

$$= \cos 65^\circ / \sin (90^\circ - 65^\circ) + \cos 32^\circ / \sin (90^\circ - 32^\circ) - \sin 28^\circ \sec (90^\circ - 28^\circ) + \operatorname{cosec}^2 30^\circ$$

By further calculation

$$= \cos 65^\circ / \cos 65^\circ + \cos 32^\circ / \cos 32^\circ - \sin 28^\circ \operatorname{cosec} 28^\circ + \operatorname{cosec}^2 30^\circ$$

We know that $\operatorname{cosec} 30^\circ = 2$

$$= 1 + 1 - 1 + 4$$

$$= 5$$

$$(ii) \sec 29^\circ / \operatorname{cosec} 61^\circ + 2 \cot 8^\circ \cot 17^\circ \cot 45^\circ \cot 73^\circ \cot 82^\circ - 3 (\sin^2 38^\circ + \sin^2 52^\circ)$$

It can be written as

$$= \sec 29^\circ / \operatorname{cosec} (90^\circ - 29^\circ) + 2 \cot 8^\circ \cot 17^\circ \cot 45^\circ \cot (90^\circ - 17^\circ) \cot (90^\circ - 8^\circ) - 3 [\sin^2 38^\circ + \sin^2 (90^\circ - 38^\circ)]$$

By further calculation

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \sec 29^\circ / \sec 29^\circ + 2 \cot 8^\circ \cot 17^\circ \times 1 \times \tan 17^\circ \tan 8^\circ - 3 (\sin^2 38^\circ + \cos^2 38^\circ)$$

So we get

$$= 1 + 2 \cot 8^\circ \tan 8^\circ \cot 17^\circ \tan 17^\circ \times 1 - 3 \times 1$$

We know that

$$\operatorname{cosec} (90^\circ - \theta) = \sec \theta$$

$$\cot (90^\circ - \theta) = \tan \theta$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

Here

$$= 1 + 2 \times 1 \times 1 \times 1 - 3$$

$$= 1 + 2 - 3$$

$$= 0$$

6. Express each of the following in terms of trigonometric ratios of angles between 0° to 45° :

(i) $\tan 81^\circ + \cos 72^\circ$

(ii) $\cot 49^\circ + \operatorname{cosec} 87^\circ$.

Solution:

(i) $\tan 81^\circ + \cos 72^\circ$

It can be written as

$$= \tan (90^\circ - 9^\circ) + \cos (90^\circ - 18^\circ)$$

So we get

$$= \cot 9^\circ + \sin 18^\circ$$

(ii) $\cot 49^\circ + \operatorname{cosec} 87^\circ$

It can be written as

$$= \cot (90^\circ - 41^\circ) + \operatorname{cosec} (90^\circ - 3^\circ)$$

So we get

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \tan 41^\circ + \sec 3^\circ$$

Without using trigonometric tables, prove that (7 to 11):

7. (i) $\sin^2 28^\circ - \cos^2 62^\circ = 0$

(ii) $\cos^2 25^\circ + \cos^2 65^\circ = 1$

(iii) $\operatorname{cosec}^2 67^\circ - \tan^2 23^\circ = 1$

(iv) $\sec^2 22^\circ - \cot^2 68^\circ = 1$.

Solution:

(i) $\sin^2 28^\circ - \cos^2 62^\circ = 0$

Consider

$$\text{LHS} = \sin^2 28^\circ - \cos^2 62^\circ$$

It can be written as

$$= \sin^2 28^\circ - \cos^2 (90^\circ - 28^\circ)$$

So we get

$$= \sin^2 28^\circ - \sin^2 28^\circ$$

$$= 0$$

$$= \text{RHS}$$

(ii) $\cos^2 25^\circ + \cos^2 65^\circ = 1$

Consider

$$\text{LHS} = \cos^2 25^\circ + \cos^2 65^\circ$$

It can be written as

$$= \cos^2 25^\circ + \cos^2 (90^\circ - 25^\circ)$$

$$\text{We know that } \sin^2 \theta + \cos^2 \theta = 1$$

So we get

$$= \cos^2 25^\circ + \sin^2 25^\circ$$

$$= 1$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(iii) \operatorname{cosec}^2 67^\circ - \tan^2 23^\circ = 1$$

Consider

$$\text{LHS} = \operatorname{cosec}^2 67^\circ - \tan^2 23^\circ$$

It can be written as

$$= \operatorname{cosec}^2 67^\circ - \tan^2 (90^\circ - 67^\circ)$$

$$\text{We know that } \operatorname{cosec}^2 \theta - \cot^2 \theta = 1$$

So we get

$$= \operatorname{cosec}^2 67^\circ - \cot^2 67^\circ$$

$$= 1$$

$$(iv) \sec^2 22^\circ - \cot^2 68^\circ = 1$$

Consider

$$\text{LHS} = \sec^2 22^\circ - \cot^2 68^\circ$$

It can be written as

$$= \sec^2 22^\circ - \cot^2 (90^\circ - 22^\circ)$$

$$\text{We know that } \sec^2 \theta - \tan^2 \theta = 1$$

So we get

$$= \sec^2 22^\circ - \tan^2 22^\circ$$

$$= 1$$

$$8. (i) \sin 63^\circ \cos 27^\circ + \cos 63^\circ \sin 27^\circ = 1$$

$$(ii) \sec 31^\circ \sin 59^\circ + \cos 31^\circ \operatorname{cosec} 59^\circ = 2.$$

Solution:

$$(i) \sin 63^\circ \cos 27^\circ + \cos 63^\circ \sin 27^\circ = 1$$

Consider

$$\text{LHS} = \sin 63^\circ \cos 27^\circ + \cos 63^\circ \sin 27^\circ$$

It can be written as

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \sin 63^\circ \cos (90^\circ - 63^\circ) + \cos 63^\circ \sin (90^\circ - 63^\circ)$$

$$= \sin 63^\circ \sin 63^\circ + \cos 63^\circ \cos 63^\circ$$

We know that $\sin^2 \theta + \cos^2 \theta = 1$

So we get

$$= \sin^2 63^\circ + \cos^2 63^\circ$$

$$= 1$$

$$(ii) \sec 31^\circ \sin 59^\circ + \cos 31^\circ \operatorname{cosec} 59^\circ = 2$$

Consider

$$\text{LHS} = \sec 31^\circ \sin 59^\circ + \cos 31^\circ \operatorname{cosec} 59^\circ$$

It can be written as

$$= 1/\cos 31^\circ \times \sin 59^\circ + \cos 31^\circ \times 1/\sin 59^\circ$$

By further calculation

$$= \sin 59^\circ / \cos (90^\circ - 59^\circ) + \cos 31^\circ / \sin (90^\circ - 31^\circ)$$

So we get

$$= \sin 59^\circ / \sin 59^\circ + \cos 31^\circ / \cos 31^\circ$$

$$= 1 + 1$$

$$= 2$$

$$= \text{RHS}$$

$$9. (i) \sec 70^\circ \sin 20^\circ - \cos 20^\circ \operatorname{cosec} 70^\circ = 0$$

$$(ii) \sin^2 20^\circ + \sin^2 70^\circ - \tan^2 45^\circ = 0.$$

Solution:

$$(i) \sec 70^\circ \sin 20^\circ - \cos 20^\circ \operatorname{cosec} 70^\circ = 0$$

Consider

$$\text{LHS} = \sec 70^\circ \sin 20^\circ - \cos 20^\circ \operatorname{cosec} 70^\circ$$

By further simplification

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \sin 20^\circ / \cos 70^\circ - \cos 20^\circ / \sin 70^\circ$$

It can be written as

$$= \sin 20^\circ / \cos (90^\circ - 20^\circ) - \cos 20^\circ / \sin (90^\circ - 20^\circ)$$

So we get

$$= \sin 20^\circ / \sin 20^\circ - \cos 20^\circ / \cos 20^\circ$$

$$= 1 - 1$$

$$= 0$$

$$= \text{RHS}$$

$$(ii) \sin^2 20^\circ + \sin^2 70^\circ - \tan^2 45^\circ = 0$$

Consider

$$\text{LHS} = \sin^2 20^\circ + \sin^2 70^\circ - \tan^2 45^\circ$$

It can be written as

$$= \sin^2 20^\circ + \sin^2 (90^\circ - 20^\circ) - \tan^2 45^\circ$$

By further calculation

$$= \sin^2 20^\circ + \cos^2 20^\circ - \tan^2 45^\circ$$

$$\text{We know that } \sin^2 \theta + \cos^2 \theta = 1 \text{ and } \tan 45^\circ = 1$$

So we get

$$= 1 - 1$$

$$= 0$$

$$= \text{RHS}$$

$$10. (i) \cot 54^\circ / \tan 36^\circ + \tan 20^\circ / \cot 70^\circ - 2 = 0$$

$$(ii) \sin 50^\circ / \cos 40^\circ + \operatorname{cosec} 40^\circ / \sec 50^\circ - 4 \cos 50^\circ \operatorname{cosec} 40^\circ + 2 = 0.$$

Solution:

$$(i) \cot 54^\circ / \tan 36^\circ + \tan 20^\circ / \cot 70^\circ - 2 = 0$$

Consider

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\text{LHS} = \cot 54^\circ / \tan 36^\circ + \tan 20^\circ / \cot 70^\circ - 2$$

It can be written as

$$= \cot 54^\circ / \tan (90^\circ - 54^\circ) + \tan 20^\circ / \cot (90^\circ - 20^\circ) - 2$$

By further calculation

$$= \cot 54^\circ / \cot 54^\circ + \tan 20^\circ / \tan 20^\circ - 2$$

So we get

$$= 1 + 1 - 2$$

$$= 0$$

$$= \text{RHS}$$

$$(ii) \sin 50^\circ / \cos 40^\circ + \operatorname{cosec} 40^\circ / \sec 50^\circ - 4 \cos 50^\circ \operatorname{cosec} 40^\circ + 2 = 0$$

Consider

$$\text{LHS} = \sin 50^\circ / \cos 40^\circ + \operatorname{cosec} 40^\circ / \sec 50^\circ - 4 \cos 50^\circ \operatorname{cosec} 40^\circ + 2$$

It can be written as

$$= \sin 50^\circ / \cos (90^\circ - 50^\circ) + \operatorname{cosec} 40^\circ / \sec (90^\circ - 40^\circ) - 4 \cos 50^\circ \operatorname{cosec} (90^\circ - 50^\circ) + 2$$

By further calculation

$$= \sin 50^\circ / \sin 50^\circ + \operatorname{cosec} 40^\circ / \operatorname{cosec} 40^\circ - \cos 50^\circ \sec 50^\circ + 2$$

So we get

$$= 1 + 1 - 4 \cos 50^\circ / \cos 50^\circ + 2$$

$$= 1 + 1 - 4 + 2$$

$$= 4 - 4$$

$$= 0$$

$$= \text{RHS}$$

$$11. (i) \cos 70^\circ / \sin 20^\circ + \cos 59^\circ / \sin 31^\circ - 8 \sin^2 30^\circ = 0$$

$$(ii) \cos 80^\circ / \sin 10^\circ + \cos 59^\circ \operatorname{cosec} 31^\circ = 2.$$

Solution:

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(i) \cos 70^\circ / \sin 20^\circ + \cos 59^\circ / \sin 31^\circ - 8 \sin^2 30^\circ = 0$$

Consider

$$\text{LHS} = \cos 70^\circ / \sin 20^\circ + \cos 59^\circ / \sin 31^\circ - 8 \sin^2 30^\circ$$

It can be written as

$$= \cos 70^\circ / \sin (90^\circ - 70^\circ) + \cos 59^\circ / \sin (90^\circ - 59^\circ) - 8 \sin^2 30^\circ$$

We know that $\sin 30^\circ = \frac{1}{2}$

$$= \cos 70^\circ / \cos 70^\circ + \cos 59^\circ / \cos 59^\circ - 8 \left(\frac{1}{2}\right)^2$$

By further calculation

$$= 1 + 1 - 8 \times \frac{1}{4}$$

So we get

$$= 2 - 2$$

$$= 0$$

$$= \text{RHS}$$

$$(ii) \cos 80^\circ / \sin 10^\circ + \cos 59^\circ \operatorname{cosec} 31^\circ = 2$$

Consider

$$\text{LHS} = \cos 80^\circ / \sin 10^\circ + \cos 59^\circ \operatorname{cosec} 31^\circ$$

It can be written as

$$= \cos 80^\circ / \sin (90^\circ - 80^\circ) + \cos 59^\circ / \sin 31^\circ$$

By further simplification

$$= \cos 80^\circ / \cos 80^\circ + \cos 59^\circ / \sin (90^\circ - 59^\circ)$$

So we get

$$= 1 + \cos 59^\circ / \cos 59^\circ$$

$$= 1 + 1$$

$$= 2$$

$$= \text{RHS}$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

12. Without using trigonometrical tables, evaluate:

$$(i) 2\left(\frac{\tan 35^\circ}{\cot 55^\circ}\right)^2 + \left(\frac{\cot 55^\circ}{\tan 35^\circ}\right) - 3\left(\frac{\sec 40^\circ}{\operatorname{cosec} 50^\circ}\right)$$

$$(ii) \frac{\sin 35^\circ \cos 55^\circ + \cos 35^\circ \sin 55^\circ}{\operatorname{cosec}^2 10^\circ - \tan^2 80^\circ}$$

$$(iii) \sin^2 34^\circ + \sin^2 56^\circ + 2 \tan 18^\circ \tan 72^\circ - \cot^2 30^\circ.$$

Solution:

$$(i) 2\left(\frac{\tan 35^\circ}{\cot 55^\circ}\right)^2 + \left(\frac{\cot 55^\circ}{\tan 35^\circ}\right) - 3\left(\frac{\sec 40^\circ}{\operatorname{cosec} 50^\circ}\right)$$

It can be written as

$$= 2\left[\frac{\tan(90^\circ - 55^\circ)}{\cot 55^\circ}\right] + \left[\frac{\cot(90^\circ - 35^\circ)}{\tan 35^\circ}\right] - 3\left[\frac{\sec(90^\circ - 50^\circ)}{\operatorname{cosec} 50^\circ}\right]$$

By further calculation

$$= 2\left(\frac{\cot 55^\circ}{\cot 55^\circ}\right)^2 + \left(\frac{\tan 35^\circ}{\tan 35^\circ}\right) - 3\left(\frac{\operatorname{cosec} 50^\circ}{\operatorname{cosec} 50^\circ}\right)$$

So we get

$$= 2 + 1 - 3$$

$$= 0$$

$$(ii) \frac{\sin 35^\circ \cos 55^\circ + \cos 35^\circ \sin 55^\circ}{\operatorname{cosec}^2 10^\circ - \tan^2 80^\circ}$$

It can be written as

$$= \frac{\sin 35^\circ \cos(90^\circ - 35^\circ) + \cos 35^\circ \sin(90^\circ - 35^\circ)}{\operatorname{cosec}^2 10^\circ - \tan^2(90^\circ - 10^\circ)}$$

By further calculation

$$= \frac{\sin 35^\circ \cdot \sin 35^\circ + \cos 35^\circ \cdot \cos 35^\circ}{\operatorname{cosec}^2 10^\circ - \cot^2 10^\circ}$$

So we get

$$= \frac{\sin^2 35^\circ + \cos^2 35^\circ}{\operatorname{cosec}^2 10^\circ - \cot^2 10^\circ}$$

We know that $\sin^2 \theta + \cos^2 \theta = 1$ and $\operatorname{cosec}^2 \theta - \cot^2 \theta = 1$

$$= 1/1$$

$$= 1$$

$$(iii) \sin^2 34^\circ + \sin^2 56^\circ + 2 \tan 18^\circ \tan 72^\circ - \cot^2 30^\circ$$

It can be written as

$$= \sin^2 34^\circ + [\sin(90^\circ - 34^\circ)]^2 + 2 \tan 18^\circ \tan(90^\circ - 18^\circ) - \cot^2 30^\circ$$

By further simplification

$$= \sin^2 34^\circ + \cos^2 34^\circ + 2 \tan 18^\circ \cot 18^\circ - (\sqrt{3})^2$$

So we get

$$= 1 + 2 \tan 18^\circ \times 1/\tan 18^\circ - 3$$

$$= 1 + 2 - 3$$

$$= 0$$

13. Prove the following:

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(i) \frac{\cos \theta}{\sin(90^\circ - \theta)} + \frac{\sin \theta}{\cos(90^\circ - \theta)} = 2$$

$$(ii) \cos \theta \sin(90^\circ - \theta) + \sin \theta \cos(90^\circ - \theta) = 1$$

$$(iii) \frac{\tan \theta}{\tan(90^\circ - \theta)} + \frac{\sin(90^\circ - \theta)}{\cos \theta} = \sec^2 \theta.$$

Solution:

$$(i) \frac{\cos \theta}{\sin(90^\circ - \theta)} + \frac{\sin \theta}{\cos(90^\circ - \theta)} = 2$$

We know that

LHS =

$$\frac{\cos \theta}{\sin(90^\circ - \theta)} + \frac{\sin \theta}{\cos(90^\circ - \theta)}$$

So we get

$$= \cos \theta / \cos \theta + \sin \theta / \sin \theta$$

$$= 1 + 1$$

$$= 2$$

$$= \text{RHS}$$

$$(ii) \cos \theta \sin(90^\circ - \theta) + \sin \theta \cos(90^\circ - \theta) = 1$$

Consider

$$\text{LHS} = \cos \theta \sin(90^\circ - \theta) + \sin \theta \cos(90^\circ - \theta)$$

It can be written as

$$= \cos \theta \cdot \cos \theta + \sin \theta \cdot \sin \theta$$

So we get

$$= \cos^2 \theta + \sin^2 \theta$$

$$= 1$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

= RHS

$$(iii) \frac{\tan \Theta}{\tan(90^\circ - \Theta)} + \frac{\sin(90^\circ - \Theta)}{\cos \Theta} = \sec^2 \Theta$$

Consider

LHS =

$$\frac{\tan \Theta}{\tan(90^\circ - \Theta)} + \frac{\sin(90^\circ - \Theta)}{\cos \Theta}$$

By further calculation

$$= \tan \theta / \cot \theta + \cos \theta / \cos \theta$$

So we get

$$= \tan \theta \times \tan \theta + 1$$

$$= \tan^2 \theta + 1$$

$$= \sec^2 \theta$$

= RHS

14. Prove the following:

$$(i) \frac{\cos(90^\circ - A) \sin(90^\circ - A)}{\tan(90^\circ - A)} = 1 - \cos^2 A$$

$$(ii) \frac{\sin(90^\circ - A)}{\operatorname{cosec}(90^\circ - A)} + \frac{\cos(90^\circ - A)}{\sec(90^\circ - A)} = 1.$$

Solution:

$$(i) \frac{\cos(90^\circ - A) \sin(90^\circ - A)}{\tan(90^\circ - A)} = 1 - \cos^2 A$$

Consider

LHS =

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\frac{\cos(90^\circ - A)\sin(90^\circ - A)}{\tan(90^\circ - A)}$$

It can be written as

$$= \sin A \cos A / \cot A$$

So we get

$$= (\sin A \cos A \times \sin A) / \cos A$$

$$= \sin^2 A$$

$$= 1 - \cos^2 A$$

$$= \text{RHS}$$

$$(ii) \frac{\sin(90^\circ - A)}{\operatorname{cosec}(90^\circ - A)} + \frac{\cos(90^\circ - A)}{\sec(90^\circ - A)} = 1$$

Consider

LHS =

$$\frac{\sin(90^\circ - A)}{\operatorname{cosec}(90^\circ - A)} + \frac{\cos(90^\circ - A)}{\sec(90^\circ - A)}$$

It can be written as

$$= \cos A / \sec A + \sin A / \operatorname{cosec} A$$

So we get

$$= \cos A \times \cos A + \sin A \times \sin A$$

$$= \cos^2 A + \sin^2 A$$

$$= 1$$

$$= \text{RHS}$$

15. Simplify the following:

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(i) \frac{\cos \Theta}{\sin(90^\circ - \Theta)} + \frac{\cos(90^\circ - \Theta)}{\sec(90^\circ - \Theta)} - 3 \tan^2 30^\circ$$

$$(ii) \frac{\operatorname{cosec}(90^\circ - \Theta) \sin(90^\circ - \Theta) \cot(90^\circ - \Theta)}{\cos(90^\circ - \Theta) \sec(90^\circ - \Theta) \tan \Theta} + \frac{\cot \Theta}{\tan(90^\circ - \Theta)}.$$

Solution:

$$(i) \frac{\cos \Theta}{\sin(90^\circ - \Theta)} + \frac{\cos(90^\circ - \Theta)}{\sec(90^\circ - \Theta)} - 3 \tan^2 30^\circ$$

It can be written as

$$= \cos \theta / \cos \theta + \sin \theta / \operatorname{cosec} \theta - 3 \tan^2 30^\circ$$

By further calculation

$$= 1 + \sin \theta \times \sin \theta - 3 (1/\sqrt{3})^2$$

So we get

$$= \sin^2 \theta + 1 - 3 \times 1/3$$

$$= \sin^2 \theta + 1 - 1$$

$$= \sin^2 \theta +$$

$$(ii) \frac{\operatorname{cosec}(90^\circ - \Theta) \sin(90^\circ - \Theta) \cot(90^\circ - \Theta)}{\cos(90^\circ - \Theta) \sec(90^\circ - \Theta) \tan \Theta} + \frac{\cot \Theta}{\tan(90^\circ - \Theta)}$$

It can be written as

$$= (\sec \theta \cos \theta \tan \theta) / (\sin \theta \operatorname{cosec} \theta \tan \theta) + \cot \theta / \cot \theta$$

So we get

$$= \sec \theta \cos \theta / \sin \theta \operatorname{cosec} \theta + 1$$

$$= 1/1 + 1$$

$$= 1 + 1$$

$$= 2$$

16. Show that

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\frac{\cos^2(45^\circ + \Theta) + \cos^2(45^\circ - \Theta)}{\tan(60^\circ + \Theta)\tan(30^\circ - \Theta)} = 1.$$

Solution:

Consider

$$LHS = \frac{\cos^2(45^\circ + \Theta) + \cos^2(45^\circ - \Theta)}{\tan(60^\circ + \Theta)\tan(30^\circ - \Theta)}$$

It can be written as

$$= \frac{\cos^2(45^\circ + \Theta) + \cos^2[90^\circ - (45^\circ - \Theta)]}{\tan(60^\circ + \Theta)\tan[90^\circ - (60^\circ - \Theta)]}$$

By further calculation

$$= \frac{\cos^2(45^\circ + \Theta) + \sin^2(45^\circ - \Theta)}{\tan(60^\circ + \Theta)\cot(60^\circ - \Theta)}$$

We know that $\cos(90^\circ - \theta) = \sin \theta$, $\tan(90^\circ - \theta) = \cot \theta$ and $\tan \theta \cot \theta = 1$

So we get

$$= 1/1$$

$$= 1$$

$$= \text{RHS}$$

17. Find the value of A if

(i) $\sin 3A = \cos(A - 6^\circ)$, where $3A$ and $A - 6^\circ$ are acute angles

(ii) $\tan 2A = \cot(A - 18^\circ)$, where $2A$ and $A - 18^\circ$ are acute angles

(iii) If $\sec 2A = \operatorname{cosec}(A - 27^\circ)$ where $2A$ is an acute angle, find the measure of $\angle A$.

Solution:

(i) $\sin 3A = \cos(A - 6^\circ)$, where $3A$ and $A - 6^\circ$ are acute angles

It is given that

$$\sin 3A = \cos(A - 6^\circ)$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

We know that $\cos (90^\circ - \theta) = \sin \theta$

$$\cos (90^\circ - 3A) = \cos (A - 6^\circ)$$

By comparing both

$$90^\circ - 3A = A - 6^\circ$$

By further calculation

$$90^\circ + 6^\circ = A + 3A$$

$$96^\circ = 4A$$

So we get

$$A = 96^\circ/4 = 24^\circ$$

Hence, the value of A is 24° .

$$(ii) \tan 2A = \cot (A - 18^\circ)$$

We know that $\cot (90^\circ - \theta) = \tan \theta$

$$\cot (90^\circ - 2A) = \cot (A - 18^\circ)$$

By comparing both

$$90^\circ - 2A = A - 18^\circ$$

By further calculation

$$90^\circ + 18^\circ = A + 2A$$

So we get

$$3A = 108^\circ$$

$$A = 108^\circ/3 = 36^\circ$$

Hence, the value of A is 36° .

$$(iii) \sec 2A = \operatorname{cosec} (A - 27^\circ)$$

We know that $\operatorname{cosec} (90^\circ - \theta) = \sec \theta$

$$\operatorname{cosec} (90^\circ - 2A) = \sec (A - 27^\circ)$$

By comparing both

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$90^\circ - 2A = A - 27^\circ$$

By further calculation

$$90^\circ + 27^\circ = A + 2A$$

So we get

$$3A = 117^\circ$$

$$A = 117^\circ/3 = 39^\circ$$

Hence, the value of A is 39° .

18. Find the value of θ ($0^\circ < \theta < 90^\circ$) if:

(i) $\cos 63^\circ \sec (90^\circ - \theta) = 1$

(ii) $\tan 35^\circ \cot (90^\circ - \theta) = 1$.

Solution:

(i) $\cos 63^\circ \sec (90^\circ - \theta) = 1$

It can be written as

$$\cos 63^\circ = 1/\sec (90^\circ - \theta)$$

We know that $1/\sec \theta = \cos \theta$

$$\cos 63^\circ = \cos (90^\circ - \theta)$$

By comparing both

$$90^\circ - \theta = 63^\circ$$

By further calculation

$$\theta = 90^\circ - 63^\circ = 27^\circ$$

(ii) $\tan 35^\circ \cot (90^\circ - \theta) = 1$

It can be written as

$$\tan 35^\circ = 1/\cot (90^\circ - \theta)$$

We know that $1/\cot \theta = \tan \theta$

$$\tan 35^\circ = \tan (90^\circ - \theta)$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

By comparing both

$$35^\circ = 90^\circ - \theta$$

By further calculation

$$\theta = 90^\circ - 35^\circ = 55^\circ$$

19. If A, B and C are the interior angles of a $\triangle ABC$, show that

(i) $\cos (A + B)/2 = \sin C/2$

(ii) $\tan (C + A)/2 = \cot B/2$.

Solution:

A, B and C are the interior angles of a $\triangle ABC$

It can be written as

$$\angle A + \angle B + \angle C = 180^\circ$$

Dividing both sides by 2

$$(\angle A + \angle B + \angle C)/2 = 180^\circ/2$$

$$A/2 + B/2 + C/2 = 90^\circ$$

(i) $\cos (A + B)/2 = \sin C/2$

We can write it as

$$(A + B)/2 = 90^\circ - C/2$$

We know that

$$\cos (90^\circ - C/2) = \sin C/2$$

Here $\cos (90^\circ - \theta) = \sin \theta$

$$\sin C/2 = \sin C/2$$

(ii) $\tan (C + A)/2 = \cot B/2$

We know that $(A + C)/2 = 90^\circ - B/2$

$$= \tan (90^\circ - B/2)$$

So we get

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= \cot B/2$$

$$= \text{RHS}$$

Chapter Test

1. Find the values of:

(i) $\sin^2 60^\circ - \cos^2 45^\circ + 3\tan^2 30^\circ$

(ii)

$$\frac{2 \cos^2 45^\circ + 3 \tan^2 30^\circ}{\sqrt{3} \cos 30^\circ + \sin 30^\circ}$$

(iii) $\sec 30^\circ \tan 60^\circ + \sin 45^\circ \operatorname{cosec} 45^\circ + \cos 30^\circ \cot 60^\circ$

Solution:

(i) $\sin^2 60^\circ - \cos^2 45^\circ + 3\tan^2 30^\circ$

$$\begin{aligned} &= \left(\frac{\sqrt{3}}{2}\right)^2 - \left(\frac{1}{\sqrt{2}}\right)^2 + 3\left(\frac{1}{\sqrt{3}}\right)^2 \\ &= \frac{3}{4} - \frac{1}{2} + 3 \times \frac{1}{3} = \frac{3}{4} - \frac{1}{2} + \frac{1}{1} \\ &= \frac{3-2+4}{4} = \frac{7-2}{4} = \frac{5}{4} = 1\frac{1}{4} \end{aligned}$$

Therefore, $\sin^2 60^\circ - \cos^2 45^\circ + 3\tan^2 30^\circ = 1\frac{1}{4}$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\begin{aligned} \text{(ii)} \quad & \frac{2\cos^2 45^\circ + 3\tan^2 30^\circ}{\sqrt{3}\cos 45^\circ + \sin 30^\circ} \\ &= \frac{2\left(\frac{1}{\sqrt{2}}\right)^2 + 3 \times \left(\frac{1}{\sqrt{3}}\right)^2}{\sqrt{3} \times \left(\frac{\sqrt{3}}{2}\right) + \frac{1}{2}} \\ &= \frac{2 \times \frac{1}{2} + 3 \times \frac{1}{3}}{\frac{3}{2} + \frac{1}{2}} = \frac{1+1}{\frac{3+1}{2}} = \frac{2}{\frac{4}{2}} = \frac{2}{2} = 1 \end{aligned}$$

Hence,

$$\frac{2\cos^2 45^\circ + 3\tan^2 30^\circ}{\sqrt{3}\cos 30^\circ + \sin 30^\circ} = 1$$

(iii) $\sec 30^\circ \tan 60^\circ + \sin 45^\circ \operatorname{cosec} 45^\circ + \cos 30^\circ \cot 60^\circ$

$$= \frac{2}{\sqrt{3}} \times \sqrt{3} + \frac{1}{\sqrt{2}} \times \sqrt{2} + \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{3}} = \frac{2}{1} + \frac{1}{1} + \frac{1}{2}$$

$$= 2 + 1 + \frac{1}{2} = 3 + \frac{1}{2} = (6 + 1)/2$$

$$= 7/2 = 3\frac{1}{2}$$

Thus, $\sec 30^\circ \tan 60^\circ + \sin 45^\circ \operatorname{cosec} 45^\circ + \cos 30^\circ \cot 60^\circ = 3\frac{1}{2}$

2. Taking $A = 30^\circ$, verify that

(i) $\cos^4 A - \sin^4 A = \cos 2A$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(ii) $4\cos A \cos (60^\circ - A) \cos (60^\circ + A) = \cos 3A$.

Solution:

(i) $\cos^4 A - \sin^4 A = \cos 2A$

Let's take $A = 30^\circ$

so, we have

L.H.S. = $\cos^4 A - \sin^4 A = \cos^4 30^\circ - \sin^4 30^\circ$

$$\begin{aligned} &= \left(\frac{\sqrt{3}}{2}\right)^4 - \left(\frac{1}{2}\right)^4 \\ &= \frac{\sqrt{3} \times \sqrt{3} \times \sqrt{3} \times \sqrt{3}}{2 \times 2 \times 2 \times 2} - \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} \times \frac{1}{2} = \frac{9}{16} - \frac{1}{16} \\ &= \frac{9-1}{16} = \frac{8}{16} = \frac{1}{2} \end{aligned}$$

Now,

R.H.S. = $\cos 2A = \cos 2(30^\circ) = \frac{1}{2}$

Therefore, L.H.S. = R.H.S. hence verified.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(ii) $4 \cos A \cos (60^\circ - A) \cos (60^\circ + A) = \cos 3A$

Let's take $A = 30^\circ$

$$\text{L.H.S.} = 4 \cos A \cos (60^\circ - A) \cos (60^\circ + A)$$

$$= 4 \cos 30^\circ \cos (60^\circ - 30^\circ) \cos (60^\circ + 30^\circ)$$

$$= 4 \cos 30^\circ \cos 30^\circ \cos 90^\circ$$

$$= 4 \times (\sqrt{3}/2) \times (\sqrt{3}/2) \times 0$$

$$= 0$$

Now,

$$\text{R.H.S.} = \cos 3A$$

$$= \cos (3 \times 30^\circ) = \cos 90^\circ = 0$$

Hence, L.H.S. = R.H.S. hence verified.

3. If $A = 45^\circ$ and $B = 30^\circ$, verify that $\sin A / (\cos A + \sin A + \sin B) = 2/3$

Solution:

Taking,

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\begin{aligned} \text{L.H.S. } & \frac{\sin A}{\cos A + \sin A \sin B} \\ &= \frac{\sin 45^\circ}{\cos 45^\circ + \sin 45^\circ \sin 30^\circ} \\ &= \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2} \times \frac{1}{2}} = \frac{\frac{\sqrt{2}}{2}}{\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{4}} \\ &= \frac{\frac{\sqrt{2}}{2}}{\frac{2\sqrt{2} + \sqrt{2}}{4}} = \frac{\frac{\sqrt{2}}{2}}{\frac{3\sqrt{2}}{4}} \\ &= \frac{\sqrt{2}}{2} \times \frac{4}{3\sqrt{2}} = \frac{2}{3} = \text{R.H.S.} \end{aligned}$$

Hence verified.

4. Taking $A = 60^\circ$ and $B = 30^\circ$, verify that

(i) $\sin(A + B) / \cos A \cos B = \tan A + \tan B$

(ii) $\sin(A - B) / \sin A \sin B = \cot B - \cot A$

Solution:

(i) Here, $A = 60^\circ$ and $B = 30^\circ$

$$\begin{aligned}\text{LHS} &= \frac{\sin(A+B)}{\cos A \cos B} = \frac{\sin(60^\circ + 30^\circ)}{\cos 60^\circ \cos 30^\circ} \\ &= \frac{\sin 90^\circ}{\cos 60^\circ \cos 30^\circ} = \frac{1}{\frac{1}{2} \times \frac{\sqrt{3}}{2}} \\ &= \frac{1 \cdot 4}{\sqrt{3}} = \frac{4}{\sqrt{3}}\end{aligned}$$

Now,

$$\begin{aligned}\text{R.H.S.} &= \tan A + \tan B \\ &= \tan 60^\circ + \tan 30^\circ \\ &= \sqrt{3} + \frac{1}{\sqrt{3}} \\ &= \frac{3+1}{\sqrt{3}} = \frac{4}{\sqrt{3}}\end{aligned}$$

$$\therefore \text{L.H.S.} = \text{R.H.S.}$$

(ii) $A = 60^\circ$, $B = 30^\circ$

$$\begin{aligned}\text{L.H.S.} &= \frac{\sin(A - B)}{\sin A \sin B} = \frac{\sin(60^\circ - 30^\circ)}{\sin 60^\circ \sin 30^\circ} \\ &= \frac{\sin 30^\circ}{\sin 60^\circ \sin 30^\circ} = \frac{1}{\sin 60^\circ} = \frac{1}{\frac{\sqrt{3}}{2}} \\ &= \frac{2}{\sqrt{3}}\end{aligned}$$

$$\begin{aligned}\text{R.H.S.} &= \cot B - \cot A \\ &= \cot 30^\circ - \cot 60^\circ \\ &= \sqrt{3} - \frac{1}{\sqrt{3}} = \frac{3-1}{\sqrt{3}} = \frac{2}{\sqrt{3}}\end{aligned}$$

$\therefore \text{L.H.S.} = \text{R.H.S.}$

5. If $\sqrt{2} \tan 2\theta = \sqrt{6}$ and $0^\circ < 2\theta < 90^\circ$, find the value of $\sin \theta + \sqrt{3} \cos \theta - 2 \tan^2 \theta$.

Solution:

Given,

$$\sqrt{2} \tan 2\theta = \sqrt{6}$$

$$\tan 2\theta = \sqrt{6} / \sqrt{2}$$

$$= \sqrt{3}$$

$$= \tan 60^\circ$$

$$\Rightarrow 2\theta = 60^\circ$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$\theta = 30^\circ$$

Now,

$$\sin \theta + \sqrt{3} \cos \theta - 2 \tan^2 \theta$$

$$= \sin 30^\circ + \sqrt{3} \cos 30^\circ - 2 \tan^2 30^\circ$$

$$= \frac{1}{2} + \sqrt{3} \times \frac{\sqrt{3}}{2} - 2 \left(\frac{1}{\sqrt{3}}\right)^2$$

$$= \frac{1}{2} + \frac{3}{2} - \frac{2}{3}$$

$$= \frac{4}{2} - \frac{2}{3}$$

$$= \frac{(12 - 4)}{6}$$

$$= \frac{8}{6}$$

$$= \frac{4}{3}$$

6. If 3θ is an acute angle, solve the following equations for θ :

(i) $(\operatorname{cosec} 3\theta - 2)(\cot 2\theta - 1) = 0$

(ii) $(\tan \theta - 1)(\operatorname{cosec} 3\theta - 1) = 0$

Solution:

(i) $(\operatorname{cosec} 3\theta - 2)(\cot 2\theta - 1) = 0$

Now, either

$$\operatorname{cosec} 3\theta - 2 \text{ or } \cot 2\theta - 1 = 0$$

$$\Rightarrow \operatorname{cosec} 3\theta = 2 \text{ or } \cot 2\theta = 1$$

So,

$$\operatorname{cosec} 3\theta = \operatorname{cosec} 30^\circ \text{ or } \cot 2\theta = \cot 45^\circ$$

$$\Rightarrow 3\theta = 30^\circ \text{ or } 2\theta = 45^\circ$$

Thus, $\theta = 30^\circ$ or 45° .

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$(ii) (\tan \theta - 1) (\operatorname{cosec} 3\theta - 1) = 0$$

Now, either

$$\tan \theta - 1 = 0 \text{ or } \operatorname{cosec} 3\theta - 1 = 0$$

$$\Rightarrow \tan \theta = 1 \text{ or } \operatorname{cosec} 3\theta = 1$$

So,

$$\tan \theta = \tan 45^\circ \text{ or } \operatorname{cosec} 3\theta = \operatorname{cosec} 90^\circ$$

$$\Rightarrow \theta = 45^\circ \text{ or } 3\theta = 90^\circ \text{ i.e. } \theta = 30^\circ$$

Thus, $\theta = 45^\circ$ or 30° .

7. If $\tan (A + B) = \sqrt{3}$ and $\tan (A - B) = 1$ and A, B ($B < A$) are acute angles, find the values of A and B.

Solution:

$$\text{Given, } \tan (A + B) = \sqrt{3}$$

$$\text{So, } \tan (A + B) = \tan 60^\circ \text{ [Since, } \tan 60^\circ = \sqrt{3}]$$

$$\Rightarrow A + B = 60^\circ \dots\dots\dots (i)$$

Also, given

$$\tan (A - B) = 1$$

$$\text{So, } \tan (A - B) = \tan 45^\circ \text{ [tan } 45^\circ = 1]$$

$$\Rightarrow A - B = 45^\circ \dots\dots\dots (ii)$$

From equation (1) and (2), we get

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$A + B = 60^\circ$$

$$A - B = 45^\circ$$

$$2A = 105^\circ$$

$$A = 52\frac{1}{2}^\circ$$

Now, on substituting the value of A in equation (i), we get

$$52\frac{1}{2}^\circ + B = 60^\circ$$

$$B = 60^\circ - 52\frac{1}{2}^\circ = 7\frac{1}{2}^\circ$$

Therefore, the value of $A = 52\frac{1}{2}^\circ$ and $B = 7\frac{1}{2}^\circ$

8. Without using trigonometrical tables, evaluate the following:

(i) $\sin^2 28^\circ + \sin^2 62^\circ - \tan^2 45^\circ$

(ii)

$$2 \frac{\cos 27^\circ}{\sin 63^\circ} + \frac{\tan 27^\circ}{\cot 63^\circ} + \cos 0^\circ$$

(iii) $\cos 18^\circ \sin 72^\circ + \sin 18^\circ \cos 72^\circ$

(iv) $5 \sin 50^\circ \sec 40^\circ - 3 \cos 59^\circ \operatorname{cosec} 31^\circ$

Solution:

(i) $\sin^2 28^\circ + \sin^2 62^\circ - \tan^2 45^\circ$

$$= \sin^2 28^\circ + \sin^2 (90^\circ - 28^\circ) - \tan^2 45^\circ$$

$$= \sin^2 28^\circ + \cos^2 28^\circ - \tan^2 45^\circ$$

$$= 1 - (1)^2 \quad (\because \sin^2 \theta + \cos^2 \theta = 1 \text{ and } \tan 45^\circ = 1)$$

$$= 1 - 1$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$= 0$$

$$\begin{aligned} (ii) \quad & 2 \frac{\cos 27^\circ}{\sin 63^\circ} + \frac{\tan 27^\circ}{\cot 63^\circ} + \cos 0^\circ \\ &= 2 \frac{\cos 27^\circ}{\sin (90^\circ - 27^\circ)} + \frac{\tan 27^\circ}{\cot (90^\circ - 27^\circ)} + \cos 0^\circ \\ &= 2 \frac{\cos 27^\circ}{\cos 27^\circ} + \frac{\tan 27^\circ}{\tan 27^\circ} + 1 \quad (\because \cos 0^\circ = 1) \end{aligned}$$

$$= 2 \times 1 + 1 + 1$$

$$= 2 + 1 + 1$$

$$= 4$$

$$(iii) \cos 18^\circ \sin 12^\circ + \sin 18^\circ \cos 12^\circ$$

$$= \cos (90^\circ - 12^\circ) \sin 12^\circ + \sin (90^\circ - 12^\circ) \cos 12^\circ$$

$$= \sin 72^\circ \cdot \sin 12^\circ + \cos 12^\circ \cos 12^\circ$$

$$= \sin^2 12^\circ + \cos^2 12^\circ$$

$$= 1 \quad (\because \sin^2 \theta + \cos^2 \theta = 1)$$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(iv) $5 \sin 50^\circ \sec 40^\circ - 3 \cos 59^\circ \operatorname{cosec} 31^\circ$

$$\begin{aligned} &= 5 \frac{\sin 50^\circ}{\cos 40^\circ} - 3 \frac{\cos 59^\circ}{\sin 31^\circ} \\ &= 5 \frac{\sin 50^\circ}{\cos (90^\circ - 50^\circ)} - 3 \frac{\cos 59^\circ}{\sin (90^\circ - 59^\circ)} \\ &= 5 \frac{\sin 50^\circ}{\sin 50^\circ} - 3 \frac{\cos 59^\circ}{\cos 59^\circ} = 5 \times 1 - 3 \times 1 \end{aligned}$$

$$= 5 - 3$$

$$= 2$$

9. Prove that:

$$\frac{\cos (90^\circ - \theta) \sec (90^\circ - \theta) \tan \theta}{\operatorname{cosec} (90^\circ - \theta) \sin (90^\circ - \theta) \cot (90^\circ - \theta)} + \frac{\tan (90^\circ - \theta)}{\cot \theta} = 2$$

Solution:

$$\begin{aligned} \text{L.H.S.} &= \frac{\cos (90^\circ - \theta) \sec (90^\circ - \theta) \tan \theta}{\operatorname{cosec} (90^\circ - \theta) \sin (90^\circ - \theta) \cot (90^\circ - \theta)} + \frac{\tan (90^\circ - \theta)}{\cot \theta} \\ &= \frac{\sin \theta \operatorname{cosec} \theta \tan \theta}{\sec \theta \cos \theta \tan \theta} + \frac{\cot \theta}{\cot \theta} \\ &= \frac{1 \times \tan \theta}{1 \times \tan \theta} + 1 = 1 + 1 = 2 = \text{R.H.S.} \end{aligned}$$

Thus, L.H.S. = R.H.S.

Hence proved.

10. When $0^\circ < A < 90^\circ$, solve the following equations:

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(i) $\sin 3A = \cos 2A$

(ii) $\tan 5A = \cot A$

Solution:

(i) $\sin 3A = \cos 2A$

$$\Rightarrow \sin 3A = \sin (90^\circ - 2A)$$

So,

$$3A = 90^\circ - 2A$$

$$3A + 2A = 90^\circ$$

$$5A = 90^\circ$$

$$\therefore A = 90^\circ / 5 = 18^\circ$$

(ii) $\tan 5A = \cot A$

$$\Rightarrow \tan 5A = \tan (90^\circ - A)$$

So,

$$5A = 90^\circ - A$$

$$5A + A = 90^\circ$$

$$6A = 90^\circ$$

$$\therefore A = 90^\circ / 6 = 15^\circ$$

11. Find the value of θ if

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

(i) $\sin (\theta + 36^\circ) = \cos \theta$, where θ and $\theta + 36^\circ$ are acute angles.

(ii) $\sec 4\theta = \operatorname{cosec} (\theta - 20^\circ)$, where 4θ and $\theta - 20^\circ$ are acute angles.

Solution:

(i) Given, θ and $(\theta + 36^\circ)$ are acute angles

And,

$$\sin (\theta + 36^\circ) = \cos \theta = \sin (90^\circ - \theta) \text{ [As, } \sin (90^\circ - \theta) = \cos \theta \text{]}$$

On comparing, we get

$$\theta + 36^\circ = 90^\circ - \theta$$

$$\theta + \theta = 90^\circ - 36^\circ$$

$$2\theta = 54^\circ$$

$$\theta = 54^\circ/2$$

$$\therefore \theta = 27^\circ$$

(ii) Given, θ and $(\theta - 20^\circ)$ are acute angles

And,

$$\sec 4\theta = \operatorname{cosec} (\theta - 20^\circ)$$

$$\operatorname{cosec} (90^\circ - 4\theta) = \operatorname{cosec} (\theta - 20^\circ) \text{ [Since, } \operatorname{cosec} (90^\circ - \theta) = \sec \theta \text{]}$$

On comparing, we get

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

$$90^\circ - 4\theta = \theta - 20^\circ$$

$$90^\circ + 20^\circ = \theta + 4\theta$$


$$5\theta = 110^\circ$$

$$\theta = 110^\circ/5$$

$$\therefore \theta = 22^\circ$$

12. In the adjoining figure, ABC is right-angled triangle at B and ABD is right angled triangle at A. If $BD \perp AC$ and $BC = 2\sqrt{3}\text{cm}$, find the length of AD.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>



Solution:

Given, $\triangle ABC$ and $\triangle ABD$ are right angled triangles in which $\angle A = 90^\circ$ and $\angle B = 90^\circ$

And,

$BC = 2\sqrt{3}$ cm. AC and BD intersect each other at E at right angle and $\angle CAB = 30^\circ$.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

Now in right $\triangle ABC$, we have

$$\tan \theta = BC/AB$$

$$\Rightarrow \tan 30^\circ = 2\sqrt{3}/AB$$

$$\Rightarrow 1/\sqrt{3} = 2\sqrt{3}/AB$$

$$\Rightarrow AB = 2\sqrt{3} \times \sqrt{3} = 2 \times 3 = 6 \text{ cm.}$$

In $\triangle ABE$, $\angle EAB = 30^\circ$ and $\angle EAB = 90^\circ$

Hence,

$$\angle ABE \text{ or } \angle ABD = 180^\circ - 90^\circ - 30^\circ$$

$$= 60^\circ$$

Now in right $\triangle ABD$, we have

$$\tan 60^\circ = AD/AB$$

$$\Rightarrow \sqrt{3} = AD/6$$

Thus, $AD = 6\sqrt{3} \text{ cm.}$

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>



ML Aggarwal Solutions Class 9 Maths

- Chapter 1- Rational and Irrational Numbers
- Chapter 2- Compound Interest
- Chapter 3- Expansions
- Chapter 4- Factorization
- Chapter 5- Simultaneous Linear Equations
- Chapter 6- Problems on Simultaneous Linear Equations
- Chapter 7- Quadratic Equations
- Chapter 8- Indices
- Chapter 9- Logarithms
- Chapter 10- Triangles
- Chapter 11- Mid Point Theorem
- Chapter 12- Pythagoras Theorem
- Chapter 13- Rectilinear Figures
- Chapter 14- Theorems on Area
- Chapter 15- Circle
- Chapter 16- Mensuration
- Chapter 17- Trigonometric Ratios
- Chapter 18- Trigonometric Ratios and Standard Angles
- Chapter 19- Coordinate Geometry
- Chapter 20- Statistics

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

About ML Aggarwal

M. L. Aggarwal, is an Indian mechanical engineer, educator. His achievements include research in solutions of industrial problems related to fatigue design. Recipient Best Paper award, Manipal Institute of Technology, 2004. Member of TSTE.

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>

About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to **info@indcareer.com**

Postal Address

IndCareer.com
52, Shilpa Nagar,
Somalwada
Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888

Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/ml-aggarwal-solutions-for-class-9-maths-chapter-18-trigonometric-ratios-and-standard-angles/>