


Maharashtra Board Solutions Class 11-Chemistry: Chapter 10- States of Matter

Class 11 - Chapter 10 States of Matter



For any clarifications or questions you can write to info@indcareer.com

Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888, **Website:** <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>



Maharashtra Board Solutions Class 11-Chemistry: Chapter 10- States of Matter

Class 11: Chemistry Chapter 10 solutions. Complete Class 11 Chemistry Chapter 10 Notes.

Maharashtra Board Solutions Class 11-Chemistry: Chapter 10- States of Matter

Maharashtra Board 11th Chemistry Chapter 10, Class 11 Chemistry Chapter 10 solutions

1. Select and write the most appropriate alternatives from the given choices.

Question A.

The unit of viscosity is

- a. dynes
- b. newton
- c. gram
- d. poise

Answer:

- d. poise

Question B.

Which of the following is true for 2 moles of an ideal gas?

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

- a. $PV = nRT$
- b. $PV = RT$
- c. $PV = 2RT$
- d. $PV = T$

Answer:

- c. $PV = 2RT$

Question C.

Intermolecular forces in liquid are

- a. greater than gases
- b. less than solids
- c. both a and b
- d. greater than solids

Answer:

- c. both a and b

Question D.

Interactive forces are in ideal gas.

- a. nil
- b. small
- c. large
- d. same as that of real gases

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

a. nil

Question E.

At constant temperature the pressure of 22.4 dm^3 volume of an ideal gas was increased from 10^5 kPa to 210 kPa , New volume could be-

a. 44.8 dm^3

b. 11.2 dm^3

c. 22.4 dm^3

d. 5.6 dm^3

Answer:

b. 11.2 dm^3

2. Answer in one sentence.

Question A.

Name the term used for mixing of different gases by random molecular motion and frequent collision.

Answer:

The mixing of different gases by random molecular motion and frequent collision is called diffusion.

Question B.

The pressure that each individual gas would exert if it were alone in the container, what do we call it as ?

Answer:

The pressure that each individual gas would exert if it were alone in the container is called as partial pressure.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Question C.

When a gas is heated the particles move more quickly. What is the change in volume of a heated gas if the pressure is kept constant ?

Answer:

The volume of the gas increases on heating if pressure is kept constant.

Question D.

A bubble of methane gas rises from the bottom of the North sea. What will happen to the size of the bubble as it rises to the surface ?

Answer:

According to Boyle's law, the size of the bubble of methane gas increases as it rises to the surface.

Question E.

Convert the following temperatures from degree celcius to kelvin.

- a. -15°C
- b. 25°C
- c. -197°C
- d. 273°C

Answer:

a. $T(\text{K}) = t^{\circ}\text{C} + 273.15$

$\therefore T(\text{K}) = -15^{\circ}\text{C} + 273.15 = 258.15\text{ K}$

b. $T(\text{K}) = t^{\circ}\text{C} + 273.15$

$\therefore T(\text{K}) = 25^{\circ}\text{C} + 273.15 = 298.15\text{ K}$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

c. $T(K) = t^{\circ}C + 273.15$

$\therefore T(K) = -197^{\circ}C + 273.15 = 76.15\text{ K}$

d. $T(K) = t^{\circ}C + 273.15$

$\therefore T(K) = 273^{\circ}C + 273.15 = 546.15\text{ K}$

Question F.

Convert the following pressure values into Pascals.

a. 10 atmosphere

b. 1 kPa.

c. 107000 Nm^{-2}

d. 1 atmosphere

Answer:

a. 10 atmosphere:

$$1\text{ atm} = 101325\text{ Pa}$$

$$\therefore 10\text{ atm} = 1013250\text{ Pa}$$

$$= 1.01325 \times 10^6\text{ Pa}$$

b. 1 kPa:

$$1\text{ kPa} = 1000\text{ Pa}$$

c. 107000 N m^{-2} :

$$1\text{ N m}^{-2} = 1\text{ Pa}$$

$$\therefore 107000\text{ Nm}^{-2} = 107000\text{ Pa}$$

$$= 1.07 \times 10^5\text{ Pa}$$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

d. 1 atmosphere:

$$1 \text{ atm} = 101325 \text{ Pa}$$

$$= 1.01325 \times 10^5 \text{ Pa}$$

Question G.

Convert:

a. Exactly 1.5 atm to pascals

b. 89 kPa to newton per square metre (N m^{-2})

c. 101.325 kPa to bar

d. -100°C to Kelvin

e. 0.124 torr to standard atmosphere

Answer:

a. Exactly 1.5 atm to pascals:

$$1 \text{ atm} = 101325 \text{ Pa}$$

$$\therefore 1.5 \text{ atm} = 1.5 \times 101325$$

$$= 151987.5 \text{ Pa}$$

b. 89 kPa to newton per square metre (N m^{-2}):

$$1 \text{ Pa} = 1 \text{ N m}^{-2} \text{ and } 1 \text{ Pa} = 10^{-3} \text{ kPa}$$

$$\therefore 10^{-3} \text{ kPa} = 1 \text{ N m}^{-2}$$

$$\therefore 89 \text{ kPa} = 1 \times 89 \times 10^{-3} \text{ N m}^{-2} = 89000 \text{ N m}^{-2}$$

c. 101.325 kPa to bar:

$$1 \text{ bar} = 1.0 \times 10^5 \text{ Pa}$$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

$$= 1.0 \times 10^2 \text{ kPa}$$

$$\therefore 100 \text{ kPa} = 1 \text{ bar}$$

$$\therefore 101.325 \text{ kPa} = 1 \times 101.325$$

$$= 1.01325 \text{ bar}$$

d. -100°C to Kelvin:

$$T(\text{K}) = t^\circ\text{C} + 273.15$$

$$\therefore T(\text{K}) = (-100^\circ\text{C}) + 273.15 = 173.15 \text{ K}$$

e. 0.124 torr to standard atmosphere:

$$1 \text{ atm} = 760 \text{ torr}$$

$$\therefore 1 \text{ torr} = \frac{1}{760} \text{ atm}$$

$$\therefore 0.124 \text{ torr} = 0.124 \times \frac{1}{760}$$

$$= 1.632 \times 10^{-4} \text{ atm}$$

Question H.

If density of a gas is measured at constant temperature and pressure then which of the following statement is correct ?

- a. Density is directly proportional to molar mass of the gas.
- b. Greater the density greater is the molar mass of the gas.
- c. If density, temperature and pressure is given ideal gas equation can be used to find molar mass.
- d. All the above statements are correct.


Answer:

- d. All the above statements are correct.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Question I.

Observe the following conversions.



Which of the above reactions is in accordance with the principle of stoichiometry?

Answer:

Both the reactions are in accordance with the principle of stoichiometry.

In the first reaction, both the reactants are completely consumed to form product according to reaction stoichiometry.



In the second reaction, chlorine is the limiting reagent and it is completely consumed to form hydrogen chloride. Excess hydrogen remains unreacted at the end of the reaction. This reaction also follows principle of stoichiometry.

**Question J.**

Hot air balloons float in air because of the low density of the air inside the balloon. Explain this with the help of an appropriate gas law.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>



Answer:

The working of hot air balloon can be explained with the help of Charles' law. According to Charles' law, at constant pressure, the volume of a fixed amount of a gas varies directly with the temperature. This means that as the temperature increases, the air inside the balloon expands and occupies more volume. Thus, hot air inside the balloon is less dense than the surrounding cold air. This causes the hot air balloon to float in air.

3. Answer the following questions.


Question A.

Identify the gas laws from the following diagrams.

Diagrams


Gas laws

a.




.....

b.



.....

c.



.....


<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Answer:

- a. Boyle's law
- b. Charles' law
- c. Avogadro's law [Note: Assuming, T constant]

Question B.

Consider a sample of a gas in a cylinder with a movable piston.




Show diagrammatically the changes in the position of piston, if

- a. Pressure is increased from 1.0 bar to 2.0 bar at constant temperature.
- b. Temperature is decreased from 300 K to 150 K at constant pressure
- c. Temperature is decreased from 400 K to 300 K and pressure is decreased from 4 bar to 3 bar.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>


i.



At constant T, $P \propto \frac{1}{V}$

Since, pressure doubles, volume will become half.

ii.



At constant P, $V \propto T$

Since, temperature becomes half, volume will also become half.

iii.



According to combined gas law,

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$

$$\therefore \frac{4 \times 1}{400} = \frac{3 \times V_2}{300} \quad [\text{Assuming } V_1 = 1 \text{ L}]$$

$$\therefore V_2 = 1 \text{ L}$$

Thus, the volume of the gas remains the same.

Hence, there will be no change in the position of the piston.

Question D.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

List the characteristic physical properties of the gases.

Answer:

Characteristic physical properties of the gases:

- Gases are lighter than solids and liquids (i.e., possess lower density).
- Gases do not possess a fixed volume and shape. They occupy entire space available and take the shape of the container.
- Gas molecules are widely separated and are in continuous, random motion. Therefore, gases exert pressure equally in all directions due to collision of gas molecules, on the walls of the container.
- In case of gases, intermolecular forces are weakest.
- Gases possess the property of diffusion, which is a spontaneous homogeneous inter mixing of two or more gases.
- Gases are highly compressible.

Question E.

Define the terms:

- a. Polarizability
- b. Hydrogen bond
- c. Aqueous tension
- d. Dipole moment

Answer:

- a. Polarizability is defined as the ability of an atom or a molecule to form momentary dipoles, that means, the ability of the atom or molecule to become polar by redistributing its electrons.
- b. The electrostatic force of attraction between positively polarised hydrogen atom of one molecule and a highly electronegative atom (which may be negatively charged) of other molecule is called as hydrogen bond.
- c. The pressure exerted by saturated water vapour is called aqueous tension.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

d. Dipole moment (p) is the product of the magnitude of the charge (Q) and the distance between the centres of positive and negative charge (r). It is designated by a Greek Letter (p) and its unit is Debye (D).

Question F.



Would it be easier to drink water with a straw on the top of the Mount Everest or at the base ? Explain.

Answer:

When you drink through a straw, the pressure inside the straw reduces (as the air is withdrawn by mouth) and the liquid is pushed up to your mouth by atmospheric pressure. Thus, drinking with a straw makes use of pressure difference to force the liquid into your mouth. So, if the pressure difference is less it will be difficult to drink through a straw. On the top of the Mount Everest, atmospheric pressure is very low. Hence, it will be difficult to drink water with a straw on the top of Mount Everest as compared to at the base.

Question G.

Identify type of the intermolecular forces in the following compounds.



Answer:

a. Hydrogen bonding (dipole-dipole attraction) and London dispersion forces

b. London dispersion forces

c. Dipole-dipole interactions and London dispersion forces

d. Dipole-dipole interactions and London dispersion forces

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Question H.

Name the types of intermolecular forces present in Ar, Cl₂, CCl₄ and HNO₃.

Answer:

- a. Ar: London dispersion forces
- b. Cl₂: London dispersion forces
- c. CCl₄: London dispersion forces
- d. HNO₃: Hydrogen bonding (dipole-dipole attraction) and London dispersion forces.

Question I.

Match the pairs of the following :

A

B

- | | |
|-----------------|------------------------------------|
| a. Boyle's law | i. At constant pressure and volume |
| b. Charles' law | ii. At constant temperature |
| | iii. At constant pressure |

Answer:

a – ii,

b – iii

Question J.

Write the statement for :

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

- (a) Boyle's law
- (b) Charles' law

Answer:

a. Statement of Boyle's law: For a fixed mass (number of moles 'n') of a gas at constant temperature, the pressure (P) of the gas is inversely proportional to the volume (V) of the gas.

OR

At constant temperature, the pressure of fixed amount (number of moles) of a gas varies inversely with its volume.

b. Statement for Charles' law:

'At constant pressure, the volume of a fixed mass of a gas is directly proportional to its temperature in Kelvin.

Question K.

Differentiate between Real gas and Ideal gas.

Answer:

Ideal gas:

Real gas:

1. Strictly obeys Boyle's and Charles' law.
 $PVnRT = 1$
2. Molecules are perfectly elastic.
3. No attraction or repulsion between the gas molecules i.e. collision without loss of kinetic energy (K.E.)
4. Actual volume of the gas molecules is negligible as compared to total volume of the gas.
5. Ideal gases cannot be liquified even at low temperature but continues to obey Charles' law and finally occupies zero volume at 0 K.
6. Practically, ideal gas does not exist.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

1. Shows deviation from Boyle's and Charles' law at high pressure and temperature, i.e. obeys Boyle's law and Charles' law at low pressure and high temperature.
 $PVnRT \neq 1$
2. Molecules are not perfectly elastic.
3. Intermolecular attraction is present, hence collision takes place with loss of kinetic energy.
4. Actual volume of individual gas molecule is significant at high pressure and low-temperature.
5. Real gases undergo liquefaction at low' temperature when cooled and compressed.
6. Gases that exist in nature like H_2 , O_2 , CO_2 , N_2 , He, etc. are real gases.

4. Answer the following questions

Question A.

State and write mathematical expression for Dalton's law of partial pressure and explain it with suitable example.

Answer:

i. Statement: The total pressure of a mixture of two or more non-reactive gases is the sum of the partial pressures of the individual gases in the mixture.

ii. Explanation:

Dalton's law can be mathematically expressed as:

$$P_{\text{Total}} = P_1 + P_2 + P_3 \dots (\text{at constant } T \text{ and } V)$$


where, P_{Total} is the total pressure of the mixture and $P_1, P_2, P_3, \dots$ are the partial pressures of individual gases 1, 2, 3, ... in the mixture.

For example, consider two non-reactive gases A and B. On mixing the two gases, pressure exerted by individual gas A in the mixture of both the gases is called partial pressure of gas A (say P_1). Likewise, partial pressure of gas B is P_2 . According to Dalton's law, total pressure of the mixture of gas A and B at constant T and V will be given as:

$$P_{\text{Total}} = P_1 + P_2$$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

iii. Schematic illustration of Dalton's law of partial pressures:



Question B.

Derive an Ideal gas equation. Mention the terms involved in it. Also write how it is utilised to obtain combined gas law.

Answer:

According to Boyle's law,

$$V \propto 1/P \text{ (at constant } T \text{ and } n) \text{(1)}$$

According to Charles' law,

$$V \propto T \text{ (at constant } P \text{ and } n) \text{(2)}$$

According to Avogadro's law,

$$V \propto n \text{ (at constant } P \text{ and } T) \text{(3)}$$

Combining relations (1), (2) and (3), we get

$$V \propto nTP$$

Converting this proportionality into an equation by introducing a constant of proportionality ('R' known as gas constant), we get

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

$$\therefore V = nRT$$

On rearranging the above equation, we get

$$PV = nRT$$

where,

P = Pressure of gas,

V = Volume of gas,

n = number of moles of gas,

R = Gas constant,

T = Absolute temperature of gas.

This is the ideal gas equation or equation of state.

[Note: In the ideal gas equation, R is called gas constant or universal gas constant, whose value is same for all the gases. In this equation, if three variables are known, fourth can be calculated. The equation describes the state of an ideal gas. Hence, it is also called as equation of state.]

The ideal gas equation is written as $PV = nRT$...(1)

On rearranging equation (1), we get,

$$\therefore \frac{PV}{T} = nR = \text{constant}$$


$$\therefore \frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2} \quad \dots(2)$$

The ideal gas equation used in this form is called combined gas law.

Question C.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

With the help of graph answer the following –



At constant temperature,

- Graph shows relation between pressure and volume. Represent the relation mathematically.
- Identify the law.
- Write the statement of law.

Answer:

a. $P \propto \frac{1}{V}$

b. The graph represents Boyle's law as it gives relation between pressure and volume at constant temperature.

c. Statement of Boyle's law: For a fixed mass (number of moles 'n') of a gas at constant temperature, the pressure (P) of the gas is inversely proportional to the volume (V) of the gas.

OR

At constant temperature, the pressure of fixed amount (number of moles) of a gas varies inversely with its volume.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Question D.

Write Postulates of kinetic theory of gases.

Answer:

Postulates of kinetic theory of gases:

- Gases consist of tiny particles (molecules or atoms).
- On an average, gas molecules remain far apart from each other. Therefore, the actual volume of the gas molecules is negligible as compared to the volume of the container. Hence, gases are highly compressible.
- The attractive forces between the gas molecules are negligible at ordinary temperature and pressure. As a result, gas expands to occupy entire volume of the container.
- Gas molecules are in constant random motion and move in all possible directions in straight lines. They collide with each other and with the walls of the container.
- Pressure of the gas is due to the collision of gas molecules with the walls of the container.
- The collisions of the gas molecules are perfectly elastic in nature, which means that the total energy of the gaseous particle remains unchanged after collision.
- The different gas molecules move with different velocities at any instant and hence have different kinetic energies. However, the average kinetic energy of the gas molecules is directly proportional to the absolute temperature.

Question E.

Write a short note on


- a. Vapour pressure.
- b. Surface tension
- c. Viscosity.

Answer:

- a. Vapour pressure:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

- Molecules of liquid have tendency to escape from its surface to form vapour above it. This called evaporation.
- When a liquid is placed in a closed container, the liquid undergoes evaporation and vapours formed undergo condensation.
- At equilibrium, the rate of evaporation and rate of condensation are equal.
- The pressure exerted by the vapour in equilibrium with the liquid is known as saturated vapour pressure or simply vapour pressure.
- Vapour pressure is measured by means of a manometer.
- The most common unit for vapour pressure is torr. $1 \text{ torr} = 1 \text{ mm Hg}$.



[Note: i. The vapour pressure of water is also called aqueous tension.


ii. Water has a vapour pressure of approximately 20 torr at room temperature.]

b. Surface tension:

- The particles in the bulk of liquid are uniformly attracted in all directions and the net force acting on the molecules present inside the bulk is zero.
- But the molecules at the surface experience a net attractive force towards the interior of the liquid, or the forces acting on the molecules on the surface are imbalanced.
- Therefore, liquids have tendency to minimize their surface area and the surface acts as a stretched membrane.
- The force acting per unit length perpendicular to the line drawn on the surface of liquid is called surface tension.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

- Unit: Surface tension is measured in SI unit, N m^{-1} and is denoted by Greek letter ' γ '



c. Viscosity:

i. Liquids (fluids) have tendency to flow.


ii. Viscosity measures the magnitude of internal friction in a liquid or fluid to flow as measured by the force per unit area resisting uniform flow.

iii. Different layers of a liquid flow with different velocity. This called laminar flow. Here, the layers of molecules in the immediate contact of the fixed surface remains stationary. The subsequent layers slip over one another. Strong intermolecular forces obstruct the layers from slipping over one another, resulting in a friction between the layers.

iv. Viscosity is defined as the force of friction between the successive layers of a flowing liquid. It is also the resistance to the flow of a liquid.

v. When a liquid flow through a tube, the central layer has the highest velocity, whereas the layer along the inner wall in the tube remains stationary. This is a result of the viscosity of a liquid. Hence, a velocity gradient exists across the cross-section of the tube.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>



vi. Viscosity is expressed in terms of coefficient of viscosity, ' η ' (Eta). The SI unit of viscosity coefficient is N s m^{-2} (newton second per square meter). In CGS system, the unit (η) is measured in poise.

$$1 \text{ poise} = 1 \text{ g cm}^{-1} \text{ s}^{-1} = 10^{-1} \text{ kg m}^{-1} \text{ s}^{-1}$$

5. Solve the following

Question A.

A balloon is inflated with helium gas at room temperature of 25°C and at 1 bar pressure when its initial volume is 2.27L and allowed to rise in air. As it rises in the air external pressure decreases and the volume of the gas increases till finally it bursts when external pressure is 0.3bar. What is the limit at which volume of the balloon can stay inflated ?

Answer:

Given: P_1 = Initial pressure = 1 bar

V_1 = Initial volume = 2.27 L

P_2 = Final pressure = 0.3 bar

To find: V_2 = Final volume

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Formula: $P_1V_1 = P_2V_2$ (at constant n and T)

Calculation: According to Boyle's law,

$P_1V_1 = P_2V_2$ (at constant n and T)

$$\therefore V_2 = \frac{P_1V_1}{P_2} = \frac{1 \times 2.27 \times 0.3}{0.3} = 7.566667 \text{ L} \approx 7.567 \text{ L}$$

Ans: The balloon can stay inflated below the volume of 7.567 L.

Question B.

A syringe has a volume of 10.0 cm³ at pressure 1 atm. If you plug the end so that no gas can escape and push the plunger down, what must be the final volume to change the pressure to 3.5 atm?



Answer:

Given: P_1 = Initial pressure = 1 atm

V_1 = Initial volume = 10.0 cm³

P_2 = Final pressure = 3.5 atm

To find: V_2 = Final volume

Formula: $P_1V_1 = P_2V_2$ (at constant n and T)

Calculation: According to Boyle's law,

$P_1V_1 = P_2V_2$ (at constant n and T)

$$\therefore V_2 = \frac{P_1V_1}{P_2} = \frac{1 \times 10.0}{3.5}$$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

$$= 2.857 \text{ L}$$

Ans: The final volume of the gas in the syringe is 2.857 L.

Question C.

The volume of a given mass of a gas at 0°C is 2 dm³. Calculate the new volume of the gas at constant pressure when

a. The temperature is increased by 10°C.

b. The temperature is decreased by 10°C.

Answer:

Given: T_1 = Initial temperature = 0 °C = 0 + 273.15 = 273.15 K,

V_1 = Initial volume = 2 dm³

a. T_2 = Final temperature = 273.15 K + 10 = 283.15 K

b. T_2 = Final temperature = 273.15 K – 10 = 263.15 K

To find: V_2 = Final volume in both the cases

Formula: $V_1 T_1 = V_2 T_2$ (at constant n and P)

Calculation: According to Charles' law,

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \text{ (at constant n and P)}$$

$$\therefore \text{ i. } V_2 = \frac{V_1 T_2}{T_1} = \frac{2 \times 283.15}{273.15} = \mathbf{2.073 \text{ dm}^3}$$

$$\text{ ii. } V_2 = \frac{V_1 T_2}{T_1} = \frac{2 \times 263.15}{273.15} = \mathbf{1.927 \text{ dm}^3}$$

Ans: The new volume of a given mass of gas is:

a. 2.073 dm³

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

b. 1.927 dm^3

Question D.

A hot air balloon has a volume of 2800 m^3 at 99°C . What is the volume if the air cools to 80°C ?



Answer:

Given: V_1 = Initial volume = 2800 m^3 , T_1 = Initial temperature = $99^\circ\text{C} = 99 + 273.15 = 372.15 \text{ K}$,

T_2 = Final temperature = $80^\circ\text{C} = 80 + 273.15 \text{ K} = 353.15 \text{ K}$

To find: V_2 = Final volume

Formula: $V_1 T_1 = V_2 T_2$ (at constant n and P)

Calculation: According to Charles' law,

$V_1 T_1 = V_2 T_2$ (at constant n and P)

$$\therefore V_2 = \frac{V_1 T_2}{T_1} = \frac{2800 \times 353.15}{372.15} = 2657 \text{ m}^3$$

Ans: The volume of the balloon when the air cools to 80°C is 2657 m^3 .

Question E.

At 0°C , a gas occupies 22.4 liters. How much hotter must be the gas in celsius and in kelvin to reach volume of 25.0 liters?

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

V_1 = Initial volume of the gas = 22.4 L,

T_1 = Initial temperature = 0 + 273.15 = 273.15 K,

V_2 = Final volume = 25.0 L

To find: T_2 = Final temperature in Celsius and in Kelvin

Formula: $\frac{V_1}{T_1} = \frac{V_2}{T_2}$ (at constant n and P)

Calculation: According to Charles' law,

$$\frac{V_1}{T_1} = \frac{V_2}{T_2} \text{ (at constant n and P)}$$

$$\therefore T_2 = \frac{V_2 \times T_1}{V_1}$$

$$\therefore T_2 = \frac{25.0 \times 273.15}{22.4} = 304.85 \text{ K} \approx \mathbf{304.9 \text{ K}}$$

Converting to Celsius scale:

$$304.85 \text{ K} = 304.85 - 273.15 \text{ }^\circ\text{C} = \mathbf{31.7 \text{ }^\circ\text{C}}$$

Calculation using log table:

$$\frac{25.0 \times 273.15}{22.4}$$

$$= \text{Antilog}_{10} [\log_{10} (25.0) + \log_{10} (273.15) - \log_{10} (22.4)]$$

$$= \text{Antilog}_{10} [1.3979 + 2.4365 - 1.3502]$$

$$= \text{Antilog}_{10} [2.4842] = 304.9$$

Ans: The temperature of the gas must be 31.7 °C or 304.9 K.

Question F.

A 20 L container holds 0.650 mol of He gas at 37 °C at a pressure of 628.3 bar. What will be new pressure inside the container if the volume is reduced to 12 L. The temperature is increased to 177 °C and 1.25 mol of additional He gas was added to it?

Answer:

Given: V_1 = Initial volume = 20 L, n_1 = Initial number of moles = 0.650 mol

P_1 = Initial pressure = 628.3 bar

T_1 = Initial temperature = 37 °C = 37 + 273.15 K = 310.15 K

n_2 = Final number of moles = 0.650 + 1.25 = 1.90 mol, V_2 = Final volume = 12 L

T_2 = Final temperature = 177 °C = 177 + 273.15 K = 450.15 K, $R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$

To find: P_2 = Final pressure

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Formula: $PV = nRT$

Calculation: According to ideal gas equation,

$$P_2V_2 = n_2RT_2.$$

$$\therefore P_2 = \frac{n_2RT_2}{V_2} = \frac{1.90 \times 0.0821 \times 450}{1512} = 5.852 \text{ atm}$$

Ans: The final pressure of the gas is 5.852 atm.

[Note: In the above numerical, converting the pressure value to different units, we get:
 $5.852 \text{ atm} = 4447.52 \text{ torr} = 5.928 \text{ bar}$]

Question G.

Nitrogen gas is filled in a container of volume 2.32 L at 32 °C and 4.7 atm pressure. Calculate the number of moles of the gas.

Answer:

Given: $V = 2.32 \text{ L}$, $P = 4.7 \text{ atm}$, $T = 32^\circ\text{C} = 32 + 273.15 \text{ K} = 305.15 \text{ K}$

$$R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

To find: n = number of moles of gas

Formula: $PV = nRT$

Calculation: According to ideal gas equation,

$$PV = nRT$$

$$\therefore n = \frac{PV}{RT} = \frac{4.7 \times 2.32}{0.0821 \times 305.15} = 0.435 \text{ moles}$$

Ans: Number of moles of N_2 gas in the given volume is 0.435 moles.

Question H.

At 25 °C and 760 mm of Hg pressure a gas occupies 600 mL volume. What will be its pressure at the height where temperature is 10 °C and volume of the gas 640 mL ?

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Answer:

Given: V_1 = Initial volume = 600 mL, V_2 = Final volume = 640 mL

P_1 = Initial pressure = 760 mm Hg

T_1 = Initial temperature = $25^\circ\text{C} = 25 + 273.15\text{ K} = 298.15\text{ K}$

T_2 = Final temperature = $10^\circ\text{C} = 10 + 273.15\text{ K} = 283.15\text{ K}$

P_2 = Final pressure

Formula: $P_1 V_1 T_1 = P_2 V_2 T_2$

Calculation: According to combined gas law.

$$\frac{P_1 V_1}{T_1} = \frac{P_2 V_2}{T_2}$$
$$\therefore P_2 = \frac{P_1 V_1 T_2}{T_1 V_2} = \frac{760 \times 600 \times 283.15}{298.15 \times 640} = 676.654\text{ mm Hg}$$

Ans: The final pressure of a gas is 676.654 mm Hg.

Question I.

A neon-dioxygen mixture contains 70.6 g dioxygen and 167.5g neon. If pressure of the mixture of the gases in the cylinder is 25 bar. What is the partial pressure of dioxygen and neon in the mixture?

Answer:

Given: $m_{\text{O}_2} = 70.6\text{ g}$, $m_{\text{Ne}} = 167.5\text{ g}$,

$P_{\text{Total}} = 25\text{ bar}$

To find: Partial pressure of each gas

Formula: $P_1 = x_1 \times P_{\text{Total}}$

Calculation: Determine number of moles (n) of each gas using formula: $n = \frac{m}{M}$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

$$\therefore n_{\text{O}_2} = \frac{70.6 \text{ g}}{32 \text{ g mol}^{-1}} = 2.206 \text{ mol}$$

$$n_{\text{Ne}} = \frac{167.5 \text{ g}}{20 \text{ g mol}^{-1}} = 8.375 \text{ mol}$$

Determine the mole fraction of each gas using the formula:

$$x = \frac{n}{n_{\text{Total}}}$$

$$x_{\text{O}_2} = \frac{n_{\text{O}_2}}{n_{\text{O}_2} + n_{\text{Ne}}} = \frac{2.206 \text{ mol}}{(2.206 + 8.375) \text{ mol}} = \frac{2.206 \text{ mol}}{10.581 \text{ mol}} = 0.208$$

$$x_{\text{Ne}} = \frac{n_{\text{Ne}}}{n_{\text{Total}}} = \frac{8.375 \text{ mol}}{10.581 \text{ mol}} = 0.792$$

Calculate the partial pressure of each gas:

$$P_{\text{O}_2} = x_{\text{O}_2} \times P_{\text{Total}} = 0.208 \times 25 \text{ bar} = \mathbf{5.2 \text{ bar}}$$

$$P_{\text{Ne}} = x_{\text{Ne}} \times P_{\text{Total}} = 0.792 \times 25 \text{ bar} = \mathbf{19.8 \text{ bar}}$$

Ans: The partial pressure of dioxygen and neon are 5.2 bar and 19.8 bar respectively

Question J.

Calculate the pressure in atm of 1.0 mole of helium in a 2.0 dm³ container at 20.0 °C.

Answer:

Given: n = number of moles = 1.0 mol, V = volume = 2.0 dm³

T = Temperature = 20.0 °C = 20.0 + 273.15 K = 293.15 K

R = 0.0821 L atm K⁻¹ mol⁻¹

To find: Pressure (P)

Formula: $PV = nRT$

Calculation: According to ideal gas equation,

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

$$\begin{aligned} PV &= nRT \\ \therefore P &= \frac{nRT}{V} \\ &= \frac{1.0 \times 0.0821 \times 293.15}{2.0} \\ &= 12.03 \text{ atm} \end{aligned}$$

Calculation using log table:

$$\begin{aligned} &\frac{0.0821 \times 293.15}{2.0} \\ &= \text{Antilog}_{10} [\log_{10} (0.0821) + \log_{10} (293.15) - \log_{10} (2.0)] \\ &= \text{Antilog}_{10} [2.9143 + 2.4672 - 0.3010] \\ &= \text{Antilog}_{10} [1.0805] = 12.03 \end{aligned}$$

Ans: The pressure of the given helium gas is 12.03 atm.

Question K.

Calculate the volume of 1 mole of a gas at exactly 20 °C at a pressure of 101.35 kPa.

Answer:

Given: n = number of moles = 1 mol, P = pressure = 101.35 kPa = 1.00025 atm $\approx$ 1 atm

T = Temperature = 20 °C = 20 + 273.15 K = 293.15 K

R = 0.0821 dm³ atm K⁻¹ mol⁻¹

To find: Volume (V)

Formula: PV = nRT

Calculation: According to ideal gas equation,

PV = nRT

$$\begin{aligned} \therefore V &= \frac{nRT}{P} \\ &= \frac{1 \times 0.0821 \times 293.15}{1} \\ &= 24.07 \text{ dm}^3 \end{aligned}$$

Calculation using log table:

$$\begin{aligned} &0.0821 \times 293.15 \\ &= \text{Antilog}_{10} [\log_{10} (0.0821) + \log_{10} (293.15)] \\ &= \text{Antilog}_{10} [2.9143 + 2.4672] \\ &= \text{Antilog}_{10} [1.3815] = 24.07 \end{aligned}$$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Ans: The volume of the given gas is 24.07 dm^3 .

Question L.

Calculate the number of molecules of methane in 0.50 m^3 of the gas at a pressure of $2.0 \times 10^2 \text{ kPa}$ and a temperature of exactly 300 K .

Answer:

$$V = 0.5 \text{ m}^3, P = 2.0 \times 10^2 \text{ kPa} = 2.0 \times 10^5 \text{ Pa}$$

$$T = 300 \text{ K}, R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$$

To find: Number of molecules of methane gas

$$\text{Formula: } PV = nRT$$

Calculation: According to ideal gas equation,

$$n = \frac{PV}{RT} = \frac{2.0 \times 10^5 \times 0.5}{8.314 \times 300} = 40 \text{ mol}$$

$$\text{Number of molecules} = n \times N_A = 40 \times 6.022 \times 10^{23} = 2.4088 \times 10^{25} \approx 2.409 \times 10^{25}$$

Ans: The number of molecules of methane gas present is 2.409×10^{25} molecules.

Do you know? (Textbook Page No. 140)

Question 1.

Consider three compounds: H_2S , H_2Se and H_2O . Identify which has the highest boiling point. Justify.

Answer:

Among the three compounds H_2O , H_2S and H_2Se , the first one, H_2O has the smallest molecular mass. But it has the highest B.P. of 100°C . B.P. of H_2S is -60°C and of H_2Se is -41.25°C . The extraordinary high B.P. of H_2O is due to very strong hydrogen bonding even though it has the lowest molecular mass.

Can you tell? (Textbook Page No. 140)

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Question i.

What are the various components present in the atmosphere?

Answer:

Various components present in the atmosphere are as follows:

- a. Nitrogen (78%)
- b. Oxygen (21%)
- c. Carbon dioxide and other gases (0.03%)
- d. Inert gases (mainly argon) (0.97%)
- e. Traces of water vapour

Question ii.

Name five elements and five compounds those exist as gases at room temperature.

Answer:

Five elements and five compounds that exist as gases at room temperature are as follows:

No	Element
.	

- a. Nitrogen
- b. Oxygen
- c. Hydrogen
- d. Chlorine

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

e. Argon

No Compounds

.

- a. Carbon dioxide
- b. Carbon monoxide
- c. Nitrogen dioxide
- d. Sulphur dioxide
- e. Methane

Just think. (Textbook Page No. 140)

Question 1.

What is air?

Answer:

- Air is a mixture of various gases.
- One cannot see air but can feel the cool breeze.
- The composition of air by volume is around 78 percent N_2 , 21 percent O_2 and 1 percent other gases including CO_2 .

Use your brainpower. (Textbook Page No. 141)

Question 1.

Find the unit in which car-tyre pressure is measured.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Car-tyre pressure is measured in the units of pounds per square inch (psi) or Newton per metre square (N m^{-2}).

Do you know? (Textbook Page No. 142)

Question 1.

How does a bicycle pump work?

Answer:

A bicycle pump works on Boyle's law. Pushing a bicycle pump squashes the same number of particles into a smaller volume. This squashing means particles hit the walls of the pump more often, increasing the pressure. The increased pressure of a gas can be felt on palm by pushing in the piston of a bicycle pump.

Internet my friend (Textbook Page No. 143)

Question 1.

i. Watch Boyle's law experiment.

ii. Find applications of Boyle's law.

iii. Try to study how Boyle's law helps in 'scuba-diving' i.e., importance of Boyle's law in scuba diving an exhilarating sport.

Answer:

i. Students can refer to 'Boyle's law experiment' on YouTube channel of 'Socratica'.

ii. a. Syringes: When the plunger of a syringe is pulled back out, it causes the volume of the gas inside it to increase due to the reduction of pressure. This creates a vacuum in the syringe, which is constantly trying to adjust the pressure back to normal. However, since the only substance available, such as the blood or medication, is on the other side of the needle, this liquid is sucked into the vacuum, increasing the pressure and decreasing the volume of the gas. When we push the plunger back down, the pressure again increases, lowering the volume inside the syringe, and pushing the fluid out.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

b. Respiration: Boyle's law is essential for the human breathing process. When person breathes in, his/her lung volume increases and the pressure within decreases. Since air always moves from areas of high pressure to areas of low pressure, air is drawn into the lungs. The opposite happens when person exhales. Since the lung volume decreases, the pressure within increases, forcing the air out of the lungs

c. Storage of Gases: Many industries store gases under high pressure. This allows the gas to be stored at a low volume, saving plenty of storage space.

[Note: Students are expected to search more on the internet about various other applications of Boyle's law on their own.]

iii. Importance of Boyle's law in scuba diving:

a. Boyle's law affects scuba diving in many ways.

b. It explains the role of pressure in the aquatic environment.

c. As divers descend, the water pressure surrounding them increases, causing air in their body and equipment to have a smaller volume. As the divers ascend, water pressure decreases, causing their body and equipment to expand to acquire a greater volume.

d. Furthermore, it is crucial that scuba divers never attempt to hold their breath when immersed in water.

e. According to Boyle's law, if divers attempt this when they ascend to a body of water of less pressure, then the air that is trapped in their lungs will over-expand and rupture. This is known as Pulmonary Barotrauma. Thus, it is important for scuba divers to exhale as they ascend because the external pressure increases.

f. Also, if a diver returns to the surface too quickly, dissolved gases in the blood expand and form bubbles, which can get stuck in capillaries and organs (causing the 'bends').

[Note: Students are expected to collect additional information their own.]

Just think. (Textbook Page No. 144)

Question i.

Why does bicycle tyre burst during summer?

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Answer:

- According to Charles' law, at constant pressure, the volume of a fixed amount of a gas varies directly with the temperature. This means that as the temperature increases, the volume also increases.
- During summer, the temperature of the surrounding air is high. Due to the high temperature, the air inside the tyre gets heated. This will increase the volume of the tyres and it will burst.

Question ii.**Why do the hot air balloons fly high?****Answer:**

- According to Charles' law, at constant pressure, gases expand on heating and become less dense. Thus, hot air is less dense than cold air.
- In a hot air balloon, the air inside it is heated by a burner. Upon heating, the air inside the balloon expands and becomes lighter (less dense) than the cooler air on the outside. This causes the hot air balloon to fly high in air.

Just think. (Textbook Page No. 145)**Question 1.****i. List out various real-life examples of Charles' law.****ii. Refer and watch Charles' law experiments.****Answer:****i. Few real-life examples of Charles' law:**

a. Helium balloon: If we fill a helium balloon in a warm or hot room, and then take it into a cold room, it shrinks up and will look like it has lost some of the air inside it. This shows that gases expand on heating and contract on cooling.

b. A bottle of deodorant: If we expose a bottle of deodorant to sunlight and high temperatures, the air molecules inside the bottle will expand which can lead to the bursting of the deodorant bottle. This is another example of Charles' law.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

c. Basketball: You may have noticed that a basketball has less responsive bounce during winter than in summer. This yet another example of Charles' law. When a basketball is inflated, the air pressure inside it is set to a fixed value. As the temperature falls, the volume of the gas inside the ball also decreases proportionally.

[Note: Students are expected to collect additional real-life examples on their own,]

ii. pi [Note: Students can scan the adjacent QR code to visualize Charles' law with the aid of a relevant video.]

Use your brainpower. (Textbook Page No. 146)

Question 1.

Why does the pressure in the automobile tyres change during hot summer or winter season?

Answer:

- According to Gay-Lussac's law, at constant volume, pressure of a fixed amount of a gas is directly proportional to its absolute temperature.
- During hot summer, the temperature of automobile tyre increases faster. Consequently, the air inside the tyre gets heated and the gas molecules starts moving faster.
- As the volume of the tyre remains constant, the pressure inside it increases.
- During winter, the temperature of automobile tyre decreases. Consequently, the air inside the tyre gets cooled and the gas molecules starts moving much slower and the pressure inside the tyre decreases.

Just think. (Textbook Page No. 149)

Question 1.

Do all pure gases and mixtures of gases obey the gas laws?

Answer:

Yes, the gas laws are also applicable to the mixtures of gases. The measurable properties of a mixture of the gases such as pressure, temperature, volume, and amount of gaseous mixture are all related by an ideal gas law.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Just think. (Textbook Page No. 150)

Question 1.

Where is Dalton's law applicable?

Answer:

Air is a gaseous mixture of different gases. Dalton's law is useful for the study of various phenomena in air, for example, air pollution.

Just think. (Textbook Page No. 155)

Question 1.

What makes the oil rise through the wick in an oil lamp?

Answer:

In an oil lamp, oil rises through the wick due to the capillary action. Such a capillary rise of oil is due to the surface tension of oil. The wick acts as a capillary tube. When the wick is placed in oil, the attractive forces between the oil and the inner wall of the capillary (wick) pull the oil up through the wick.



<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

Maharashtra Board Solutions

Class 11 Chemistry

- Chapter 1- Some Basic Concepts of Chemistry
- Chapter 2- Introduction to Analytical Chemistry
- Chapter 3- Basic Analytical Techniques
- Chapter 4- Structure of Atom
- Chapter 5- Chemical Bonding
- Chapter 6- Redox Reactions
- Chapter 7- Modern Periodic Table
- Chapter 8- Elements of Group 1 and 2
- Chapter 9- Elements of Group 13, 14 and 15
- Chapter 10- States of Matter
- Chapter 11- Adsorption and Colloids
- Chapter 12- Chemical Equilibrium
- Chapter 13- Nuclear Chemistry and Radioactivity
- Chapter 14- Basic Principles of Organic Chemistry
- Chapter 15- Hydrocarbons
- Chapter 16- Chemistry in Everyday Life

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

About About Maharashtra State Board (MSBSHSE)

The Maharashtra State Board of Secondary and Higher Secondary Education or MSBSHSE (Marathi: महाराष्ट्र राज्य माध्यमिक आणि उच्च माध्यमिक शिक्षण मंडळ), is an **autonomous and statutory body established in 1965**. The board was amended in the year 1977 under the provisions of the Maharashtra Act No. 41 of 1965.

The Maharashtra State Board of Secondary & Higher Secondary Education (MSBSHSE), Pune is an independent body of the Maharashtra Government. There are more than 1.4 million students that appear in the examination every year. The Maha State Board conducts the board examination twice a year. This board conducts the examination for SSC and HSC.

The Maharashtra government established the Maharashtra State Bureau of Textbook Production and Curriculum Research, also commonly referred to as Ebalbharati, in 1967 to take up the responsibility of providing quality textbooks to students from all classes studying under the Maharashtra State Board. MSBHSE prepares and updates the curriculum to provide holistic development for students. It is designed to tackle the difficulty in understanding the concepts with simple language with simple illustrations. Every year around 10 lakh students are enrolled in schools that are affiliated with the Maharashtra State Board.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

FAQs

Where do I get the Maharashtra State Board Books PDF For free download?

You can download the Maharashtra State Board Books from the eBalbharti official website, i.e. cart.ebalbharati.in or from this article.

Add image

How to Download Maharashtra State Board Books?

Students can get the Maharashtra Books for primary, secondary, and senior secondary classes from here. You can view or download the Maharashtra State Board Books from this page or from the official website for free of cost. Students can follow the detailed steps below to visit the official website and download the e-books for all subjects or a specific subject in different mediums.

Step 1: Visit the official website ebalbharati.in

Step 2: On the top of the screen, select "Download PDF textbooks"

Step 3: From the "Classes" section, select your class.

Step 4: From "Medium", select the medium suitable to you.

Step 5: All Maharashtra board books for class 11th will now be displayed on the right side.

Step 6: Click on the "Download" option to download the PDF book.

Who developed the Maharashtra State board books?

As of now, the MSCERT and Balbharti are responsible for the syllabus and textbooks of Classes 1 to 8, while Classes 9 and 10 are under the Maharashtra State Board of Secondary and Higher Secondary Education (MSBSHSE).

How many state boards are there in Maharashtra?

The Maharashtra State Board of Secondary & Higher Secondary Education, conducts the HSC and SSC Examinations in the state of Maharashtra through its nine Divisional Boards located at Pune, Mumbai, Aurangabad, Nasik, Kolhapur, Amravati, Latur, Nagpur and Ratnagiri.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>

About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to **info@indcareer.com**

Postal Address

IndCareer.com
52, Shilpa Nagar,
Somalwada
Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888

Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-chemistry-chapter-10-states-of-matter/>