


M.Sc sem. 3 Examination
July-sep -2021
Subject : Physical chemistry
Paper CHNN – 602 (New course)

કુલ સમય :- ૬૦ મીનીટ કુલ પ્રશ્નો : ૫૦ વિદ્યાર્થીએ ટીક કરવાના પ્રશ્નો : ૩૫
નોંધ : વિદ્યાર્થીએ માત્ર ૩૫ પ્રશ્નો જ ટીક કરવાના રહેશે. જો વધારે ટીક કરેલ હશે તો પ્રથમ ૩૫ પ્રશ્નો જ ગણતરીમાં લેવામાં આવશે.

1. Amalgum cell shape.....
(A) T-shape b.Z-shape C.A-shape D.H-shape
2. Amalgum cell known as.....
(A) Concentration cell B.Volumetry cell C.Thermal cell D.None of these
3. According to debye-huckelonsagar equation $\Delta m =$ _____
(A) $\Delta m^{\circ} + b\sqrt{c}$ (B) $\Delta m^{\circ} - b\sqrt{c}$ (C) A and B both (D)None of these
- 4.The Conductance of weak electrolytic solution_____with the increase in solution.
(A)Decrease (B)Increase (C)None of these (D)A and B both
- 5.The theoretical explanation of wein effect has been proposed by.....
(A)Debye (B)Onsagar (C)Huckel (D)None of above
- 6.The_____effect is the experimentally observed increase in ionic mobility or conductivity of electrolytes at very high gradient of electrical potential.
(A) Field effect (B) Direct effect (C) wein effect (D) None of these
7. Unit of ionic mobility is.....
(A) Ohm⁻¹ Cm² (B) V⁻¹ S⁻¹ Cm² (C) None of these (D) A and B both
8. Hydroquinone also none as.....
(A)Benzene (B)Phenol (C)Benzene 1,4 diol (D)None of these
9. The name “hydroquinone” was coined by.....
(A) Vander vaals B.Friedrich wolher (C) Thomson (D)Debye-huckel
- 10.Hydroquinone molecular structure.....
(A)  B.  C.  D. 
11. In dissociation constants of polybasic acid the ionization occur in increasing dilution shows that.....
(A) $K_1 < K_2 < K_3$ (B) $K_1 > K_2 > K_3$ (C) $K_1 < K_2 > K_3$ (D) $K_1 > K_2 < K_3$
12. What is protophilic solvents.....
(A)Solvents which have greater tendency the accept protons
(B)Solvents which have lower tendency the accept protons
(C)Solvents which have the tendency to produce protons (D)None of these
- 13.What is monobasic acid.....
(A)Have one replaceable hydrogen atom per one acid molecule (B)Have no stepwise dissociation
(C)Have only one dissociation constant (D)all of these
- 14.Colourimetric determination of dissociation constants function.....
(A) $\frac{CH \cdot CHA}{CA}$ (B) $\frac{CH \cdot CA}{CHA}$ (C) $\frac{CHA \cdot CH}{CA}$ (D) $\frac{CHA \cdot CA}{CH}$
- 15.In colourimetric determination of dissociation constants the values of the function in different solution may be extrapolated to.....
(A)One ionic strength (B)Two ionic strength (C)Three ionic strength (D)Negative ionic strength
- 16.What is aprotic solvent.....
(A)Does not have H atom and attached to electronegative atom
(B)Have H atom and attached to electronegative atom
(C)Have a H atom and attached to electropositive atom
(D)Does not have H atom and attached to electropositive atom
- 17.The electrostatic attraction between two oppositely charged particles.....
(A)Increasing with increasing dielectric constant
(B)Decreasing with decreasing dielectric constant
(C)Decreasing with increasing dielectric constant (D)Both are equal
- 18.The Pka value for ammonium ion acid for in water and in methanol is.....
(A)9.3 and 11.0 (B)11.0 and 9.3 (C)11.3 and 9.0 (D)10.3 and 11.0
- 19.The specific conductance of the water found to be.....
(A) $0.43 \times 10^{-3} \Omega^{-1}$ (B) $0.0043 \times 10^{-3} \Omega^{-1}$ (C) $0.040 \times 10^{-3} \Omega^{-1}$ (D) $0.043 \times 10^{-3} \Omega^{-1}$
- 20.What is isoelectric point.....
(A)When concentration of positive ions is equal to negative ions
(B)When concentration of positive ions is greater than to negative ions
(C)When concentration of negative ions is greater than to positive ions (D)None of these
- 21.A window which can transmit all the incident light without any reflection is called...
a) Polarized window b.Malus window c.Brewster window d.Non-reflecting window

22. What should be the phase difference between the two plane polarized waves vibrating at right angles to each other, to produce Circularly polarized right?
- a) $\pi/6$ b. $\pi/2$ c. $\pi/4$ d. $\pi/3$
23. Which of the following is a Uniaxial crystal?
- a) Borax b. Mica c. Quartz d. Selenite
24. If the phase difference between two rays is $\pi/2$ and the angle of incidence is not equal to $\pi/4$ the emergent light is...
- a) Linearly polarized b. Elliptically polarized c. Circularly polarized d. Non-polarized
25. The velocity of light in water is 1.5×10^8 m/s what is the polarizing angle of incidence?
- a) 47.23° b. 51.02° c. 53.74° d. 63.43°
26. The electro magnetic beam has an intensity 28 W/m^2 and is linearly polarized vertically what is the intensity of the transmitted beam, if the angle of incidence on the Polaroid is 45° vertical?
- a) 10 W/m b. 12 W/m c. 14 W/m d. 16 W/m
27. What should be the thickness of quarter wave plate for u light of wave length $5890 \text{ \AA}$ if $u_e = 1.553$ and $u_0 = 1.544$
- a) $1.33 \times 10^{-3} \text{ cm}$ b. $1.43 \times 10^{-3} \text{ cm}$ c. $1.53 \times 10^{-3} \text{ cm}$ d. $1.63 \times 10^{-3} \text{ cm}$
28. A plate which induces the desired amount of phase difference between two rays is known as...
- a) Polaroid b. Phasor plates c. Retardation plates d. Quartz plates
29. What does the effectiveness of the catalyst in the initiation process of anionic polymerization depend on?
- a) Acidity of the monomer b. Basicity of the catalyst
c. All of the mentioned d. None of the mentioned
30. Which of the following monomers is more prone to form low molecular weight polymer when Polymerized in liquid ammonia?
- a) Acrylonitrile b. Methacrylonitrile c. Styrene d. Methyl methacrylonitrile
31. A linearly polarized wave is always...
- a) In X-Y plane b. In transverse wave c. In longitudinal wave d. In Y-Z plane
32. How the termination in anionic polymerization most probably does take place?
- a) Unimolecular termination b. Bimolecular
b) Termination by solvent transfer d. All of the mentioned
33. Which of the following nature of solvent can exhibit the growth of polymer in anionic polymerization?
- a) Acidic b. Basic c. Neutral d. None of this
34. Which of the following can assist as a catalyst in anionic polymerization?
- a) Sulphuric Acid b. Boron trifluoride c. Grignard reagent d. All of the mentioned
35. Styrene is polymerized using amide catalyst in liquid ammonia, anionically how will the rate of reaction change, if the monomer concentration is doubled?
- a) Doubled b. Halved c. Quadrupled d. No change
36. The auxiliary electrode in polarography is
- a. Dropping mercury b. Mercury pool c. Graphite d. Rotating platinum electrode
37. The electroanalytical technique that involves the measurement of electricity consumed in a Redox reaction of the analyte is.....
- a. Potentiometry b. Conductometry c. Polarography d. Coulometry
38. Electrophoresis was developed by
- (A) Tswett (B) Tsvetberg (C) Tiselius (D) Sanger
39. In polarography the Mercury drop is issued from the capillary by
- (A) a constant Pressure Pump (B) Gravity (C) Reciprocal pump (D) Pneumatic pump
40. Maxima in polarography curve is observed due to...
- A) Migration current B) Convection current C) Residual Current D) Diffusive current
41. The Drop life of mercury in dropping mercury voltammetry is
- A) $< 2 \text{ sec}$ B) $2-5 \text{ sec}$ C) $5-7 \text{ sec}$ D) $5-10 \text{ sec}$
42. The ratio of concentration of titration to the concentration of analyte in amperometric titration ideal should be..
- A) 2 B) 5 C) 10 D) 0.5
43. The current is due to concentration gradient of Electro active species in polarography is
- A) diffusion current B) migration current C) convection current D) residual current
44. Secondary colorimetric titration involves
- A) Addition of titrant B) Titrant Precursor C) use of secondary standards D) constant potential
45. What is diameter DEM?
- A) $10-30 \text{ micrometer}$ B) $20-50 \text{ micrometer}$ C) $0-5 \text{ micrometer}$ D) $20-30 \text{ micrometer}$
46. Which of the following is not a type of electrochemical cell ?
- A) Voltaic cell B) Photovoltaic cell C) Electrolytic cell D) Fuel cell
47. Wien effect is very important at....
- A) Very high electrical fields B) Very low electrical fields C) A & B both D) None
48. Which statement is true for Wien effect ?
- A) Decrease with ionic mobility b. Increase in ionic mobility c. Increase in conductivity d. B & C both
49. The diffusion current in polarography is expressed by the equation
- A) Beer-Lambert law B) Boltzmann equation C) Ilkovic equation D) Bragg's equation
50. Which technique separates charged particles using electrified ?
- A) Hydrolysis B) Electrophoresis C) Protein synthesis D) Protein denaturing