



	Or	
	<p>At cathode: $\text{Ag}^+ + \text{e}^- \longrightarrow \text{Ag}_{(s)}$</p> <p>108g of Ag require 1F</p> <p>\therefore 1.5g of Ag require $\frac{1.5}{108} \text{ F} = \frac{1.5 \times 96500}{108} = 1340.27 \text{ C}$</p> <p>$t = \frac{Q}{i} = \frac{1340.27}{1.5}$</p> <p>=893.51s or 14.89 min</p>	<p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p>
11.	<p>(i) $\text{CH}_3\text{-CH=CH}_2 \xrightarrow{\text{H}_2\text{O} / \text{H}^+} \text{CH}_3\text{-CH(OH)-CH}_3$</p> <p>(ii) </p> <p>(iii) $\text{CH}_3\text{-CH}_2\text{-CH(Br)-CH}_3 \xrightarrow{\text{KOH (Alc)}} \text{CH}_3\text{-CH=CH-CH}_3$</p> <p style="text-align: right;">(or any other correct method)</p> <p style="text-align: center;">Or</p>	<p>1</p> <p>1</p> <p>1</p>
11.	<p>(i) $\text{C}_2\text{H}_5\text{Cl} + \text{NaI} \xrightarrow{\text{Acetone}} \text{C}_2\text{H}_5\text{I} + \text{NaCl}$</p> <p>(ii) </p> <p>(iii) $\text{CH}_3\text{Cl} + \text{KNO}_2 \xrightarrow{\Delta} \text{CH}_3\text{-ONO} + \text{KCl}$</p>	<p>1</p> <p>1</p> <p>1</p>
12.	<p>(i) Due to -I / -R effect of $-\text{NO}_2$ group & +I / +R effect of $-\text{CH}_3$ group or 4-nitrophenoxide ion is more stable than 4-methylphenoxide ion</p> <p>(ii) Due to +R effect of $-\text{OH}$ group in phenol / due to sp^2 hybridization of C-atom in C-OH group in phenol whereas sp^3 hybridization of C-atom in C-OH group in methanol.</p> <p>(iii) $(\text{CH}_3)_3\text{C-Br}$ being a 3° halide prefers to undergo β - elimination on reacting with strong base like NaOCH_3.</p>	<p>1</p> <p>1</p> <p>1</p>
13.	<p>$P_A^0 = 17.5 \text{ mm of Hg}$ $W_B = 15 \text{ g}$ $M_B = 180 \text{ g/mol}$</p> <p>$W_A = 150 \text{ g}$ $P_s = ?$</p> <p>$\frac{P_A^0 - P_s}{P_A^0} = \frac{W_B \times M_A}{M_B \times W_A} \quad \therefore \frac{P_A^0 - P_s}{P_A^0} = \frac{15 \times 18}{180 \times 150} = 0.01$</p> <p>$\frac{P_A^0 - P_s}{P_A^0} = \frac{17.5 - P_s}{17.5} = 0.01$</p> <p>$\therefore p_s = 17.325 \text{ mm of Hg}$</p>	<p>1</p> <p>1</p> <p>1</p>
14.	(i) Non - Stoichiometric defect	1

26.	(ii) Average rate during the interval 30 - 60 sec = $-\frac{\text{Change in concentration}}{\text{Change in time}}$	1/2
	$= -\frac{0.15 - 0.30}{60 - 30}$	1/2
	$= -\frac{-0.15}{30} = 0.005 \text{ mol L}^{-1} \text{ S}^{-1}.$	1
	Or	
	(a) (i) rate increases by 4 times	1
	(ii) 2 nd order	1
	(b) Reaction is 50% completed in 23.1 min i.e. Half-life is 23.1 min	
	$\therefore k = \frac{0.693}{t_{1/2}}$	1/2
	$= \frac{0.693}{23.1} = 0.03 \text{ min}^{-1}$	
	$k = \frac{2.303}{t} \log \frac{[A_0]}{[A]}$	1/2
$0.03 \text{ min}^{-1} = \frac{2.303}{t} \log \frac{100}{25}$	1/2	
$0.03 = \frac{2.303}{t} \log 4$	1/2	
$t = \frac{2.303}{0.03} \times 0.6021 = \frac{1.3866}{0.03} :$		
$= 46.221 \text{ min}$	1	