NCERT

National Council Of Educational Research And Training

NCERT Solutions for 11th Class Maths: Chapter 4-Principle of Mathematical Induction

IndCareer Schools

indCareer indCareer indCareer

NCERT Solutions for 11th Class Maths: Chapter 4-Principle of Mathematical Induction

Class 11: Maths Chapter 4 solutions. Complete Class 11 Maths Chapter 4 Notes.

NCERT Solutions for 11th Class Maths: Chapter 4-Principle of Mathematical Induction

NCERT 11th Maths Chapter 4, class 11 Maths Chapter 4 solutions

Exercise 4.1

https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-4-principle-of-mathematical-induction/

Question 1:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1+3+3^2+...+3^{n-1}=\frac{(3^n-1)}{2}$$

Ans :

Let the given statement be P(n), i.e.,

P(n): $1 + 3 + 3^2 + \ldots + 3^{n-1} = \frac{(3^n - 1)}{2}$

For $n = 1$, we have

P(1): $1 = \frac{(3^1 - 1)}{2} = \frac{3-1}{2} = \frac{2}{2} = 1$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1 + 3 + 3^2 + \ldots + 3^{k-1} = \frac{(3^k - 1)}{2} \quad \ldots(i)$$

We shall now prove that P(k + 1) is true.

Consider

$1 + 3 + 3^2 + \ldots + 3^{k-1} + 3^{(k+1)-1}$

$= (1 + 3 + 3^2 + \ldots + 3^{k-1}) + 3^k$

$= \frac{(3^k - 1)}{2} + 3^k$ [Using (i)]

$= \frac{(3^k - 1) + 2.3^k}{2}$

$= \frac{(1+2)3^k - 1}{2}$

$= \frac{3.3^k - 1}{2}$

$= \frac{3^{k+1} - 1}{2}$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 2:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1^3 + 2^3 + 3^3 + ... + n^3 = \left(\frac{n(n+1)}{2}\right)^2$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): 1^3 + 2^3 + 3^3 + ... + n^3 = \left(\frac{n(n+1)}{2}\right)^2$$

For $n = 1$, we have

$$P(1): 1^3 = 1 = \left(\frac{1(1+1)}{2}\right)^2 = \left(\frac{1.2}{2}\right)^2 = 1^2 = 1$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1^3 + 2^3 + 3^3 + ... + k^3 = \left(\frac{k(k+1)}{2}\right)^2 \qquad ... (i)$$

We shall now prove that P(k + 1) is true.

Consider

$1^3 + 2^3 + 3^3 + \ldots + k^3 + (k + 1)^3$

$= (1^3 + 2^3 + 3^3 + \ldots . + k^3) + (k + 1)^3$

$$=\left(\frac{k(k+1)}{2}\right)^2+(k+1)^3 \qquad [\text{Using (i)}]$$

$$=\frac{k^2(k+1)^2}{4}+(k+1)^3$$

$$=\frac{k^2(k+1)^2+4(k+1)^3}{4}$$

$$=\frac{(k+1)^2\{k^2+4(k+1)\}}{4}$$

$$=\frac{(k+1)^2\{k^2+4k+4\}}{4}$$

$$=\frac{(k+1)^2(k+2)^2}{4}$$

$$=\frac{(k+1)^2(k+1+1)^2}{4}$$

$$=\left(\frac{(k+1)(k+1+1)}{2}\right)^2$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 3:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1+\frac{1}{(1+2)}+\frac{1}{(1+2+3)}+\ldots+\frac{1}{(1+2+3+\ldots n)}=\frac{2n}{(n+1)}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): 1+\frac{1}{1+2}+\frac{1}{1+2+3}+\ldots+\frac{1}{1+2+3+\ldots n}=\frac{2n}{n+1}$$

For n = 1, we have

$$P(1): 1=\frac{2.1}{1+1}=\frac{2}{2}=1 \text{ which is true.}$$

Let P(k) be true for some positive integer k, i.e.,

$$1+\frac{1}{1+2}+\ldots+\frac{1}{1+2+3}+\ldots+\frac{1}{1+2+3+\ldots+k}=\frac{2k}{k+1} \qquad \ldots \text{(i)}$$

We shall now prove that P(k + 1) is true.

Consider

$$1+\frac{1}{1+2}+\frac{1}{1+2+3}+\ldots+\frac{1}{1+2+3+\ldots+k}+\frac{1}{1+2+3+\ldots+k+(k+1)}$$

$$=\left(1+\frac{1}{1+2}+\frac{1}{1+2+3}+\ldots+\frac{1}{1+2+3+\ldots k}\right)+\frac{1}{1+2+3+\ldots+k+(k+1)}$$

$$=\frac{2k}{k+1}+\frac{1}{1+2+3+\ldots+k+(k+1)} \qquad [\text{Using (i)}]$$

$$= \frac{2k}{k+1} + \frac{1}{\left(\frac{(k+1)(k+1+1)}{2}\right)} \quad \left[1+2+3+...+n = \frac{n(n+1)}{2}\right]$$

$$= \frac{2k}{(k+1)} + \frac{2}{(k+1)(k+2)}$$

$$= \frac{2}{(k+1)}\left(k + \frac{1}{k+2}\right)$$

$$= \frac{2}{k+1}\left(\frac{k(k+2)+1}{k+2}\right)$$

$$= \frac{2}{(k+1)}\left(\frac{k^2+2k+1}{k+2}\right)$$

$$= \frac{2\cdot(k+1)^2}{(k+1)(k+2)}$$

$$= \frac{2(k+1)}{(k+2)}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 4:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1.2.3 + 2.3.4 + ... + n(n + 1)(n + 2) = \frac{n(n+1)(n+2)(n+3)}{4}$$

Ans :

Let the given statement be P(n), i.e.,

P(n): $1.2.3 + 2.3.4 + \ldots + n(n+1)(n+2) = \frac{n(n+1)(n+2)(n+3)}{4}$

For $n = 1$, we have

P(1): $1.2.3 = 6 = \frac{1(1+1)(1+2)(1+3)}{4} = \frac{1.2.3.4}{4} = 6$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.2.3 + 2.3.4 + \ldots + k(k+1)(k+2) = \frac{k(k+1)(k+2)(k+3)}{4}$$

We shall now prove that P(k + 1) is true.

Consider

$1.2.3 + 2.3.4 + \ldots + k(k+1)(k+2) + (k+1)(k+2)(k+3)$

$= \{1.2.3 + 2.3.4 + \ldots + k(k+1)(k+2)\} + (k+1)(k+2)(k+3)$

$$= \frac{k(k+1)(k+2)(k+3)}{4} + (k+1)(k+2)(k+3) \qquad \text{[Using (i)]}$$

$$= (k+1)(k+2)(k+3)\left(\frac{k}{4}+1\right)$$

$$= \frac{(k+1)(k+2)(k+3)(k+4)}{4}$$

$$= \frac{(k+1)(k+1+1)(k+1+2)(k+1+3)}{4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 5:

Let the given statement be P(n), i.e.,

$$P(n): 1.3 + 2.3^2 + 3.3^3 + ... + n3^n = \frac{(2n-1)3^{n+1} + 3}{4}$$

For $n = 1$, we have

$$P(1): 1.3 = 3 = \frac{(2.1-1)3^{1+1} + 3}{4} = \frac{3^2 + 3}{4} = \frac{12}{4} = 3$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.3 + 2.3^2 + 3.3^3 + ... + k3^k = \frac{(2k-1)3^{k+1} + 3}{4} \quad ... (i)$$

We shall now prove that P(k + 1) is true.

Consider

$1.3 + 2.3^2 + 3.3^3 + \ldots + k3^k + (k + 1)\ 3^{k+1}$

$= (1.3 + 2.3^2 + 3.3^3 + \ldots + k.3^k) + (k + 1)\ 3^{k+1}$

Ans :

$$= \frac{(2k-1)3^{k+1}+3}{4} + (k+1)3^{k+1} \qquad [\text{Using (i)}]$$

$$= \frac{(2k-1)3^{k+1}+3+4(k+1)3^{k+1}}{4}$$

$$= \frac{3^{k+1}\{2k-1+4(k+1)\}+3}{4}$$

$$= \frac{3^{k+1}\{6k+3\}+3}{4}$$

$$= \frac{3^{k+1}.3\{2k+1\}+3}{4}$$

$$= \frac{3^{(k+1)+1}\{2k+1\}+3}{4}$$

$$= \frac{\{2(k+1)-1\}3^{(k+1)+1}+3}{4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

$$= \frac{(2k-1)3^{k+1}+3}{4} + (k+1)3^{k+1} \qquad \text{[Using (i)]}$$

$$= \frac{(2k-1)3^{k+1}+3+4(k+1)3^{k+1}}{4}$$

$$= \frac{3^{k+1}\{2k-1+4(k+1)\}+3}{4}$$

$$= \frac{3^{k+1}\{6k+3\}+3}{4}$$

$$= \frac{3^{k+1}.3\{2k+1\}+3}{4}$$

$$= \frac{3^{(k+1)+1}\{2k+1\}+3}{4}$$

$$= \frac{\{2(k+1)-1\}3^{(k+1)+1}+3}{4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 6:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1.2+2.3+3.4+...+n.(n+1) = \left[\frac{n(n+1)(n+2)}{3}\right]$$

Ans :

Let the given statement be P(n), i.e.,

P(n): $1.2+2.3+3.4+...+n.(n+1)=\left[\frac{n(n+1)(n+2)}{3}\right]$

For n = 1, we have

P(1): $1.2=2=\frac{1(1+1)(1+2)}{3}=\frac{1.2.3}{3}=2$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.2+2.3+3.4+.....+k.(k+1)=\left[\frac{k(k+1)(k+2)}{3}\right] \quad ...\text{(i)}$$

We shall now prove that P(k + 1) is true.

Consider

$1.2 + 2.3 + 3.4 + \ldots + k.(k + 1) + (k + 1).(k + 2)$

$= [1.2 + 2.3 + 3.4 + \ldots + k.(k + 1)] + (k + 1).(k + 2)$

$$= \frac{k(k+1)(k+2)}{3} + (k+1)(k+2) \qquad \text{[Using (i)]}$$

$$= (k+1)(k+2)\left(\frac{k}{3}+1\right)$$

$$= \frac{(k+1)(k+2)(k+3)}{3}$$

$$= \frac{(k+1)(k+1+1)(k+1+2)}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 7:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1.3 + 3.5 + 5.7 + \ldots + (2n-1)(2n+1) = \frac{n(4n^2+6n-1)}{3}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n):\ 1.3+3.5+5.7+...+(2n-1)(2n+1)=\frac{n\left(4n^2+6n-1\right)}{3}$$

For $n = 1$, we have

$$P(1):1.3=3=\frac{1\left(4.1^2+6.1-1\right)}{3}=\frac{4+6-1}{3}=\frac{9}{3}=3$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.3+3.5+5.7+......+(2k-1)(2k+1)=\frac{k\left(4k^2+6k-1\right)}{3} \qquad ...\text{(i)}$$

We shall now prove that P(k + 1) is true.

Consider

$(1.3 + 3.5 + 5.7 + \ldots + (2k - 1)(2k + 1) + \{2(k + 1) - 1\}\{2(k + 1) + 1\}$

$$= \frac{k(4k^2+6k-1)}{3} + (2k+2-1)(2k+2+1) \qquad \text{[Using (i)]}$$

$$= \frac{k(4k^2+6k-1)}{3} + (2k+1)(2k+3)$$

$$= \frac{k(4k^2+6k-1)}{3} + (4k^2+8k+3)$$

$$= \frac{k(4k^2+6k-1)+3(4k^2+8k+3)}{3}$$

$$= \frac{4k^3+6k^2-k+12k^2+24k+9}{3}$$

$$= \frac{4k^3+18k^2+23k+9}{3}$$

$$= \frac{4k^3+14k^2+9k+4k^2+14k+9}{3}$$

$$= \frac{k(4k^2+14k+9)+1(4k^2+14k+9)}{3}$$

$$= \frac{(k+1)(4k^2+14k+9)}{3}$$

$$= \frac{(k+1)\{4k^2+8k+4+6k+6-1\}}{3}$$

$$= \frac{(k+1)\{4(k^2+2k+1)+6(k+1)-1\}}{3}$$

$$= \frac{(k+1)\{4(k+1)^2+6(k+1)-1\}}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 8:

Prove the following by using the principle of mathematical induction for all $n \in N$: $1.2 + 2.2^2 + 3.2^2 + ... + n.2^n = (n - 1)\, 2^{n+1} + 2$

Ans :

Let the given statement be P(n), i.e.,

P(n): $1.2 + 2.2^2 + 3.2^2 + \ldots + n.2^n = (n - 1)\, 2^{n+1} + 2$

For $n = 1$, we have

P(1): $1.2 = 2 = (1 - 1)\, 2^{1+1} + 2 = 0 + 2 = 2$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$1.2 + 2.2^2 + 3.2^2 + \ldots + k.2^k = (k - 1)\, 2^{k+1} + 2$ … (i)

We shall now prove that P(k + 1) is true.

Consider

$$\left\{1.2 + 2.2^2 + 3.2^3 + \ldots + k.2^k\right\} + (k+1)\cdot 2^{k+1}$$
$$= (k-1)2^{k+1} + 2 + (k+1)2^{k+1}$$
$$= 2^{k+1}\left\{(k-1) + (k+1)\right\} + 2$$
$$= 2^{k+1}.2k + 2$$
$$= k.2^{(k+1)+1} + 2$$
$$= \left\{(k+1) - 1\right\}2^{(k+1)+1} + 2$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 9:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\frac{1}{2}+\frac{1}{4}+\frac{1}{8}+\ldots+\frac{1}{2^n}=1-\frac{1}{2^n}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{2}+\frac{1}{4}+\frac{1}{8}+\ldots+\frac{1}{2^n}=1-\frac{1}{2^n}$$

For $n = 1$, we have

$$P(1): \frac{1}{2}=1-\frac{1}{2^1}=\frac{1}{2}$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{2}+\frac{1}{4}+\frac{1}{8}+\ldots.+\frac{1}{2^k}=1-\frac{1}{2^k} \qquad \ldots \text{(i)}$$

We shall now prove that P(k + 1) is true.

Consider

$$\left(\frac{1}{2}+\frac{1}{4}+\frac{1}{8}+\ldots\ldots.+\frac{1}{2^k}\right)+\frac{1}{2^{k+1}}$$

$$=\left(1-\frac{1}{2^k}\right)+\frac{1}{2^{k+1}} \qquad \text{[Using (i)]}$$

$$= 1 - \frac{1}{2^k} + \frac{1}{2.2^k}$$

$$= 1 - \frac{1}{2^k}\left(1 - \frac{1}{2}\right)$$

$$= 1 - \frac{1}{2^k}\left(\frac{1}{2}\right)$$

$$= 1 - \frac{1}{2^{k+1}}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 10:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \ldots + \frac{1}{(3n-1)(3n+2)} = \frac{n}{(6n+4)}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{2.5}+\frac{1}{5.8}+\frac{1}{8.11}+...+\frac{1}{(3n-1)(3n+2)}=\frac{n}{(6n+4)}$$

For n = 1, we have

$$P(1)=\frac{1}{2.5}=\frac{1}{10}=\frac{1}{6.1+4}=\frac{1}{10}$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{2.5}+\frac{1}{5.8}+\frac{1}{8.11}+...+\frac{1}{(3k-1)(3k+2)}=\frac{k}{6k+4} \qquad ... (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$\frac{1}{2.5}+\frac{1}{5.8}+\frac{1}{8.11}+.......+\frac{1}{(3k-1)(3k+2)}+\frac{1}{\{3(k+1)-1\}\{3(k+1)+2\}}$$

$$=\frac{k}{6k+4}+\frac{1}{(3k+3-1)(3k+3+2)} \qquad [\text{Using (i)}]$$

$$= \frac{k}{6k+4} + \frac{1}{(3k+2)(3k+5)}$$

$$= \frac{k}{2(3k+2)} + \frac{1}{(3k+2)(3k+5)}$$

$$= \frac{1}{(3k+2)}\left(\frac{k}{2} + \frac{1}{3k+5}\right)$$

$$= \frac{1}{(3k+2)}\left(\frac{k(3k+5)+2}{2(3k+5)}\right)$$

$$= \frac{1}{(3k+2)}\left(\frac{3k^2+5k+2}{2(3k+5)}\right)$$

$$= \frac{1}{(3k+2)}\left(\frac{(3k+2)(k+1)}{2(3k+5)}\right)$$

$$= \frac{(k+1)}{6k+10}$$

$$= \frac{(k+1)}{6(k+1)+4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 11:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \ldots + \frac{1}{n(n+1)(n+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{1.2.3}+\frac{1}{2.3.4}+\frac{1}{3.4.5}+\ldots+\frac{1}{n(n+1)(n+2)}=\frac{n(n+3)}{4(n+1)(n+2)}$$

For n = 1, we have

$$P(1): \frac{1}{1\cdot 2\cdot 3}=\frac{1\cdot(1+3)}{4(1+1)(1+2)}=\frac{1\cdot 4}{4\cdot 2\cdot 3}=\frac{1}{1\cdot 2\cdot 3}$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{1\cdot 2\cdot 3}+\frac{1}{2\cdot 3\cdot 4}+\frac{1}{3\cdot 4\cdot 5}+\ldots+\frac{1}{k(k+1)(k+2)}=\frac{k(k+3)}{4(k+1)(k+2)} \quad \ldots \text{(i)}$$

We shall now prove that P(k + 1) is true.

Consider

$$\left[\frac{1}{1\cdot 2\cdot 3}+\frac{1}{2\cdot 3\cdot 4}+\frac{1}{3\cdot 4\cdot 5}+\ldots\ldots+\frac{1}{k(k+1)(k+2)}\right]+\frac{1}{(k+1)(k+2)(k+3)}$$

$$=\frac{k(k+3)}{4(k+1)(k+2)}+\frac{1}{(k+1)(k+2)(k+3)} \quad \text{[Using (i)]}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k(k+3)}{4}+\frac{1}{k+3}\right\}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k(k+3)^2+4}{4(k+3)}\right\}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k(k^2+6k+9)+4}{4(k+3)}\right\}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k^3+6k^2+9k+4}{4(k+3)}\right\}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k^3+2k^2+k+4k^2+8k+4}{4(k+3)}\right\}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k(k^2+2k+1)+4(k^2+2k+1)}{4(k+3)}\right\}$$

$$= \frac{1}{(k+1)(k+2)}\left\{\frac{k(k+1)^2+4(k+1)^2}{4(k+3)}\right\}$$

$$= \frac{(k+1)^2(k+4)}{4(k+1)(k+2)(k+3)}$$

$$= \frac{(k+1)\{(k+1)+3\}}{4\{(k+1)+1\}\{(k+1)+2\}}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 12:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$a+ar+ar^2+\ldots+ar^{n-1} = \frac{a(r^n-1)}{r-1}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): a + ar + ar^2 + ... + ar^{n-1} = \frac{a(r^n - 1)}{r - 1}$$

For n = 1, we have

$$P(1): a = \frac{a(r^1 - 1)}{(r - 1)} = a$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$a + ar + ar^2 + ... + ar^{k-1} = \frac{a(r^k - 1)}{r - 1} \quad ... (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$\{a + ar + ar^2 + ... + ar^{k-1}\} + ar^{(k+1)-1}$$
$$= \frac{a(r^k - 1)}{r - 1} + ar^k \quad [\text{Using (i)}]$$

Let the given statement be P(n), i.e.,

$$P(n): a + ar + ar^2 + \dots + ar^{n-1} = \frac{a(r^n - 1)}{r-1}$$

For $n = 1$, we have

$$P(1): a = \frac{a(r^1 - 1)}{(r-1)} = a$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$a + ar + ar^2 + \dots\dots + ar^{k-1} = \frac{a(r^k - 1)}{r-1} \qquad \dots \text{(i)}$$

We shall now prove that P(k + 1) is true.

Consider

$$\left\{a + ar + ar^2 + \dots\dots + ar^{k-1}\right\} + ar^{(k+1)-1}$$
$$= \frac{a(r^k - 1)}{r-1} + ar^k \qquad \left[\text{Using (i)}\right]$$

Question 13:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)\dots\left(1+\frac{(2n+1)}{n^2}\right) = (n+1)^2$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n):\left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)\ldots\left(1+\frac{(2n+1)}{n^2}\right)=(n+1)^2$$

For $n = 1$, we have

$$P(1):\left(1+\frac{3}{1}\right)=4=(1+1)^2=2^2=4, \text{ which is true.}$$

Let P(k) be true for some positive integer k, i.e.,

$$\left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)\ldots\left(1+\frac{(2k+1)}{k^2}\right)=(k+1)^2 \qquad \ldots (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\left[\left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)\ldots\left(1+\frac{(2k+1)}{k^2}\right)\right]\left\{1+\frac{\{2(k+1)+1\}}{(k+1)^2}\right\}$$

$$=(k+1)^2\left(1+\frac{2(k+1)+1}{(k+1)^2}\right) \qquad [\text{Using}(1)]$$

$$=(k+1)^2\left[\frac{(k+1)^2+2(k+1)+1}{(k+1)^2}\right]$$

$$=(k+1)^2+2(k+1)+1$$

$$=\{(k+1)+1\}^2$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 14:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)\ldots\left(1+\frac{1}{n}\right)=(n+1)$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n):\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)\ldots\left(1+\frac{1}{n}\right)=(n+1)$$

For $n = 1$, we have

$$P(1):\left(1+\frac{1}{1}\right)=2=(1+1)$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k):\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)\ldots\left(1+\frac{1}{k}\right)=(k+1) \quad \ldots (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\left[\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)\dots\left(1+\frac{1}{k}\right)\right]\left(1+\frac{1}{k+1}\right)$$

$$=(k+1)\left(1+\frac{1}{k+1}\right) \qquad \left[\text{Using }(1)\right]$$

$$=(k+1)\left(\frac{(k+1)+1}{(k+1)}\right)$$

$$=(k+1)+1$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 15:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1^2+3^2+5^2+\dots+(2n-1)^2=\frac{n(2n-1)(2n+1)}{3}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n) = 1^2 + 3^2 + 5^2 + ... + (2n-1)^2 = \frac{n(2n-1)(2n+1)}{3}$$

For $n = 1$, we have

$$P(1) = 1^2 = 1 = \frac{1(2.1-1)(2.1+1)}{3} = \frac{1.1.3}{3} = 1, \text{ which is true.}$$

Let P(k) be true for some positive integer k, i.e.,

$$P(k) = 1^2 + 3^2 + 5^2 + ... + (2k-1)^2 = \frac{k(2k-1)(2k+1)}{3} \quad ... (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\left\{1^2 + 3^2 + 5^2 + ... + (2k-1)^2\right\} + \left\{2(k+1) - 1\right\}^2$$

$$= \frac{k(2k-1)(2k+1)}{3} + (2k+2-1)^2 \quad [\text{Using } (1)]$$

$$= \frac{k(2k-1)(2k+1)}{3} + (2k+1)^2$$

$$= \frac{k(2k-1)(2k+1)+3(2k+1)^2}{3}$$

$$= \frac{(2k+1)\{k(2k-1)+3(2k+1)\}}{3}$$

$$= \frac{(2k+1)\{2k^2-k+6k+3\}}{3}$$

$$= \frac{(2k+1)\{2k^2+5k+3\}}{3}$$

$$= \frac{(2k+1)\{2k^2+2k+3k+3\}}{3}$$

$$= \frac{(2k+1)\{2k(k+1)+3(k+1)\}}{3}$$

$$= \frac{(2k+1)(k+1)(2k+3)}{3}$$

$$= \frac{(k+1)\{2(k+1)-1\}\{2(k+1)+1\}}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 16:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\frac{1}{1.4}+\frac{1}{4.7}+\frac{1}{7.10}+...+\frac{1}{(3n-2)(3n+1)}=\frac{n}{(3n+1)}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{1.4}+\frac{1}{4.7}+\frac{1}{7.10}+\ldots+\frac{1}{(3n-2)(3n+1)}=\frac{n}{(3n+1)}$$

For $n=1$, we have

$$P(1)=\frac{1}{1.4}=\frac{1}{3.1+1}=\frac{1}{4}=\frac{1}{1.4}, \text{ which is true.}$$

Let P(k) be true for some positive integer k, i.e.,

$$P(k)=\frac{1}{1.4}+\frac{1}{4.7}+\frac{1}{7.10}+\ldots+\frac{1}{(3k-2)(3k+1)}=\frac{k}{3k+1} \quad \ldots (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\left\{\frac{1}{1.4}+\frac{1}{4.7}+\frac{1}{7.10}+\ldots+\frac{1}{(3k-2)(3k+1)}\right\}+\frac{1}{\{3(k+1)-2\}\{3(k+1)+1\}}$$

$$=\frac{k}{3k+1}+\frac{1}{(3k+1)(3k+4)} \quad \left[\text{Using } (1)\right]$$

$$=\frac{1}{(3k+1)}\left\{k+\frac{1}{(3k+4)}\right\}$$

$$=\frac{1}{(3k+1)}\left\{\frac{k(3k+4)+1}{(3k+4)}\right\}$$

$$=\frac{1}{(3k+1)}\left\{\frac{3k^2+4k+1}{(3k+4)}\right\}$$

$$=\frac{1}{(3k+1)}\left\{\frac{3k^2+3k+k+1}{(3k+4)}\right\}$$

$$=\frac{(3k+1)(k+1)}{(3k+1)(3k+4)}$$

$$=\frac{(k+1)}{3(k+1)+1}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n

Question 17:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$\frac{1}{3.5}+\frac{1}{5.7}+\frac{1}{7.9}+\ldots+\frac{1}{(2n+1)(2n+3)}=\frac{n}{3(2n+3)}$$

Ans :

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + ... + \frac{1}{(2n+1)(2n+3)} = \frac{n}{3(2n+3)}$$

For $n = 1$, we have

$$P(1): \frac{1}{3.5} = \frac{1}{3(2.1+3)} = \frac{1}{3.5}$$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k): \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + ... + \frac{1}{(2k+1)(2k+3)} = \frac{k}{3(2k+3)} \quad ... (1)$$

We shall now prove that P(k + 1) is true.

$$\left[\frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + ... + \frac{1}{(2k+1)(2k+3)}\right] + \frac{1}{\{2(k+1)+1\}\{2(k+1)+3\}}$$

$$= \frac{k}{3(2k+3)} + \frac{1}{(2k+3)(2k+5)} \quad [\text{Using } (1)]$$

$$= \frac{1}{(2k+3)}\left[\frac{k}{3} + \frac{1}{(2k+5)}\right]$$

$$= \frac{1}{(2k+3)}\left[\frac{k(2k+5)+3}{3(2k+5)}\right]$$

$$= \frac{1}{(2k+3)}\left[\frac{2k^2+5k+3}{3(2k+5)}\right]$$

$$= \frac{1}{(2k+3)}\left[\frac{2k^2+2k+3k+3}{3(2k+5)}\right]$$

$$= \frac{1}{(2k+3)}\left[\frac{2k(k+1)+3(k+1)}{3(2k+5)}\right]$$

$$= \frac{(k+1)(2k+3)}{3(2k+3)(2k+5)}$$

$$= \frac{(k+1)}{3\{2(k+1)+3\}}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 18:

Prove the following by using the principle of mathematical induction for all $n \in N$:

$$1+2+3+...+n < \frac{1}{8}(2n+1)^2$$

Ans :

Let P(k) be true for some positive integer k, i.e.,

$$1+2+...+k<\frac{1}{8}(2k+1)^2 \qquad ... (1)$$

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider inequality (1)

$$1+2+...+k<\frac{1}{8}(2k+1)^2$$

Adding (k+1) on both the sides of the inequality, we have,

$$(1+2+...+k)+(k+1)<\frac{1}{8}(2k+1)^2+(k+1)$$

$$(1+2+...+k)+(k+1)<\frac{1}{8}\left\{(2k+1)^2+8(k+1)\right\}$$

$$(1+2+...+k)+(k+1)<\frac{1}{8}\left\{4k^2+4k+1+8k+8\right\}$$

$$(1+2+...+k)+(k+1)<\frac{1}{8}\left\{4k^2+12k+9\right\}$$

$$(1+2+...+k)+(k+1)<\frac{1}{8}(2k+3)^2$$

$$(1+2+...+k)+(k+1)<\frac{1}{8}\left\{2(k+1)+1\right\}^2$$

Hence, $(1+2+3+...+k)+(k+1)<\frac{1}{8}(2k+1)^2+(k+1)$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 19:

Prove the following by using the principle of mathematical induction for all $n \in N$: $10^{2n-1} + 1$ is divisible by 11.

Ans :

Let the given statement be P(n), i.e.,

P(n): $10^{2n-1}+1$ is divisible by 11.

It can be observed that P(n) is true for $n = 1$ since P(1) = $10^{2.1-1}+1 = 11$, which is divisible by 11.

Let P(k) be true for some positive integer k, i.e.,

$10^{2k-1}+1$ is divisible by 11.

$\therefore 10^{2k-1}+1 = 11m$, where $m \in \mathbf{N}$ … (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$10^{2(k+1)-1}+1$$
$$=10^{2k+2-1}+1$$
$$=10^{2k+1}+1$$
$$=10^2\left(10^{2k-1}+1-1\right)+1$$
$$=10^2\left(10^{2k-1}+1\right)-10^2+1$$
$$=10^2.11m-100+1 \qquad \left[\text{Using (1)}\right]$$

$$=100\times 11m-99$$
$$=11(100m-9)$$
$$=11r,\ \text{where } r = (100m-9) \text{ is some natural number}$$

Therefore, $10^{2(k+1)-1}+1$ is divisible by 11.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 20:

Prove the following by using the principle of mathematical induction for all $n \in N$: $x^{2n} - y^{2n}$ is divisible by $x + y$.

Ans :

Let the given statement be P(n), i.e.,

P(n): $x^{2n} - y^{2n}$ is divisible by $x + y$.

It can be observed that P(n) is true for $n = 1$.

This is so because $x^{2 \times 1} - y^{2 \times 1} = x^2 - y^2 = (x + y)(x - y)$ is divisible by $(x + y)$.

Let P(k) be true for some positive integer k, i.e.,

$x^{2k} - y^{2k}$ is divisible by $x + y$.

$\therefore x^{2k} - y^{2k} = m(x + y)$, where $m \in \mathbf{N}$ ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$x^{2(k+1)} - y^{2(k+1)}$$
$$= x^{2k} \cdot x^2 - y^{2k} \cdot y^2$$
$$= x^2\left(x^{2k} - y^{2k} + y^{2k}\right) - y^{2k} \cdot y^2$$
$$= x^2\left\{m(x+y) + y^{2k}\right\} - y^{2k} \cdot y^2 \qquad [\text{Using } (1)]$$
$$= m(x+y)x^2 + y^{2k} \cdot x^2 - y^{2k} \cdot y^2$$
$$= m(x+y)x^2 + y^{2k}\left(x^2 - y^2\right)$$
$$= m(x+y)x^2 + y^{2k}(x+y)(x-y)$$
$$= (x+y)\left\{mx^2 + y^{2k}(x-y)\right\}, \text{ which is a factor of } (x+y).$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 21:

Prove the following by using the principle of mathematical induction for all $n \in N$: $3^{2n+2} - 8n - 9$ is divisible by 8.

Ans :

Let the given statement be P(n), i.e.,

P(n): $3^{2n+2} - 8n - 9$ is divisible by 8.

It can be observed that P(n) is true for $n = 1$ since $3^{2 \times 1 + 2} - 8 \times 1 - 9 = 64$, which is divisible by 8.

Let P(k) be true for some positive integer k, i.e.,

$3^{2k+2} - 8k - 9$ is divisible by 8.

$\therefore 3^{2k+2} - 8k - 9 = 8m$; where $m \in \mathbf{N}$ ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$3^{2(k+1)+2} - 8(k+1) - 9$$
$$= 3^{2k+2} \cdot 3^2 - 8k - 8 - 9$$
$$= 3^2\left(3^{2k+2} - 8k - 9 + 8k + 9\right) - 8k - 17$$
$$= 3^2\left(3^{2k+2} - 8k - 9\right) + 3^2\left(8k + 9\right) - 8k - 17$$
$$= 9.8m + 9\left(8k + 9\right) - 8k - 17$$
$$= 9.8m + 72k + 81 - 8k - 17$$
$$= 9.8m + 64k + 64$$
$$= 8\left(9m + 8k + 8\right)$$

Therefore, $3^{2(k+1)+2} - 8(k+1) - 9$ is divisible by 8.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 22:

Prove the following by using the principle of mathematical induction for all $n \in N$: $41^n - 14^n$ is a multiple of 27.

Ans :

Let the given statement be P(n), i.e.,

P(n): $41^n - 14^n$ is a multiple of 27.

It can be observed that P(n) is true for $n = 1$ since $41^1 - 14^1 = 27$, which is a multiple of 27.

Let P(k) be true for some positive integer k, i.e.,

$41^k - 14^k$ is a multiple of 27

$\therefore 41^k - 14^k = 27m$, where $m \in \mathbf{N}$ … (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$
\begin{aligned}
&41^{k+1} - 14^{k+1} \\
&= 41^k \cdot 41 - 14^k \cdot 14 \\
&= 41\left(41^k - 14^k + 14^k\right) - 14^k \cdot 14 \\
&= 41\left(41^k - 14^k\right) + 41.14^k - 14^k \cdot 14 \\
&= 41.27m + 14^k\left(41 - 14\right) \\
&= 41.27m + 27.14^k \\
&= 27\left(41m - 14^k\right) \\
&= 27 \times r, \text{ where } r = \left(41m - 14^k\right) \text{ is a natural number}
\end{aligned}
$$

Therefore, $41^{k+1} - 14^{k+1}$ is a multiple of 27.

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Question 23:

Prove the following by using the principle of mathematical induction for all $n \in \mathbf{N}$:

$(2n+7) < (n+3)^2$

Ans :

Let the given statement be P(n), i.e.,

P(n): $(2n+7) < (n+3)^2$

It can be observed that P(n) is true for $n = 1$ since $2.1 + 7 = 9 < (1 + 3)^2 = 16$, which is true.

Let P(k) be true for some positive integer k, i.e.,

$(2k+7) < (k+3)^2$ … (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$\{2(k+1)+7\} = (2k+7)+2$$

$$\therefore \{2(k+1)+7\} = (2k+7)+2 < (k+3)^2 + 2 \qquad [\text{using } (1)]$$

$$2(k+1)+7 < k^2+6k+9+2$$

$$2(k+1)+7 < k^2+6k+11$$

$$\text{Now, } k^2+6k+11 < k^2+8k+16$$

$$\therefore 2(k+1)+7 < (k+4)^2$$

$$2(k+1)+7 < \{(k+1)+3\}^2$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Chapterwise NCERT Solutions for Class 11 Maths :

About NCERT

The National Council of Educational Research and Training is an autonomous organization of the Government of India which was established in 1961 as a literary, scientific, and charitable Society under the Societies Registration Act. The major objectives of NCERT and its constituent units are to: undertake, promote and coordinate research in areas related to school education; prepare and publish model textbooks, supplementary material, newsletters, journals and develop educational kits, multimedia digital materials, etc.

Organise pre-service and in-service training of teachers; develop and disseminate innovative educational techniques and practices;collaborate and network with state educational departments, universities, NGOs and other educational institutions; act as a clearing house for ideas and information in matters related to school education; and act as a nodal agency for achieving the goals of Universalisation of Elementary Education. In addition to research, development, training, extension, publication and dissemination activities, NCERT is an implementation agency for bilateral cultural exchange programmes with other countries in the field of school education.Its headquarters are located at Sri Aurobindo Marg in New Delhi. Visit the Official NCERT website to learn more.