



NCERT Solutions for 11th Class Maths: Chapter 8-Binomial Theorem



indCareer



indCareer



indCareer

NCERT Solutions for 11th Class Maths: Chapter 8-Binomial Theorem

Class 11: Maths Chapter 8 solutions. Complete Class 11 Maths Chapter 8 Notes.

NCERT Solutions for 11th Class Maths: Chapter 8-Binomial Theorem

NCERT 11th Maths Chapter 8, class 11 Maths Chapter 8 solutions

Exercise.8.1

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Question-1

Expand the expression $(1-2x)^5$

Ans.

By using Binomial Theorem, the expression $(1-2x)^5$ can be expanded as

$$\begin{aligned}(1-2x)^5 &= {}^5C_0(1)^5 - {}^5C_1(1)^4(2x) + {}^5C_2(1)^3(2x)^2 - {}^5C_3(1)^2(2x)^3 + {}^5C_4(1)^1(2x)^4 - {}^5C_5(2x)^5 \\&= 1 - 5(2x) + 10(4x^2) - 10(8x^3) + 5(16x^4) - (32x^5) \\&= 1 - 10x + 40x^2 - 80x^3 + 80x^4 - 32x^5\end{aligned}$$

Question-2

Expand the expression $\left(\frac{2}{x} - \frac{x}{2}\right)^5$

Ans.

By using Binomial Theorem, the expression $\left(\frac{2}{x} - \frac{x}{2}\right)^5$ can be expanded as

$$\begin{aligned}\left(\frac{2}{x} - \frac{x}{2}\right)^5 &= {}^5C_0\left(\frac{2}{x}\right)^5 - {}^5C_1\left(\frac{2}{x}\right)^4\left(\frac{x}{2}\right) + {}^5C_2\left(\frac{2}{x}\right)^3\left(\frac{x}{2}\right)^2 \\&\quad - {}^5C_3\left(\frac{2}{x}\right)^2\left(\frac{x}{2}\right)^3 + {}^5C_4\left(\frac{2}{x}\right)\left(\frac{x}{2}\right)^4 - {}^5C_5\left(\frac{x}{2}\right)^5 \\&= \frac{32}{x^5} - 5\left(\frac{16}{x^4}\right)\left(\frac{x}{2}\right) + 10\left(\frac{8}{x^3}\right)\left(\frac{x^2}{4}\right) - 10\left(\frac{4}{x^2}\right)\left(\frac{x^3}{8}\right) + 5\left(\frac{2}{x}\right)\left(\frac{x^4}{16}\right) - \frac{x^5}{32} \\&= \frac{32}{x^5} - \frac{40}{x^3} + \frac{20}{x} - 5x + \frac{5}{8}x^3 - \frac{x^5}{32}\end{aligned}$$

Question-3

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Expand the expression $(2x - 3)^6$

Ans.

By using Binomial Theorem, the expression $(2x - 3)^6$ can be expanded as

$$\begin{aligned}(2x - 3)^6 &= {}^6C_0 (2x)^6 - {}^6C_1 (2x)^5 (3) + {}^6C_2 (2x)^4 (3)^2 - {}^6C_3 (2x)^3 (3)^3 \\&\quad + {}^6C_4 (2x)^2 (3)^4 - {}^6C_5 (2x)(3)^5 + {}^6C_6 (3)^6 \\&= 64x^6 - 6(32x^5)(3) + 15(16x^4)(9) - 20(8x^3)(27) \\&\quad + 15(4x^2)(81) - 6(2x)(243) + 729 \\&= 64x^6 - 576x^5 + 2160x^4 - 4320x^3 + 4860x^2 - 2916x + 729\end{aligned}$$

Question-4

Expand the expression $\left(\frac{x}{3} + \frac{1}{x}\right)^5$

Ans.

By using Binomial Theorem, the expression $\left(\frac{x}{3} + \frac{1}{x}\right)^5$ can be expanded as

$$\begin{aligned}\left(\frac{x}{3} + \frac{1}{x}\right)^5 &= {}^5C_0 \left(\frac{x}{3}\right)^5 + {}^5C_1 \left(\frac{x}{3}\right)^4 \left(\frac{1}{x}\right) + {}^5C_2 \left(\frac{x}{3}\right)^3 \left(\frac{1}{x}\right)^2 \\&\quad + {}^5C_3 \left(\frac{x}{3}\right)^2 \left(\frac{1}{x}\right)^3 + {}^5C_4 \left(\frac{x}{3}\right) \left(\frac{1}{x}\right)^4 + {}^5C_5 \left(\frac{1}{x}\right)^5 \\&= \frac{x^5}{243} + 5\left(\frac{x^4}{81}\right)\left(\frac{1}{x}\right) + 10\left(\frac{x^3}{27}\right)\left(\frac{1}{x^2}\right) + 10\left(\frac{x^2}{9}\right)\left(\frac{1}{x^3}\right) + 5\left(\frac{x}{3}\right)\left(\frac{1}{x^4}\right) + \frac{1}{x^5} \\&= \frac{x^5}{243} + \frac{5x^3}{81} + \frac{10x}{27} + \frac{10}{9x} + \frac{5}{3x^3} + \frac{1}{x^5}\end{aligned}$$

Question-5

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>



Expand $\left(x + \frac{1}{x}\right)^6$

Ans.

By using Binomial Theorem, the expression $\left(x + \frac{1}{x}\right)^6$ can be expanded as

$$\begin{aligned}\left(x + \frac{1}{x}\right)^6 &= {}^6C_0(x)^6 + {}^6C_1(x)^5\left(\frac{1}{x}\right) + {}^6C_2(x)^4\left(\frac{1}{x}\right)^2 \\ &\quad + {}^6C_3(x)^3\left(\frac{1}{x}\right)^3 + {}^6C_4(x)^2\left(\frac{1}{x}\right)^4 + {}^6C_5(x)\left(\frac{1}{x}\right)^5 + {}^6C_6\left(\frac{1}{x}\right)^6 \\ &= x^6 + 6(x)^5\left(\frac{1}{x}\right) + 15(x)^4\left(\frac{1}{x^2}\right) + 20(x)^3\left(\frac{1}{x^3}\right) + 15(x)^2\left(\frac{1}{x^4}\right) + 6(x)\left(\frac{1}{x^5}\right) + \frac{1}{x^6} \\ &= x^6 + 6x^4 + 15x^2 + 20 + \frac{15}{x^2} + \frac{6}{x^4} + \frac{1}{x^6}\end{aligned}$$

Question-6

Using Binomial Theorem, evaluate $(96)^3$

Ans.

96 can be expressed as the sum or difference of two numbers whose powers are easier to calculate and then, binomial theorem can be applied.

It can be written that, $96 = 100 - 4$

$$\begin{aligned}\therefore (96)^3 &= (100 - 4)^3 \\ &= {}^3C_0(100)^3 - {}^3C_1(100)^2(4) + {}^3C_2(100)(4)^2 - {}^3C_3(4)^3 \\ &= (100)^3 - 3(100)^2(4) + 3(100)(4)^2 - (4)^3 \\ &= 1000000 - 120000 + 4800 - 64 \\ &= 884736\end{aligned}$$

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Question-7

Using Binomial Theorem, evaluate $(102)^5$

Ans.

102 can be expressed as the sum or difference of two numbers whose powers are easier to calculate and then, Binomial Theorem can be applied.

It can be written that, $102 = 100 + 2$

$$\begin{aligned}\therefore (102)^5 &= (100+2)^5 \\ &= {}^5C_0(100)^5 + {}^5C_1(100)^4(2) + {}^5C_2(100)^3(2)^2 + {}^5C_3(100)^2(2)^3 \\ &\quad + {}^5C_4(100)(2)^4 + {}^5C_5(2)^5 \\ &= (100)^5 + 5(100)^4(2) + 10(100)^3(2)^2 + 10(100)^2(2)^3 + 5(100)(2)^4 + (2)^5 \\ &= 10000000000 + 1000000000 + 40000000 + 800000 + 8000 + 32 \\ &= 11040808032\end{aligned}$$

Question-8

Using Binomial Theorem, evaluate $(101)^4$

Ans.

101 can be expressed as the sum or difference of two numbers whose powers are easier to calculate and then, Binomial Theorem can be applied.

It can be written that, $101 = 100 + 1$

$$\begin{aligned}\therefore (101)^4 &= (100+1)^4 \\ &= {}^4C_0(100)^4 + {}^4C_1(100)^3(1) + {}^4C_2(100)^2(1)^2 + {}^4C_3(100)(1)^3 + {}^4C_4(1)^4 \\ &= (100)^4 + 4(100)^3 + 6(100)^2 + 4(100) + (1)^4 \\ &= 100000000 + 4000000 + 60000 + 400 + 1 \\ &= 104060401\end{aligned}$$

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Question-9

Using Binomial Theorem, evaluate $(99)^5$

Ans.

99 can be written as the sum or difference of two numbers whose powers are easier to calculate and then, Binomial Theorem can be applied.

It can be written that, $99 = 100 - 1$

$$\begin{aligned}\therefore (99)^5 &= (100 - 1)^5 \\ &= {}^5C_0(100)^5 - {}^5C_1(100)^4(1) + {}^5C_2(100)^3(1)^2 - {}^5C_3(100)^2(1)^3 \\ &\quad + {}^5C_4(100)(1)^4 - {}^5C_5(1)^5 \\ &= (100)^5 - 5(100)^4 + 10(100)^3 - 10(100)^2 + 5(100) - 1 \\ &= 10000000000 - 5000000000 + 1000000000 - 100000 + 500 - 1 \\ &= 10010000500 - 500100001 \\ &= 9509900499\end{aligned}$$

Question-10

Using Binomial Theorem, indicate which number is larger $(1.1)^{10000}$ or 1000.

Ans.

By splitting 1.1 and then applying Binomial Theorem, the first few terms of $(1.1)^{10000}$ can be obtained as

$$\begin{aligned}(1.1)^{10000} &= (1 + 0.1)^{10000} \\ &= {}^{10000}C_0 + {}^{10000}C_1(1.1) + \text{Other positive terms} \\ &= 1 + 10000 \times 1.1 + \text{Other positive terms} \\ &= 1 + 11000 + \text{Other positive terms} \\ &> 1000\end{aligned}$$

Hence, $(1.1)^{10000} > 1000$

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Question-11

Find $(a + b)^4 - (a - b)^4$. Hence, evaluate $(\sqrt{3} + \sqrt{2})^4 - (\sqrt{3} - \sqrt{2})^4$.

Ans.

Using Binomial Theorem, the expressions, $(a + b)^4$ and $(a - b)^4$, can be expanded as

$$\begin{aligned}(a + b)^4 &= {}^4C_0a^4 + {}^4C_1a^3b + {}^4C_2a^2b^2 + {}^4C_3ab^3 + {}^4C_4b^4 \\(a - b)^4 &= {}^4C_0a^4 - {}^4C_1a^3b + {}^4C_2a^2b^2 - {}^4C_3ab^3 + {}^4C_4b^4 \\ \therefore (a + b)^4 - (a - b)^4 &= {}^4C_0a^4 + {}^4C_1a^3b + {}^4C_2a^2b^2 + {}^4C_3ab^3 + {}^4C_4b^4 \\ &\quad - [{}^4C_0a^4 - {}^4C_1a^3b + {}^4C_2a^2b^2 - {}^4C_3ab^3 + {}^4C_4b^4] \\ &= 2({}^4C_1a^3b + {}^4C_3ab^3) = 2(4a^3b + 4ab^3) \\ &= 8ab(a^2 + b^2)\end{aligned}$$

By putting $a = \sqrt{3}$ and $b = \sqrt{2}$, we obtain

$$\begin{aligned}(\sqrt{3} + \sqrt{2})^4 - (\sqrt{3} - \sqrt{2})^4 &= 8(\sqrt{3})(\sqrt{2})\{(\sqrt{3})^2 + (\sqrt{2})^2\} \\ &= 8(\sqrt{6})\{3 + 2\} = 40\sqrt{6}\end{aligned}$$

Question-12

Find $(x + 1)^6 + (x - 1)^6$. Hence or otherwise evaluate $(\sqrt{2} + 1)^6 + (\sqrt{2} - 1)^6$.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Using Binomial Theorem, the expressions, $(x + 1)^6$ and $(x - 1)^6$, can be expanded as

$$(x+1)^6 = {}^6C_0x^6 + {}^6C_1x^5 + {}^6C_2x^4 + {}^6C_3x^3 + {}^6C_4x^2 + {}^6C_5x + {}^6C_6$$

$$(x-1)^6 = {}^6C_0x^6 - {}^6C_1x^5 + {}^6C_2x^4 - {}^6C_3x^3 + {}^6C_4x^2 - {}^6C_5x + {}^6C_6$$

$$\begin{aligned}\therefore (x+1)^6 + (x-1)^6 &= 2[{}^6C_0x^6 + {}^6C_2x^4 + {}^6C_4x^2 + {}^6C_6] \\ &= 2[x^6 + 15x^4 + 15x^2 + 1]\end{aligned}$$

By putting $x = \sqrt{2}$, we obtain

$$\begin{aligned}(\sqrt{2}+1)^6 + (\sqrt{2}-1)^6 &= 2\left[(\sqrt{2})^6 + 15(\sqrt{2})^4 + 15(\sqrt{2})^2 + 1\right] \\ &= 2(8 + 15 \times 4 + 15 \times 2 + 1) \\ &= 2(8 + 60 + 30 + 1) \\ &= 2(99) = 198\end{aligned}$$

Question-13

Show that $9^{n+1} - 8n - 9$ is divisible by 64, whenever n is a positive integer.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

In order to show that $9^{n+1} - 8n - 9$ is divisible by 64, it has to be proved that,

$$9^{n+1} - 8n - 9 = 64k, \text{ where } k \text{ is some natural number}$$

By Binomial Theorem,

$$(1+a)^m = {}^mC_0 + {}^mC_1a + {}^mC_2a^2 + \dots + {}^mC_ma^m$$

For $a = 8$ and $m = n + 1$, we obtain

$$\begin{aligned}(1+8)^{n+1} &= {}^{n+1}C_0 + {}^{n+1}C_1(8) + {}^{n+1}C_2(8)^2 + \dots + {}^{n+1}C_{n+1}(8)^{n+1} \\ \Rightarrow 9^{n+1} &= 1 + (n+1)(8) + 8^2 \left[{}^{n+1}C_2 + {}^{n+1}C_3 \times 8 + \dots + {}^{n+1}C_{n+1}(8)^{n-1} \right] \\ \Rightarrow 9^{n+1} &= 9 + 8n + 64 \left[{}^{n+1}C_2 + {}^{n+1}C_3 \times 8 + \dots + {}^{n+1}C_{n+1}(8)^{n-1} \right] \\ \Rightarrow 9^{n+1} - 8n - 9 &= 64k, \text{ where } k = {}^{n+1}C_2 + {}^{n+1}C_3 \times 8 + \dots + {}^{n+1}C_{n+1}(8)^{n-1} \text{ is a natural number}\end{aligned}$$

Thus, $9^{n+1} - 8n - 9$ is divisible by 64, whenever n is a positive integer.

Question-14

Prove that $\sum_{r=0}^n 3^r {}^nC_r = 4^n$.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

By Binomial Theorem,

$$\sum_{r=0}^n {}^nC_r a^{n-r} b^r = (a+b)^n$$

By putting $b = 3$ and $a = 1$ in the above equation, we obtain

$$\begin{aligned}\sum_{r=0}^n {}^nC_r (1)^{n-r} (3)^r &= (1+3)^n \\ \Rightarrow \sum_{r=0}^n 3^r {}^nC_r &= 4^n\end{aligned}$$

Hence, proved.

NCERT 11th Maths Chapter 8, class 11 Maths Chapter 8 solutions

Exercise.8.2

Question-1

Find the coefficient of x^5 in $(x + 3)^8$

Ans.

It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Assuming that x^5 occurs in the $(r + 1)^{\text{th}}$ term of the expansion $(x + 3)^8$, we obtain

$$T_{r+1} = {}^8C_r (x)^{8-r} (3)^r$$

Comparing the indices of x in x^5 and in T_{r+1} , we obtain

$$r = 3$$

$$\text{Thus, the coefficient of } x^5 \text{ is } {}^8C_3 (3)^3 = \frac{8!}{3!5!} \times 3^3 = \frac{8 \cdot 7 \cdot 6 \cdot 5!}{3 \cdot 2 \cdot 5!} \cdot 3^3 = 1512$$

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Question-2

Find the coefficient of a^5b^7 in $(a - 2b)^{12}$

Ans.

It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Assuming that a^5b^7 occurs in the $(r + 1)^{\text{th}}$ term of the expansion $(a - 2b)^{12}$, we obtain

$$T_{r+1} = {}^{12}C_r (a)^{12-r} (-2b)^r = {}^{12}C_r (-2)^r (a)^{12-r} (b)^r$$

Comparing the indices of a and b in a^5b^7 and in T_{r+1} , we obtain

$$r = 7$$

Thus, the coefficient of a^5b^7 is

$${}^{12}C_7 (-2)^7 = -\frac{12!}{7!5!} \cdot 2^7 = -\frac{12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7!}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 7!} \cdot 2^7 = -(792)(128) = -101376$$

Question-3

Write the general term in the expansion of $(x^2 - y)^6$

Ans.

It is known that the general term T_{r+1} {which is the $(r + 1)^{\text{th}}$ term} in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Thus, the general term in the expansion of $(x^2 - y)^6$ is

$$T_{r+1} = {}^6C_r (x^2)^{6-r} (-y)^r = (-1)^r {}^6C_r x^{12-2r} y^r$$

Question-4

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Write the general term in the expansion of $(x^2 - yx)^{12}$, $x \neq 0$

Ans.

It is known that the general term T_{r+1} {which is the $(r + 1)^{\text{th}}$ term} in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Thus, the general term in the expansion of $(x^2 - yx)^{12}$ is

$$T_{r+1} = {}^{12}C_r (x^2)^{12-r} (-yx)^r = (-1)^r {}^{12}C_r x^{24-2r} y^r x^r = (-1)^r {}^{12}C_r x^{24-r} y^r$$

Question-5

Find the 4th term in the expansion of $(x - 2y)^{12}$.

Ans.

It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Thus, the 4th term in the expansion of $(x - 2y)^{12}$ is

$$T_4 = T_{3+1} = {}^{12}C_3 (x)^{12-3} (-2y)^3 = (-1)^3 \cdot \frac{12!}{3!9!} \cdot x^9 \cdot (2)^3 \cdot y^3 = -\frac{12 \cdot 11 \cdot 10}{3 \cdot 2} \cdot (2)^3 x^9 y^3 = -1760x^9 y^3$$

Question-6

Find the 13th term in the expansion of $\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}$, $x \neq 0$.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Thus, 13th term in the expansion of $\left(9x - \frac{1}{3\sqrt{x}}\right)^{18}$ is

$$\begin{aligned} T_{13} = T_{12+1} &= {}^{18}C_{12} (9x)^{18-12} \left(-\frac{1}{3\sqrt{x}}\right)^{12} \\ &= (-1)^{12} \frac{18!}{12!6!} (9)^6 (x)^6 \left(\frac{1}{3}\right)^{12} \left(\frac{1}{\sqrt{x}}\right)^{12} \\ &= \frac{18 \cdot 17 \cdot 16 \cdot 15 \cdot 14 \cdot 13 \cdot 12!}{12! \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2} \cdot x^6 \cdot \left(\frac{1}{x^6}\right) \cdot 3^{12} \left(\frac{1}{3^{12}}\right) \quad \left[9^6 = (3^2)^6 = 3^{12}\right] \\ &= 18564 \end{aligned}$$

Question-7

Find the middle terms in the expansions of $\left(3 - \frac{x^3}{6}\right)^7$

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

It is known that in the expansion of $(a + b)^n$, if n is odd, then there are two middle terms, namely, $\left(\frac{n+1}{2}\right)^{\text{th}}$ term and $\left(\frac{n+1}{2} + 1\right)^{\text{th}}$ term.

Therefore, the middle terms in the expansion of $\left(3 - \frac{x^3}{6}\right)^7$ are $\left(\frac{7+1}{2}\right)^{\text{th}} = 4^{\text{th}}$ term and $\left(\frac{7+1}{2} + 1\right)^{\text{th}} = 5^{\text{th}}$ term

$$\begin{aligned}T_4 = T_{4+1} &= {}^7C_3 (3)^{7-3} \left(-\frac{x^3}{6}\right)^3 = (-1)^3 \frac{7!}{3!4!} \cdot 3^4 \cdot \frac{x^9}{6^3} \\&= -\frac{7 \cdot 6 \cdot 5 \cdot 4!}{3 \cdot 2 \cdot 4!} \cdot 3^4 \cdot \frac{1}{2^3 \cdot 3^3} \cdot x^9 = -\frac{105}{8} x^9 \\T_5 = T_{4+1} &= {}^7C_4 (3)^{7-4} \left(-\frac{x^3}{6}\right)^4 = (-1)^4 \frac{7!}{4!3!} (3)^3 \cdot \frac{x^{12}}{6^4} \\&= \frac{7 \cdot 6 \cdot 5 \cdot 4!}{4! \cdot 3 \cdot 2} \cdot \frac{3^3}{2^4 \cdot 3^4} \cdot x^{12} = \frac{35}{48} x^{12}\end{aligned}$$

Thus, the middle terms in the expansion of $\left(3 - \frac{x^3}{6}\right)^7$ are $-\frac{105}{8} x^9$ and $\frac{35}{48} x^{12}$.

Question-8

Thus, the middle terms in the expansion of $\left(3 - \frac{x^3}{6}\right)^7$ are $-\frac{105}{8} x^9$ and $\frac{35}{48} x^{12}$.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

It is known that in the expansion $(a + b)^n$, if n is even, then the middle term is

$\left(\frac{n}{2} + 1\right)^{\text{th}}$ term.

Therefore, the middle term in the expansion of $\left(\frac{x}{3} + 9y\right)^{10}$ is $\left(\frac{10}{2} + 1\right)^{\text{th}} = 6^{\text{th}}$ term

$$\begin{aligned}T_6 = T_{5+1} &= {}^{10}C_5 \left(\frac{x}{3}\right)^{10-5} (9y)^5 = \frac{10!}{5!5!} \cdot \frac{x^5}{3^5} \cdot 9^5 \cdot y^5 \\&= \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5!}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 5!} \cdot \frac{1}{3^5} \cdot 3^{10} \cdot x^5 y^5 \quad \left[9^5 = (3^2)^5 = 3^{10}\right] \\&= 252 \times 3^5 \cdot x^5 \cdot y^5 = 61236 x^5 y^5\end{aligned}$$

Thus, the middle term in the expansion of $\left(\frac{x}{3} + 9y\right)^{10}$ is $61236 x^5 y^5$.

Question-9

In the expansion of $(1 + a)^{m+n}$, prove that coefficients of a^m and a^n are equal.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>



It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Assuming that a^m occurs in the $(r + 1)^{\text{th}}$ term of the expansion $(1 + a)^{m+n}$, we obtain

$$T_{r+1} = {}^{m+n}C_r (1)^{m+n-r} (a)^r = {}^{m+n}C_r a^r$$

Comparing the indices of a in a^m and in T_{r+1} , we obtain

$$r = m$$

Therefore, the coefficient of a^m is

$${}^{m+n}C_m = \frac{(m+n)!}{m!(m+n-m)!} = \frac{(m+n)!}{m!n!} \quad \dots(1)$$

Assuming that a^n occurs in the $(k + 1)^{\text{th}}$ term of the expansion $(1 + a)^{m+n}$, we obtain

$$T_{k+1} = {}^{m+n}C_k (1)^{m+n-k} (a)^k = {}^{m+n}C_k (a)^k$$

Comparing the indices of a in a^n and in T_{k+1} , we obtain

$$k = n$$

Therefore, the coefficient of a^n is

$${}^{m+n}C_n = \frac{(m+n)!}{n!(m+n-n)!} = \frac{(m+n)!}{n!m!} \quad \dots(2)$$

Thus, from (1) and (2), it can be observed that the coefficients of a^m and a^n in the expansion of $(1 + a)^{m+n}$ are equal.

Question-10

The coefficients of the $(r - 1)^{\text{th}}$, r^{th} and $(r + 1)^{\text{th}}$ terms in the expansion of

$(x + 1)^n$ are in the ratio 1:3:5. Find n and r .

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

It is known that $(k + 1)^{\text{th}}$ term, (T_{k+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{k+1} = {}^nC_k a^{n-k} b^k$.

Therefore, $(r - 1)^{\text{th}}$ term in the expansion of $(x + 1)^n$ is

$$T_{r-1} = {}^nC_{r-2} (x)^{n-(r-2)} (1)^{(r-2)} = {}^nC_{r-2} x^{n-r+2}$$

r^{th} term in the expansion of $(x + 1)^n$ is $T_r = {}^nC_{r-1} (x)^{n-(r-1)} (1)^{(r-1)} = {}^nC_{r-1} x^{n-r+1}$

$(r + 1)^{\text{th}}$ term in the expansion of $(x + 1)^n$ is $T_{r+1} = {}^nC_r (x)^{n-r} (1)^r = {}^nC_r x^{n-r}$

Therefore, the coefficients of the $(r - 1)^{\text{th}}$, r^{th} , and $(r + 1)^{\text{th}}$ terms in the expansion of $(x + 1)^n$ are ${}^nC_{r-2}$, ${}^nC_{r-1}$, and nC_r respectively. Since these coefficients are in the ratio 1:3:5, we obtain

$$\begin{aligned} \frac{{}^nC_{r-2}}{{}^nC_{r-1}} &= \frac{1}{3} \text{ and } \frac{{}^nC_{r-1}}{{}^nC_r} = \frac{3}{5} \\ \frac{{}^nC_{r-2}}{{}^nC_{r-1}} &= \frac{n!}{(r-2)!(n-r+2)!} \times \frac{(r-1)!(n-r+1)!}{n!} = \frac{(r-1)(r-2)!(n-r+1)!}{(r-2)!(n-r+2)(n-r+1)!} \\ &= \frac{r-1}{n-r+2} \end{aligned}$$

$$\therefore \frac{r-1}{n-r+2} = \frac{1}{3}$$

$$\Rightarrow 3r-3 = n-r+2$$

$$\Rightarrow n-4r+5=0 \quad \dots(1)$$

$$\therefore \frac{r-1}{n-r+2} = \frac{1}{3}$$

$$\Rightarrow 3r-3 = n-r+2$$

$$\Rightarrow n-4r+5=0 \quad \dots(1)$$

$$\begin{aligned} \frac{{}^nC_{r-1}}{{}^nC_r} &= \frac{n!}{(r-1)!(n-r+1)!} \times \frac{r!(n-r)!}{n!} = \frac{r(r-1)!(n-r)!}{(r-1)!(n-r+1)(n-r)!} \\ &= \frac{r}{n-r+1} \end{aligned}$$

$$\therefore \frac{r}{n-r+1} = \frac{3}{5}$$

$$\Rightarrow 5r = 3n - 3r + 3$$

$$\Rightarrow 3n - 8r + 3 = 0 \quad \dots(2)$$

Multiplying (1) by 3 and subtracting it from (2), we obtain

$$4r - 12 = 0$$

$$r = 3$$

Putting the value of r in (1), we obtain

$$n - 12 + 5 = 0$$

$$n = 7$$

Thus, $n = 7$ and $r = 3$

Question-11

Prove that the coefficient of x^n in the expansion of $(1+x)^{2n}$ is twice the coefficient of x^n in the expansion of $(1+x)^{2n-1}$

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Assuming that x^n occurs in the $(r + 1)^{\text{th}}$ term of the expansion of $(1 + x)^{2n}$, we obtain

$$T_{r+1} = {}^{2n}C_r (1)^{2n-r} (x)^r = {}^{2n}C_r (x)^r$$

Comparing the indices of x in x^n and in T_{r+1} , we obtain

$$r = n$$

Therefore, the coefficient of x^n in the expansion of $(1 + x)^{2n}$ is

$${}^{2n}C_n = \frac{(2n)!}{n!(2n-n)!} = \frac{(2n)!}{n!n!} = \frac{(2n)!}{(n!)^2} \quad \dots(1)$$

Assuming that x^n occurs in the $(k + 1)^{\text{th}}$ term of the expansion $(1 + x)^{2n-1}$, we obtain

$$T_{k+1} = {}^{2n-1}C_k (1)^{2n-1-k} (x)^k = {}^{2n-1}C_k (x)^k$$

Comparing the indices of x in x^n and T_{k+1} , we obtain

$$k = n$$

Therefore, the coefficient of x^n in the expansion of $(1 + x)^{2n-1}$ is

$$\begin{aligned} {}^{2n-1}C_n &= \frac{(2n-1)!}{n!(2n-1-n)!} = \frac{(2n-1)!}{n!(n-1)!} \\ &= \frac{2n \cdot (2n-1)!}{2n \cdot n!(n-1)!} = \frac{(2n)!}{2n!n!} = \frac{1}{2} \left[\frac{(2n)!}{(n!)^2} \right] \quad \dots(2) \end{aligned}$$

From (1) and (2), it is observed that

$$\frac{1}{2} \left({}^{2n}C_n \right) = {}^{2n-1}C_n$$
$$\Rightarrow {}^{2n}C_n = 2 \left({}^{2n-1}C_n \right)$$

Therefore, the coefficient of x^n in the expansion of $(1 + x)^{2n}$ is twice the coefficient of x^n in the expansion of $(1 + x)^{2n-1}$.

Hence, proved.

Question-12

Find a positive value of m for which the coefficient of x^2 in the expansion

$(1 + x)^m$ is 6.

Ans.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

It is known that $(r + 1)^{\text{th}}$ term, (T_{r+1}) , in the binomial expansion of $(a + b)^n$ is given by $T_{r+1} = {}^nC_r a^{n-r} b^r$.

Assuming that x^2 occurs in the $(r + 1)^{\text{th}}$ term of the expansion $(1 + x)^m$, we obtain

$$T_{r+1} = {}^mC_r (1)^{m-r} (x)^r = {}^mC_r (x)^r$$

Comparing the indices of x in x^2 and in T_{r+1} , we obtain

$$r = 2$$

Therefore, the coefficient of x^2 is mC_2 .

It is given that the coefficient of x^2 in the expansion $(1 + x)^m$ is 6.

$$\therefore {}^mC_2 = 6$$

$$\Rightarrow \frac{m!}{2!(m-2)!} = 6$$

$$\Rightarrow \frac{m(m-1)(m-2)!}{2 \times (m-2)!} = 6$$

$$\Rightarrow m(m-1) = 12$$

$$\Rightarrow m^2 - m - 12 = 0$$

$$\Rightarrow m^2 - 4m + 3m - 12 = 0$$

$$\Rightarrow m(m-4) + 3(m-4) = 0$$

$$\Rightarrow (m-4)(m+3) = 0$$

$$\Rightarrow (m-4) = 0 \text{ or } (m+3) = 0$$

$$\Rightarrow m = 4 \text{ or } m = -3$$

Thus, the positive value of m , for which the coefficient of x^2 in the expansion

$(1 + x)^m$ is 6, is 4.

NCERT 11th Maths Chapter 8, class 11 Maths Chapter 8 solutions



<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

Chapterwise NCERT Solutions for Class 11 Maths :

- Chapter 1-Sets
- Chapter 2-Relations and Functions
- Chapter 3-Trigonometric Functions
- Chapter 4-Principle of Mathematical Induction
- Chapter 5-Complex Numbers and Quadratic Equations
- Chapter 6-Linear Inequalities
- Chapter 7-Permutation and Combinations
- Chapter 8-Binomial Theorem
- Chapter 9-Sequences and Series
- Chapter 10-Straight Lines
- Chapter 11-Conic Sections
- Chapter 12-Introduction to three Dimensional Geometry
- Chapter 13-Limits and Derivatives
- Chapter 14-Mathematical Reasoning
- Chapter 15-Statistics
- Chapter 16-Probability

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>

About NCERT

The National Council of Educational Research and Training is an autonomous organization of the Government of India which was established in 1961 as a literary, scientific, and charitable Society under the Societies Registration Act. The major objectives of NCERT and its constituent units are to: undertake, promote and coordinate research in areas related to school education; prepare and publish model textbooks, supplementary material, newsletters, journals and develop educational kits, multimedia digital materials, etc.

Organise pre-service and in-service training of teachers; develop and disseminate innovative educational techniques and practices; collaborate and network with state educational departments, universities, NGOs and other educational institutions; act as a clearing house for ideas and information in matters related to school education; and act as a nodal agency for achieving the goals of Universalisation of Elementary Education. In addition to research, development, training, extension, publication and dissemination activities, NCERT is an implementation agency for bilateral cultural exchange programmes with other countries in the field of school education. Its headquarters are located at Sri Aurobindo Marg in New Delhi. [Visit the Official NCERT website](#) to learn more.

<https://www.indcareer.com/schools/ncert-solutions-for-11th-class-maths-chapter-8-binomial-theorem/>