



एन सी ई आर टी
NCERT

NCERT

National Council Of Educational Research
And Training

NCERT Solutions for 12th Class Chemistry: Chapter 2-Solutions



IndCareer
Schools



indCareer



indCareer



indCareer

NCERT Solutions for 12th Class Chemistry: Chapter 2-Solutions

Class 12: Chemistry Chapter 2 solutions. Complete Class 12 Chemistry Chapter 2 Notes.

NCERT Solutions for 12th Class Chemistry: Chapter 2-Solutions

NCERT 12th Chemistry Chapter 2, class 12 Chemistry chapter 2 solutions

NCERT 12th Chemistry Chapter 2

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

2.1. Calculate the mass percentage of benzene (C₆H₆) and carbon tetrachloride (CCl₄) if 22 g of benzene is dissolved in 122 g of carbon tetrachloride.

Ans: Mass of solution = Mass of C₆H₆ + Mass of CCl₄
= 22 g + 122 g = 144 g

Mass % of benzene = $\frac{22}{144} \times 100 = 15.28 \%$

Mass % of CCl₄ = $\frac{122}{144} \times 100 = 84.72 \%$

2.2. Calculate the mole fraction of benzene in solution containing 30% by mass in carbon tetrachloride.

Ans: 30% by mass of C₆H₆ in CCl₄ => 30 g C₆H₆ in 100 g solution

∴ no. of moles of C₆H₆, ($n_{C_6H_6}$) = $\frac{30}{78} = 0.385$

(molar mass of C₆H₆ = 78g)

no. of moles of
CCl₄ (n_{CCl_4}) = $\frac{70}{154} = 0.455$

$$x_{C_6H_6} = \frac{n_{C_6H_6}}{n_{C_6H_6} + n_{CCl_4}}$$
$$= \frac{0.385}{0.385 + 0.455} = \frac{0.385}{0.84} = 0.458$$
$$x_{CCl_4} = 1 - 0.458 = 0.542$$

2.3. Calculate the molarity of each of the following solutions

(a) 30 g of Co(NO₃)₂·6H₂O in 4.3 L of solution

(b) 30 mL of 0.5 M H₂SO₄ diluted to 500 mL.

Ans:

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

(a)
$$\text{Molarity of solution} = \frac{\text{Mass of solute} / \text{Molar mass of solute}}{\text{Volume of solution in litres}}$$

Mass of solute, $\text{Co}(\text{NO}_3)_2 \cdot 6 \text{H}_2\text{O} = 30 \text{ g}$.
Molar mass of solute, $\text{Co}(\text{NO}_3)_2 \cdot 6 \text{H}_2\text{O} = 59 + 2 \times 14 + 6 \times 16 + 6 \times 18 = 291 \text{ g mol}^{-1}$.
Volume of solution = 4.3 L

$$\text{Molarity (M)} = \frac{(30\text{g}) / (291\text{g mol}^{-1})}{(4.3\text{L})} = 0.024 \text{ mol L}^{-1} = \mathbf{0.024 \text{ M}}$$

(b) Volume of undiluted H_2SO_4 solution (V_1) = 30 mL
Molarity of undiluted H_2SO_4 solution (M_1) = 0.5 M
Volume of diluted H_2SO_4 solution (V_2) = 500 mL

Molarity of diluted H_2SO_4 (M_2) can be calculated as :

$$M_1 V_1 = M_2 V_2$$

or
$$M_2 = \frac{M_1 V_1}{V_2} = \frac{(30\text{mL}) \times (0.5\text{M})}{(500\text{mL})} = \mathbf{0.03 \text{ M}}$$

2.4. Calculate the mass of urea (NH_2CONH_2) required in making 2.5 kg of 0.25 molal aqueous solution.

Ans: 0.25 Molal aqueous solution to urea means that

moles of urea = 0.25 mole

mass of solvent (NH_2CONH_2) = 60 g mol⁻¹

∴ 0.25 mole of urea = 0.25 x 60 = 15g

Mass of solution = 1000 + 15 = 1015g = 1.015 kg

1.015 kg of urea solution contains 15g of urea

∴ 2.5 kg of solution contains urea = 15 / 1.015 x 2.5 = 37g

2.5. Calculate

(a) molality

(b) molarity and

(c) mole fraction of KI if the density of 20% (mass/mass) aqueous KI solution is 1.202 g mL^{-1} .

Ans:

Step I. Calculation of molality of solution

Weight of KI in 100 g of the solution = 20 g

Weight of water in the solution = $100 - 20 = 80 \text{ g} = 0.08 \text{ kg}$

Molar mass of KI = $39 + 127 = 166 \text{ g mol}^{-1}$.

$$\begin{aligned}\text{Molality of solution } (m) &= \frac{\text{No of gram moles of KI}}{\text{Mass of water in kg}} = \frac{(20\text{g})/(166\text{g mol}^{-1})}{(0.08\text{kg})} \\ &= 1.506 \text{ mol kg}^{-1} = \mathbf{1.506 \text{ m.}}\end{aligned}$$

Step II. Calculation of molarity of solution

Weight of solution = 100 g ; Density of solution = 1.202 g mL^{-1} .

$$\text{Volume of solution} = \frac{\text{Weight of solution}}{\text{Density}} = \frac{(100\text{g})}{(1.202\text{g mL}^{-1})} = 83.19 \text{ mL}$$

$$\begin{aligned}\text{Molarity of solution } (M) &= \frac{\text{No. of gram moles of KI}}{\text{Volume of solution in litres}} = \frac{(20\text{g})/(166\text{g mol}^{-1})}{(0.083\text{L})} \\ &= 1.45 \text{ mol L}^{-1} = \mathbf{1.45 \text{ M}}\end{aligned}$$

Step III. Calculation of mole fraction of KI

$$\begin{aligned}n_{\text{KI}} &= \frac{\text{Mass of KI}}{\text{Molar mass of KI}} = \frac{(20\text{g})}{(166\text{g mol}^{-1})} = 0.12 \text{ mol} \\ n_{\text{H}_2\text{O}} &= \frac{\text{Mass of water}}{\text{Molar mass of water}} = \frac{(80\text{g})}{(18\text{g mol}^{-1})} = 4.44 \text{ mol.} \\ x_{\text{KI}} &= \frac{n_{\text{KI}}}{n_{\text{KI}} + n_{\text{H}_2\text{O}}} = \frac{(0.12\text{mol})}{(0.12 + 4.44)\text{mol}} = \frac{0.12}{4.56} = \mathbf{0.0263.}\end{aligned}$$

2.6. H₂S, a toxic gas with rotten egg like smell, is used for the qualitative analysis. If the solubility of H₂S in water at STP is 0.195 m, calculate Henry's law constant.

Ans: Solubility of H₂S gas = 0.195 m

= 0.195 mole in 1 kg of solvent

1 kg of solvent = 1000g

$$= \frac{1000}{18} = 55.55 \text{ moles}$$

$$\therefore x_{\text{H}_2\text{S}} = \frac{0.195}{0.195 + 55.55}$$

$$= \frac{0.195}{55.745} = 0.0035$$

– Pressure at STP = 0.987 bar

Applying Henry's law,

$$P_{\text{H}_2\text{S}} = K_{\text{H}} \times x_{\text{H}_2\text{S}}$$

$$\Rightarrow K_{\text{H}} = \frac{P_{\text{H}_2\text{S}}}{x_{\text{H}_2\text{S}}} = \frac{0.987}{0.0035} = 282 \text{ bar}$$

2.7. Henry's law constant for CO₂ in water is 1.67 x 10⁸ Pa at 298 K. Calculate the quantity of CO₂ in 500 mL of soda water when packed under 2.5 atm CO₂ pressure at 298 K.

Ans.:

$$K_H = 1.67 \times 10^8 \text{ Pa}$$

$$P_{\text{CO}_2} = 2.5 \text{ atm} = 2.5 \times 101325 \text{ Pa}$$

$$\therefore x_{\text{CO}_2} = \frac{P_{\text{CO}_2}}{K_H} = \frac{2.5 \times 101325}{1.67 \times 10^8} = 1.517 \times 10^{-3}$$

For 500 mL of soda water, water present \approx 500 mL

$$= 500 \text{ g} = \frac{500}{18} = 27.78 \text{ moles}$$

$$\therefore n_{\text{H}_2\text{O}} = 27.78 \text{ moles}$$

$$\therefore \frac{n_{\text{CO}_2}}{27.78} = 1.517 \times 10^{-3}$$

$$\begin{aligned} n_{\text{CO}_2} &= 42.14 \times 10^{-3} \text{ mole} \\ &= 42.14 \text{ mmol} \\ &= 42.14 \times 10^{-3} \times 44 \text{ g} \\ &= 1.854 \text{ g} \end{aligned}$$

NCERT 12th Chemistry Chapter 2

2.8 The vapour pressures of pure liquids A and B are 450 mm and 700 mm of Hg respectively at 350 K. Calculate the composition of the liquid mixture if total vapour pressure is 600 mm of Hg. Also find the composition in the vapour phase.

Ans:

Vapour pressure of pure liquid A ($P^\circ \text{A}$) = 450 mm

Vapour pressure of pure liquid B ($P^\circ \text{B}$) = 700 mm

Total vapour pressure of the solution (P) = 600 mm

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

$$\begin{aligned}\text{According to Raoult's Law, } P &= P_A^\circ x_A + P_B^\circ x_B = P_A^\circ x_A + P_B^\circ (1 - x_A) \\ (600 \text{ mm}) &= 450 \text{ mm} \times x_A + 700 \text{ mm} (1 - x_A) \\ &= 700 \text{ mm} + x_A (450 - 700) \text{ mm} \\ &= 700 - x_A (250 \text{ mm})\end{aligned}$$

or

$$x_A = \frac{(600 - 700) \text{ mm}}{-(250 \text{ mm})} = 0.40$$

$$\text{Mole fraction of A } (x_A) = 0.40$$

$$\text{Mole fraction of B } (x_B) = 1 - 0.40 = 0.60$$

$$P_A = P_A^\circ x_A = (450 \text{ mm}) \times 0.40 = 180 \text{ mm}$$

$$P_B = P_B^\circ x_B = (700 \text{ mm}) \times 0.60 = 420 \text{ mm}$$

$$\text{Mole fraction of A in the vapour phase} = \frac{P_A}{P_A + P_B} = \frac{(180 \text{ mm})}{(180 + 420) \text{ mm}} = 0.30$$

$$\text{Mole fraction of B in the vapour phase} = \frac{P_B}{P_A + P_B} = \frac{(420 \text{ mm})}{(180 + 420) \text{ mm}} = 0.70$$

2.9. Vapour pressure of pure water at 298 K is 23.8 mm Hg. 50 g of urea (NH_2CONH_2) is dissolved in 850 g of water. Calculate the vapour pressure of water for this solution and its relative lowering.

Ans:

$$P^{\circ} = 23.8 \text{ mm}$$

$$w_2 = 50 \text{ g}, M_2 (\text{urea}) = 60 \text{ g mol}^{-1}$$

$$w_1 = 850 \text{ g}, M_1 (\text{water}) = 18 \text{ g mol}^{-1}$$

To find: P_s and $(P^{\circ} - P_s)/P^{\circ}$

Solution: Applying Raoult's law,

$$\frac{P^{\circ} - P_s}{P^{\circ}} = \frac{n_2}{n_1 + n_2} = \frac{w_2 / M_2}{w_1 / M_1 + w_2 / M_2}$$

$$\begin{aligned} \therefore \frac{P^{\circ} - P_s}{P^{\circ}} &= \frac{50 / 60}{850 / 18 + 50 / 60} \\ &= \frac{0.83}{47.22 + 0.83} = 0.017 \end{aligned}$$

Putting $P^{\circ} = 23.8 \text{ mm}$, we have

$$\frac{23.8 - P_s}{P_s} = 0.017$$

$$\Rightarrow 23.8 - P_s = 0.017 P_s$$

$$\text{or, } 1.017 P_s = 23.8$$

$$\text{or, } P_s = 23.4 \text{ mm}$$

2.10. Boiling point of water at 750 mm Hg is 99.63°C . How much sucrose is to be added to 500 g of water such that it boils at 100°C .

Ans:

Given $\Delta T_b = 100 - 96.63 = 3.37^\circ$

Mass of water, $w_1 = 500$ g

Molar mass of water, $M_1 = 18$ g mol⁻¹

Molar mass of sucrose, $M_2 = 342$ g mol⁻¹

To find: Mass of sucrose, $w_2 = ?$

Solution: We know, $\Delta T_b = K_b \times m$

$$= K_b \times \frac{w_2}{M_2} \times \frac{1000}{w_1}$$

$$\Rightarrow w_2 = \frac{M_2 \times w_1 \times \Delta T_b}{1000 \times K_b} = \frac{342 \times 500 \times 3.37}{1000 \times 0.52}$$

$$w_2 = 1108.2$$

\therefore Mass of solute, $w_2 = 1.11$ kg

2.11 Calculate the mass of ascorbic acid (vitamin C, C₆H₈O₆) to be dissolved in 75 g of acetic acid to lower its melting point by 1.5°C. (K_f for CH₃COOH) = 3.9 K kg mol⁻¹)

Ans:

$$W_B = \frac{M_B \times \Delta T_f \times W_A}{K_f}$$

Mass of acetic acid (W_A) = 75 g = 0.075 kg.

Depression in freezing point (ΔT_f) = 1.5°C = 1.5 K

Molar mass of ascorbic acid (M_B) = 6 × 12 + 8 × 1 + 6 × 16 = 176 g mol⁻¹

Molal depression constant (K_f) = 3.9 K kg mol⁻¹

$$W_B = \frac{(176 \text{ g mol}^{-1}) \times (1.5 \text{ K}) \times (0.075 \text{ kg})}{(3.9 \text{ K kg mol}^{-1})} = 5.08 \text{ g}$$

2.12. Calculate the osmotic pressure in pascals exerted by a solution prepared by dissolving 1.0 g of polymer of molar mass 185,000 in 450 mL of water at 37°C.

Ans:

Given: $V = 450 \text{ mL} = 0.45 \text{ L}$
 $T = 37^\circ\text{C} = 310 \text{ K}$
 $R = 8.314 \text{ kPa L K}^{-1} \text{ mol}^{-1}$

To find: $\pi = ?$

Solution: Applying the formula,

$$\pi = CRT = \frac{n}{V}RT$$

$$n = \frac{1.0\text{g}}{185,000\text{g mol}^{-1}}$$

$$\begin{aligned}\therefore P &= \frac{1}{185,000} \times \frac{1}{0.45} \times 8.314 \\ &\quad \times 10^3 \text{ Pa L K}^{-1} \text{ mol}^{-1} \times 310\text{K} \\ &= 30.96 \text{ Pa}\end{aligned}$$

EXERCISES (NCERT 12th Chemistry Chapter 2)

2.1. Define the terra solution. How many types of solutions are formed? Write briefly about each type with an example.

Sol: A solution is a homogeneous mixture of two or more chemically non-reacting substances. Types of solutions: There are nine types of solutions.

Types of Solution Examples

Gaseous solutions

- (a) Gas in gas Air, mixture of O_2 and N_2 , etc.
- (b) Liquid in gas Water vapour
- (c) Solid in gas Camphor vapours in N_2 gas, smoke etc.

Liquid solutions

- (a) Gas in liquid CO_2 dissolved in water (aerated water), and O_2 dissolved in water, etc.

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

(b) Liquid in liquid Ethanol dissolved in water, etc.

(c) Solid in liquid Sugar dissolved in water, saline water, etc.

Solid solutions

(a) Gas in solid Solution of hydrogen in palladium

(b) Liquid in solid Amalgams, e.g., Na-Hg

(c) Solid in solid Gold ornaments (Cu/Ag with Au)

2.2. Concentrated nitric acid used in the laboratory work is 68% nitric acid by mass in aqueous solution. What should be the molarity of such a sample of acid if the density of the solution is 1.504 g mL^{-1} ?

Sol: Mass of HNO_3 in solution = 68 g

Molar mass of $\text{HNO}_3 = 63 \text{ g mol}^{-1}$

Mass of solution = 100 g

Density of solution = 1.504 g mL^{-1}

$$\begin{aligned}\text{Volume of solution} &= \frac{\text{Mass of solution}}{\text{Density of solution}} \\ &= \frac{(100 \text{ g})}{(1.504 \text{ g mL}^{-1})} = 66.5 \text{ mL} = 0.0665 \text{ L} \\ \text{Molarity of solution (M)} &= \frac{\text{Mass of HNO}_3 / \text{Molar mass of HNO}_3}{\text{Volume of solution in Litres}} \\ &= \frac{(68 \text{ g} / 63 \text{ g mol}^{-1})}{(0.0665 \text{ L})} = 16.23 \text{ mol L}^{-1} = \mathbf{16.23 \text{ M.}}\end{aligned}$$

2.3. A solution of glucose in water is labelled as 10% w/w, what would be the molality and mole fraction of each component in

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

the solution? If the density of solution is 1.2 g mL^{-1} , then what shall be the molarity of the solution?

Sol: 10 percent w/w solution of glucose in water means 10g glucose and 90g of water.

Molar mass of glucose = 180 g mol^{-1} and molar mass of water = 18 g mol^{-1}

$$\begin{aligned} \therefore 10 \text{ g of glucose} &= \frac{10}{180} = 0.0555 \text{ moles} \\ \text{and } 90 \text{ g of H}_2\text{O} &= \frac{90}{18} = 5 \text{ moles} \\ \therefore \text{Molality of solution} &= \frac{\text{Moles of solute} \times 1000}{\text{Mass of solution in grams}} \\ &= \frac{0.0555}{90} \times 1000 = 0.617 \text{ m} \end{aligned}$$

Mole fraction of glucose

$$\begin{aligned} = X_g &= \frac{\text{No. of moles of glucose}}{\text{No. of moles of glucose} + \text{No. of moles of water}} \\ &= \frac{0.0555}{5 + 0.0555} = 0.01 \end{aligned}$$

Mole fraction of water

$$\begin{aligned} = X_w &= \frac{\text{No. of moles of water}}{\text{No. of moles of glucose} + \text{No. of moles of water}} \\ &= \frac{5}{5 + 0.0555} = 0.99 \end{aligned}$$

Volume of 100g of solution

$$= \frac{\text{Mass of solution}}{\text{Density}} = \frac{100}{1.2} = 83.33 \text{ mL}$$

$$\begin{aligned} \therefore \text{Molarity of solution} &= \frac{0.0555}{83.33} \times 1000 \\ &= 0.67 \text{ M.} \end{aligned}$$

2.4. How many mL of 0.1 M HCl are required to react completely with 1 g mixture of Na_2CO_3 and NaHCO_3 containing equimolar amounts of both?

Sol: Calculation of no. of moles of components in the mixture.

Let x g of Na_2CO_3 is present in the mixture.

$\therefore (1-x)$ g of NaHCO_3 is present in the mixture.

Molar mass of Na_2CO_3

$$= 2 \times 23 + 12 + 3 \times 16 = 106 \text{ g mol}^{-1}$$

and molar mass of NaHCO_3

$$= 23 \times 1 + 1 + 12 + 3 \times 16 = 84 \text{ g mol}^{-1}$$

$$\text{No. of moles of } \text{Na}_2\text{CO}_3 \text{ in } x \text{ g} = \frac{x}{106}$$

$$\text{No. of moles of } \text{NaHCO}_3 \text{ in } (1-x) \text{ g} = (1-x)/84$$

As given that the mixture contains equimolar amounts of Na_2CO_3 and NaHCO_3 , therefore

$$\frac{x}{106} = \frac{1-x}{84}$$

$$106 - 106x = 84x$$

$$106 = 190x$$

$$\therefore x = \frac{106}{190} = 0.558 \text{g}$$



\therefore No. of moles of Na_2CO_3 present

$$= \frac{0.558}{106} = 0.00526$$

and no. of moles of NaHCO_3 present

$$= \frac{1 - 0.558}{84} = 0.00526$$

Calculation of no. of moles of HCl required



As can be seen, each mole of Na_2CO_3 needs 2 moles of HCl,

\therefore 0.00526 mole of Na_2CO_3 needs

$$= 0.00526 \times 2 = 0.01052 \text{ mole}$$

Each mole of NaHCO_3 needs 1 mole of HCl.

\therefore 0.00526 mole of NaHCO_3 needs

$$= 1 \times 0.00526 = 0.00526 \text{ mole}$$

Total amount of HCl needed will be

$$= 0.01052 + 0.00526 = 0.01578 \text{ mole.}$$

0.1 mole of 0.1 M HCl are present in 1000 mL of HCl

\therefore 0.01578 mole of 0.1 M HCl will be present in

$$= \frac{1000}{0.1} \times 0.01578 = 157.8 \text{ mL.}$$

NCERT 12th Chemistry Chapter 2

2.5. Calculate the percentage composition in terms of mass of a solution obtained by mixing 300 g of a 25% and 400 g of a 40% solution by mass.

Sol:

$$\text{Mass of one component in solution} = \frac{(300 \text{ g}) \times 25}{100} = 75 \text{ g}$$

$$\text{Mass of other component in solution} = \frac{(400 \text{ g}) \times 40}{100} = 160 \text{ g}$$

$$\text{Total mass of solute} = (75 + 160) \text{ g} = 235 \text{ g}$$

$$\text{Total mass of solution} = (300 + 400) \text{ g} = 700 \text{ g}$$

$$\% \text{ of solute in the final solution} = \frac{(235 \text{ g})}{(700 \text{ g})} \times 100 = 33.57$$

$$\% \text{ of solvent in the final solution} = 100 - 33.57 = 66.43$$

2.6. An antifreeze solution is prepared from 222.6 g of ethylene glycol, (C₂H₆O₂) and 200 g of water. Calculate the molality of the solution. If the density of the solution is 1.072 g mL⁻¹, then what shall be the molarity of the solution?

Sol:

$$\text{Mass of solvent} = 200 \text{ g}$$

$$\therefore \text{Molality} = \frac{3.59}{200} \times 1000 = 17.95 \text{ mol kg}^{-1}$$

$$\text{Total mass of solution} = 422.6 \text{ g}$$

$$\text{Volume of solution} = \frac{422.6}{1.072} = 394.21 \text{ mL}$$

$$\text{Mass of solute} = 222.6 \text{ g}$$

$$\text{Molar mass of solute, C}_2\text{H}_4(\text{OH})_2 \\ = 12 \times 2 + 4 + 2(12 + 1) = 62 \text{ g mol}^{-1}$$

$$\therefore \text{Moles of solute} = \frac{222.6}{62} = 3.59$$

$$\therefore \text{Molarity} = \frac{3.59}{394.2} \times 1000 = 9.1 \text{ mol L}^{-1}$$

2.7. A sample of drinking water was found to be severely contaminated with chloroform (CHCl₃), supposed to be a carcinogen. The level of contamination was 15 ppm (by mass).

(i) express this in percent by mass.

(ii) determine the molality of chloroform in the water sample.

Sol: 15 ppm means 15 parts in million (10⁶) by mass in the solution.

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

$$\therefore \text{Percentage by mass} = \frac{15}{10^6} \times 100 = 15 \times 10^{-4} \%$$

As only 15g of chloroform is present in 10^6 g of the solution, mass of the solvent = 10^6 g

$$\begin{aligned} \text{Molar mass of CHCl}_3 &= 12 + 1 + 3 \times 35.5 \\ &= 119.5 \text{ g mol}^{-1} \end{aligned}$$

$$\text{Moles of CHCl}_3 = \frac{15}{119.5}$$

$$\therefore \text{Molality} = \frac{15/119.5 \times 1000}{10^6} = 1.25 \times 10^{-4} \text{ m}$$

2.8. What role does the molecular interaction play in solution of alcohol in water?

Sol: In case of alcohol as well as water, the molecules are interlinked by intermolecular hydrogen bonding. However, the hydrogen bonding is also present in the molecules of alcohol and water in the solution but it is comparatively less than both alcohol and water. As a result, the magnitude of attractive forces tends to decrease and the solution shows positive deviation from Raoult's Law. This will lead to increase in vapour pressure of the solution and also decrease in its boiling point.

2.9. Why do gases always tend to be less soluble in liquids as the temperature is raised?

Sol: When gases are dissolved in water, it is accompanied by a release of heat energy, i.e., process is exothermic. When the temperature is increased, according to Lechatlier's Principle, the equilibrium shifts in backward direction, and thus gases becomes less soluble in liquids.

2.10. State Henry's law and mention some of its important applications.

Sol:

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

Henry's law: The solubility of a gas in a liquid at a particular temperature is directly proportional to the pressure of the gas in equilibrium with the liquid at that temperature.

or

The partial pressure of a gas in vapour phase is proportional to the mole fraction of the gas (x) in the solution. $p = K_H X$

where K_H is Henry's law constant.

Applications of Henry's law :

(i) In order to increase the solubility of CO_2 gas in soft drinks and soda water, the bottles are normally sealed under high pressure. Increase in pressure increases the solubility of a gas in a solvent according to Henry's Law. If the bottle is opened by removing the stopper or seal, the pressure on the surface of the gas will suddenly decrease. This will cause a decrease in the solubility of the gas in the liquid i.e. water. As a result, it will rush out of the bottle producing a hissing noise or with a fiz.

(ii) As pointed above, oxygen to be used by deep sea divers is generally diluted with helium in order to reduce or minimise the painful effects during decompression.

(iii) As the partial pressure of oxygen in air is high, in lungs it combines with haemoglobin to form oxyhaemoglobin. In tissues, the partial pressure of oxygen is comparatively low. Therefore, oxyhaemoglobin releases oxygen in order to carry out cellular activities.

2.11. The partial pressure of ethane over a solution containing 6.56×10^{-3} g of ethane is 1 bar. If the solution contains 5.00×10^{-2} g of ethane, then what shall be the partial pressure of the gas?

Sol:

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

We know that, $m = K_H \times P$

$$\therefore 6.56 \times 10^{-2} \text{ g} = K_H \times 1 \text{ bar} \quad \dots(i)$$

$$\therefore 5.00 \times 10^{-2} \text{ g} = K_H \times P \quad \dots(ii)$$

$$K_H = 6.56 \times 10^{-2} / 1 \text{ bar (from i)}$$

$$K_H = 5.00 \times 10^{-2} / P \text{ bar (from ii),}$$

$$\therefore \frac{6.56 \times 10^{-2}}{1} = \frac{5.00 \times 10^{-2}}{P}$$

$$\therefore P = \frac{5.00}{6.56} = 0.762 \text{ bar.}$$

2.13. According to Raoult's law, what is meant by positive and negative deviations and how is the sign of $\Delta_{\text{sol}}H$ related to positive and negative deviations from Raoult's law?

Sol: Solutions having vapour pressures more than that expected from Raoult's law are said to exhibit positive deviation. In these solutions solvent – solute interactions are weaker and $\Delta_{\text{sol}}H$ is positive because stronger A – A or B – B interactions are replaced by weaker A – B interactions. Breaking of the stronger interactions requires more energy & less energy is released on formation of weaker interactions. So overall $\Delta_{\text{sol}}H$ is positive. Similarly $\Delta_{\text{sol}}V$ is positive i.e. the volume of solution is somewhat more than sum of volumes of solvent and solute.

So there is expansion in volume on solution formation.

Similarly in case of solutions exhibiting negative deviations, A – B interactions are stronger than A-A&B-B. So weaker interactions are replaced by stronger interactions so, there is release of energy i.e. $\Delta_{\text{sol}}H$ is negative.

2.14. An aqueous solution of 2 percent non-volatile solute exerts a pressure of 1.004 bar at the boiling point of the solvent. What is the molecular mass of the solute ?

Sol:

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

According to Raoult's Law,

$$\frac{P_A^\circ - P_S}{P_S} = \frac{n_B}{n_A} = \frac{W_B}{M_B} \times \frac{M_A}{W_A}$$

P_A° (for water) = 1.013 bar ; P_S = 1.004 bar ; W_B = 2 g ; W_A = 100 - 2 = 98 g ;
 M_A = 18 g mol⁻¹.

$$\frac{(1.013 - 1.004) \text{ bar}}{(1.004 \text{ bar})} = \frac{(2 \text{ g}) \times (18 \text{ g mol}^{-1})}{M_B \times (98 \text{ g})}$$

$$\therefore M_B = \frac{(2 \text{ g}) \times (18 \text{ g mol}^{-1}) \times (1.004 \text{ bar})}{(0.009 \text{ bar}) \times (98 \text{ g})} = 41.0 \text{ g mol}^{-1}$$

NCERT 12th Chemistry Chapter 2

2.15 Heptane and octane form an ideal solution. At 373 K, the vapour pressures of the two liquid components are 105.2 kPa and 46.8 kPa respectively. What will be the vapour pressure of a mixture of 26.0 g of heptane and 35.0 g of octane?

Sol.

Molar mass of heptane (C₇H₁₆)
 = 7 × 12 + 16 = 100 g mol⁻¹

Molar mass of octane (C₈H₁₈)
 = 8 × 12 + 18 = 114 g mol⁻¹

Moles of heptane present in mixture
 = $\frac{26.0}{100} = 0.26 \text{ mol}$

Moles of octane present in mixture
 = $\frac{35.0}{114} = 0.307 \text{ mol}$

Mole fraction of heptane x_H

$$= \frac{0.26}{0.26 + 0.307} = 0.458$$

Mole fraction of octane, $x_O = (1 - 0.458) = 0.542$

Vapour pressure of heptane = $x_H \times P^\circ$
 = 0.458 × 105.2 kPa = 48.18 kPa

Vapour pressure of octane = $x_O \times P^\circ$
 = 0.542 × 46.8 kPa = 25.36 kPa

Vapour pressure of mixture
 = 48.18 + 25.36 = 73.54 kPa

2.16. The vapour pressure of water is 12.3 kPa at 300 K. Calculate vapour pressure of 1 molal solution of a non-volatile solute in it

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

Sol: 1 molal solution of solute means 1 mole of solute in 1000g of the solvent.

Mole fraction of heptane x_H

$$= \frac{0.26}{0.26 + 0.307} = 0.458$$

Mole fraction of octane, $x_O = (1 - 0.458) = 0.542$

Vapour pressure of heptane $= x_H \times P^\circ$

$$= 0.458 \times 105.2 \text{ kPa} = 48.18 \text{ kPa}$$

Vapour pressure of octane $= x_O \times P^\circ$

$$= 0.542 \times 46.8 \text{ kPa} = 25.36 \text{ kPa}$$

Vapour pressure of mixture

$$= 48.18 + 25.36 = 73.54 \text{ kPa}$$

2.17. Calculate the mass of a non-volatile solute (molecular mass 40 g mol^{-1}) that should be dissolved in 114 g of octane to reduce its pressure to 80%. (C.B.S.E. Outside Delhi 2008)

Sol: According to Raoult's Law,

$$\frac{P_A^\circ - P_S}{P_S} = \frac{n_B}{n_A} = \frac{W_B}{M_B} \times \frac{M_A}{W_A}$$

Let $P_A^\circ = 1 \text{ atm}$, $P_S = 0.8 \text{ atm}$; $P_A^\circ - P_S = 0.2 \text{ atm}$; $M_B = 40 \text{ g mol}^{-1}$; $W_A = 114 \text{ g}$; $M_A (\text{C}_8\text{H}_{18}) = 114 \text{ g mol}^{-1}$.

$$\begin{aligned} W_B &= \frac{(P_A^\circ - P_S)}{P_S} \times \frac{M_B \times W_A}{M_A} \\ &= \frac{(0.2 \text{ atm})}{(0.8 \text{ atm})} \times \frac{(40 \text{ g mol}^{-1}) \times (114 \text{ g})}{(114 \text{ g mol}^{-1})} = 10.0 \text{ g.} \end{aligned}$$

2.18. A solution containing 30g of non-volatile solute exactly in 90 g of water has a vapour pressure of 2.8 kPa at 298 K. Further, 18g of water is then added to the solution and the new of vapour pressure becomes 2.9 kPa at 298 K. Calculate

(i) molar mass of the solute.

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

(ii) vapour pressure of water at 298 K.

Sol: Let the molar mass of solute = $M \text{ g mol}^{-1}$

∴ Moles of solute present

$$= \frac{30\text{g}}{M \text{ g mol}^{-1}} = \frac{30}{M} \text{ mol}$$

Moles of solvent present, $(n_1) = \frac{90}{18} = 5$ moles.

$$\therefore \frac{P^\circ - P_s}{P^\circ} = \frac{n_2}{n_1 + n_2}$$

$$\frac{P^\circ - 2.8}{P^\circ} = \frac{30/M}{5 + 30/M}$$

$$1 - \frac{2.8}{P^\circ} = \frac{30}{(5M + 30)}$$

$$1 - \frac{30}{5M + 30} = \frac{2.8}{P^\circ}$$

$$1 - \frac{6}{M + 6} = \frac{2.8}{P^\circ}$$

$$\frac{M + 6 - 6}{M + 6} = \frac{2.8}{P^\circ}$$

$$\frac{M}{M + 6} = \frac{2.8}{P^\circ}$$

$$\frac{P^\circ}{2.8} = 1 + \frac{6}{M} \quad \dots(i)$$

After adding 18 g of water,
Moles of water becomes

$$= \frac{90 + 18}{18} = \frac{108}{18} = 6 \text{ moles}$$

$$\therefore \frac{P^\circ - P_s}{P^\circ} = \frac{30/M}{6 + 30/M}$$

P_s New vapour pressure = 2.9 kPa

$$\frac{P^\circ - 2.9}{P^\circ} = \frac{30 M}{M (6M + 30)} = \frac{5}{M + 5}$$

$$1 - \frac{2.9}{P^\circ} = \frac{5}{M + 5}$$

$$1 - \frac{5}{M + 5} = \frac{2.9}{P^\circ}$$

$$\frac{M + 5 - 5}{M + 5} = \frac{2.9}{P^\circ}$$

$$\frac{P^\circ}{2.9} = \frac{M + 5}{M} \Rightarrow = 1 + \frac{5}{M}$$

$$\frac{P^\circ}{2.9} = 1 + \frac{5}{M}$$

...(ii)

Dividing equation (i) by (ii), we get,

$$\frac{2.9}{2.8} = \frac{1 + 6/M}{1 + 5/M}$$

$$2.9 \left(1 + \frac{5}{M}\right) = 2.8 \left(1 + \frac{6}{M}\right)$$

$$2.9 + \frac{2.9 \times 5}{M} = 2.8 + \frac{2.8 \times 6}{M}$$

$$2.9 + \frac{14.5}{M} = 2.8 + \frac{16.8}{M}$$

$$0.1 = \frac{16.8}{M} - \frac{14.5}{M} = \frac{2.3}{M}$$

$$M = \frac{2.3}{0.1}$$

$$M = 23 \text{ g mol}^{-1}$$

Putting $M = 23$, in equation (i), we get,

$$\frac{P^\circ}{2.8} = 1 + \frac{6}{23} = \frac{29}{23}$$

$$P^\circ = \frac{29}{23} \times 2.8 = 3.53 \text{ kPa.}$$

NCERT 12th Chemistry Chapter 2

2.19. A 5% solution (by mass) of cane sugar in water has freezing point of 271 K. Calculate the freezing point of 5% glucose in water if freezing point of pure water is 273.15 K.

Sol: Mass of sugar in 5% (by mass) solution means 5g in 100g of solvent (water)

Molar mass of sugar = 342 g mol^{-1}

$$\text{Molality of sugar solution} = \frac{5 \times 1000}{342 \times 100} = 0.146$$

$$\therefore \Delta T_f \text{ for sugar solution} = 273.15 - 271 = 2.15^\circ$$

$$\Delta T_f = K_f \times m$$

$$\Delta T_f = K_f \times 0.146 \Rightarrow K_f = 2.15/0.146$$

Molality of glucose solution

$$= \frac{5}{180} \times \frac{1000}{100} = 0.278$$

(Molar mass of glucose = 180 g mol^{-1})

$$\Delta T_f = K_f \times m = \frac{2.15}{0.146} \times 0.278 = 4.09^\circ$$

$$\therefore \text{Freezing point of glucose solution} \\ = 273.15 - 4.09 = 269.06 \text{ K.}$$

2.20. Two elements A and B form compounds having formula AB_2 and AB_4 . When dissolved in 20g of benzene (C_6H_6), 1 g of AB_2 lowers the freezing point by 2.3 K whereas 1.0 g of AB_4 lowers it by 1.3 K. The molar depression constant for benzene is $5.1 \text{ K kg mol}^{-1}$. Calculate atomic masses of A and B.

Sol:

Using the relation, $M_2 = \frac{1000 \times k_f \times w_2}{w_1 \times \Delta T_f}$

$$\therefore M_{AB_2} = \frac{1000 \times 5.1 \times 1}{20 \times 2.3} = 110.87 \text{ g mol}^{-1}$$

$$M_{AB_4} = \frac{1000 \times 5.1 \times 1}{20 \times 1.3} = 196.15 \text{ g mol}^{-1}$$

Let the atomic masses of *A* and *B* are 'p' and 'q' respectively.

Then molar mass of

$$AB_2 = p + 2q = 110.87 \text{ g mol}^{-1} \dots(i)$$

And molar mass of

$$AB_4 = p + 4q = 196.15 \text{ g mol}^{-1} \dots(ii)$$

Subtracting equation (ii) from equation (i), we get $2q = 85.28 \Rightarrow q = 42.64$

Putting $q = 42.64$ in equ. (i), we get

$$p = 110.87 - 85.28$$

$$p = 25.59$$

Thus, atomic mass of *A* = 25.59 g mol⁻¹ and atomic mass of *B* = 42.64 g mol⁻¹

2.21. At 300 K, 36 g glucose present per litre in its solution has osmotic pressure of 4.98 bar. If the osmotic pressure of the solution is 1.52 bar at the same temperature, what would be its concentration?

Sol:

$$\pi = CRT = \frac{W_B \times R \times T}{M_B \times V}$$

For both the solutions, R, T and V are constants

Ist case : $(4.98 \text{ bar}) = \frac{(36 \text{ g}) \times R \times T}{(180 \text{ g mol}^{-1}) \times V}$

IInd case : $(1.52 \text{ bar}) = \frac{W_B \times R \times T}{M_B \times V}$... (ii)

Divide eqn. (ii) by eqn. (i),

$$\frac{(1.52 \text{ bar})}{(4.98 \text{ bar})} = \frac{W_B}{M_B} \times (5 \text{ mol})$$

or $\frac{W_B}{M_B} = \frac{1.52}{4.98} \times \frac{1}{(5 \text{ mol})} = 0.0610 \text{ mol}^{-1}$

2.22. Suggest the most important type of intermolecular attractive interaction in the following pairs:

(i) n-hexane and n-octane

(ii) I₂ and CCl₄.

(iii) NaClO₄ and water

(iv) methanol and acetone

(v) acetonitrile (CH₃CN) and acetone (C₃H₆O)

Sol: (i) Both n-hexane and n-octane are non-polar. Thus, the intermolecular interactions will be London dispersion forces.

(ii) Both I₂ and CCl₄ are non-polar. Thus, the intermolecular interactions will be London dispersion forces.

(iii) NaClO₄ is an ionic compound and gives Na⁺ and ClO₄⁻ ions in the solution. Water is a polar molecule. Thus, the intermolecular interactions will be ion-dipole interactions.

(iv) Both methanol and acetone are polar molecules. Thus, intermolecular interactions will be dipole-dipole interactions.

(v) Both CH_3CN and $\text{C}_3\text{H}_6\text{O}$ are polar molecules. Thus, intermolecular interactions will be dipole-dipole interactions.

2.23. Based on solute solvent interactions, arrange the following in order of increasing solubility in n-octane and explain.

Cyclohexane, KCl, CH_3OH , CH_3CN .

Sol: n-octane (C_8H_{18}) is a non-polar liquid and solubility is governed by the principle that like dissolve like. Keeping this in view, the increasing order of solubility of different solutes is:

$\text{KCl} < \text{CH}_3\text{OH} < \text{CH}_3\text{C}=\text{N} < \text{C}_6\text{H}_{12}$ (cyclohexane).

2.24. Amongst the following compounds, identify which are insoluble, partially soluble and highly soluble in water?

(i) phenol

(ii) toluene

(iii) formic acid

(iv) ethylene glycol

(v) chloroform

(vi) pentanol

Sol: (i) Phenol (having polar – OH group) – Partially soluble.

(ii) Toluene (non-polar) – Insoluble.

(iii) Formic acid (form hydrogen bonds with water molecules) – Highly soluble.

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

(iv) Ethylene glycol (form hydrogen bonds with water molecules) Highly soluble.

(v) Chloroform (non-polar)- Insoluble.

(vi) Pentanol (having polar -OH) – Partially soluble.


2.25. If the density of lake water is 1.25 g mL^{-1} , and it contains 92 g of Na^+ ions per kg of water, calculate the molality of Na^+ ions in the lake. (C.B.S.E. Outside Delhi 2008)

Sol:

$$\begin{aligned}\text{Molality of Na}^+ \text{ ions } (m) &= \frac{\text{No. of moles of Na}^+ \text{ ions}}{\text{Mass of water in kg}} \\ &= \frac{(92 \text{ g}) / (23 \text{ g mol}^{-1})}{1 \text{ kg}} = 4 \text{ mol kg}^{-1} = 4 \text{ m}\end{aligned}$$

2.26. If the solubility product of CuS is 6×10^{-16} , calculate the maximum molarity of CuS in aqueous solution.

Sol:



Maximum molarity of CuS in aqueous solution means solubility of CuS.

Let the solubility of CuS be $S \text{ mol L}^{-1}$

$$\begin{aligned}\therefore K_{sp} &= [\text{Cu}^{2+}] [\text{S}^{2-}] \\ 6 \times 10^{-16} &= S \times S = S^2\end{aligned}$$

$$\therefore S = \sqrt{6 \times 10^{-16}} = 2.45 \times 10^{-8} \text{ mol L}^{-1}.$$

2.27. Calculate the mass percentage of aspirin ($\text{C}_9\text{H}_8\text{O}_4$) in acetonitrile (CH_3CN) when 6.5 g of CHO is dissolved in 450 g of CH_3CN .

Solution:

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

Mass percentage of aspirin

$$= \frac{\text{Mass of aspirin}}{\text{Mass of aspirin} + \text{Mass of acetonitrile}} \times 100$$

$$= \frac{6.5}{6.5 + 450} \times 100 = 1.424\%$$

2.28. Nalorphene ($\text{C}_{19}\text{H}_{21}\text{NO}_3$), similar to morphine, is used to combat withdrawal symptoms in narcotic users. Dose of nalorphene generally given is 1.5 mg. Calculate the mass of 1.5×10^{-3} m aqueous solution required for the above dose.

Solution:

1.5×10^{-3} m aqueous solution of nalorphene means that 1.5×10^{-3} mole of nalorphene is dissolved in 1 kg of water.

Molar mass of nalorphene, $\text{C}_{19}\text{H}_{21}\text{NO}_3$
 $= 19 \times 12 + 21 + 14 + 3 \times 16 = 311 \text{ g mol}^{-1}$

$\therefore 1.5 \times 10^{-3}$ mole of nalorphene

$$= 1.5 \times 10^{-3} \times 311 \text{ g} = 0.467 \text{ g}$$

\therefore Mass of solution = $0.467 + 1000 = 1000.467 \text{ g}$.

For 0.467g of nalorphene, mass of solution required = 1000.467g

For 1.5 mg (1.5×10^{-3} g) of nalorphene, mass of solution required

$$= \frac{1000.467}{0.467} \times 1.5 \times 10^{-3} = 3.21 \text{ g}$$

2.29. Calculate the amount of benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$) required for preparing 250 mL of 0.15 M solution in methanol.

Solution:

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

$$\text{Molarity (M)} = \frac{\text{Mass of solute/molar mass}}{\text{Volume of solution in litres}}$$

$$M = 0.15 \text{ M} = 0.15 \text{ mol L}^{-1};$$

$$\text{Molar mass of solute} = 7 \times 12 + 6 \times 1 + 2 \times 16 = 122 \text{ g mol}^{-1};$$


$$\text{Volume of solution} = 250 \text{ mL} = 0.25 \text{ L.}$$

$$(0.15 \text{ mol L}^{-1}) = \frac{\text{Mass of solute}}{(122 \text{ g mol}^{-1}) \times (0.25 \text{ L})}$$

$$\text{Mass of solute} = (0.15 \text{ mol L}^{-1}) \times (122 \text{ g mol}^{-1}) \times (0.25 \text{ L}) = 4.575 \text{ g}$$

2.30. The depression in freezing point of water observed for the same amount of acetic acid, trichloroacetic acid and trifluoroacetic acid increases in the order given above. Explain briefly.

Solution:



Fluorine being more electronegative than chlorine has the highest electron withdrawing inductive effect. Thus, trifluoroacetic acid is the strongest trichloroacetic acid is second most and acetic acid is the weakest acid due to absence of any electron withdrawing group. Thus, F_3CCOOH ionizes to the largest extent while CH_3COOH ionizes to minimum extent in water. Greater the extent of ionization greater is the depression in freezing point. Hence, the order of depression in freezing point will be $\text{CH}_3\text{COOH} < \text{Cl}_3\text{CCOOH} < \text{F}_3\text{CCOOH}$.

2.31. Vapour pressure of water at 293 K is 17.535 mm Hg. Calculate the vapour pressure of water at 293 K when 25 g of glucose is dissolved in 450 g of water.

Solution:

According to Raoult's Law,

$$\frac{P_A^\circ - P_S}{P_S} = \frac{n_B}{n_A} \text{ or } \frac{P_A^\circ}{P_S} - 1 = \frac{n_B}{n_A}$$

$$\frac{P_A^\circ}{P_S} = 1 + \frac{n_B}{n_A} = 1 + \frac{W_B}{M_B} \times \frac{M_A}{W_A}$$

$$W_B = 25 \text{ g}; W_A = 450 \text{ g}; M_B = 180 \text{ g mol}^{-1};$$

$$M_A = 18 \text{ g mol}^{-1}; P_A^\circ = 17.535 \text{ mm}$$

$$\frac{P_A^\circ}{P_S} = 1 + \frac{(25 \text{ g}) \times (18 \text{ g mol}^{-1})}{(180 \text{ g mol}^{-1}) \times (450 \text{ g})} = 1 + 0.0055 = 1.0055$$

$$P_S (\text{V.P of water in solution}) = \frac{(17.535 \text{ mm})}{(1.0055)} = 17.44 \text{ mm}$$

2.32. Henry's law constant for the molality of methane in benzene at 298 K is 4.27×10^5 mm Hg. Calculate the solubility of methane in benzene at 298 K under 760 mm Hg.

Solution:

Using relation; $P = K_H x$

$$\therefore x = \frac{P}{K_H} = \frac{760 \text{ mm Hg}}{4.27 \times 10^5 \text{ mm Hg}} = 1.78 \times 10^{-3}$$

i.e., mole fraction of methane in benzene
 $= 1.78 \times 10^{-3}$.

2.33. 100g of liquid A (molar mass 140 g mol^{-1}) was dissolved in 1000g of liquid B (molar mass 180 g mol^{-1}). The vapour pressure of pure liquid B was found to be 500 torr. Calculate the vapour

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

pressure of pure liquid A and its vapour pressure in the solution if the total vapour pressure of the solution is 475 torr.

Solution:

| | | | | | | | | |
|--------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|
| $100 \times \chi_{\text{acetone}}$ | 0 | 11.8 | 23.4 | 36.0 | 50.8 | 58.2 | 64.5 | 72.1 |
| $P_{\text{acetone}}/\text{mm Hg}$ | 0 | 54.9 | 110.1 | 202.4 | 322.7 | 405.9 | 454.1 | 521.1 |
| $P_{\text{chloroform}}/\text{mm Hg}$ | 632.8 | 548.1 | 469.4 | 359.7 | 257.7 | 193.6 | 161.2 | 120.7 |


2.34. Vapour pressures of pure acetone and chloroform at 328 K are 741.8 mm Hg and 632.8 mm Hg respectively. Assuming that they form ideal solution over the entire range of composition, plot P_{total} , $P_{\text{chloroform}}$ and P_{acetone} as a function of χ_{acetone} . The experimental data observed for different compositions of mixtures is:

| | | | | | | | | |
|--------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|
| $100 \times \chi_{\text{acetone}}$ | 0 | 11.8 | 23.4 | 36.0 | 50.8 | 58.2 | 64.5 | 72.1 |
| $P_{\text{acetone}}/\text{mm Hg}$ | 0 | 54.9 | 110.1 | 202.4 | 322.7 | 405.9 | 454.1 | 521.1 |
| $P_{\text{chloroform}}/\text{mm Hg}$ | 632.8 | 548.1 | 469.4 | 359.7 | 257.7 | 193.6 | 161.2 | 120.7 |

Plot this data also on the same graph paper. Indicate whether it has positive deviation or negative deviation from the ideal solution.

Solution:

| | | | | | | | | |
|--------------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|
| χ_{acetone} | 0.0 | 0.118 | 0.234 | 0.360 | 0.508 | 0.582 | 0.645 | 0.721 |
| $P_{\text{acetone}}/\text{mm Hg}$ | 0 | 54.9 | 110.1 | 202.4 | 322.7 | 405.9 | 454.1 | 521.1 |
| $P_{\text{chloroform}}/\text{mm Hg}$ | 632.8 | 548.1 | 469.4 | 359.7 | 257.7 | 193.6 | 161.2 | 120.7 |
| P_{total} | 632.8 | 603.0 | 579.5 | 562.1 | 580.4 | 599.5 | 615.3 | 641.8 |



As the plot for P_{total} dips downwards, hence the solution shows negative deviation from the ideal behaviour.

Mole fraction of $\text{C}_6\text{H}_5\text{CH}_3$,

$$x_T = \frac{1.087}{1.026 + 1.087} = 0.514$$

According to Raoult's Law,

$$P_B = x_B \times P_B^\circ = 0.486 \times 50.71 = 24.65 \text{ mm}$$

$$P_T = x_T \times P_T^\circ = 0.514 \times 32.06 = 16.48 \text{ mm}$$

Mole fraction of C_6H_6 in vapour phase

$$= \frac{P_B}{P_B + P_T} = \frac{24.65}{24.65 + 16.48} = 0.599$$

2.35. Benzene and toluene form ideal solution over the entire range of composition. The vapour pressure of pure benzene and toluene at 300 K are 50.71 mm Hg and 32.06 mm Hg respectively. Calculate the mole fraction of benzene in vapour phase if 80g of benzene is mixed with 100g of toluene.

Solution:

$$\text{Molar mass of } \text{C}_6\text{H}_6 = 78 \text{ g mol}^{-1}$$

$$\text{Molar mass of } \text{C}_6\text{H}_5\text{CH}_3 = 92 \text{ g mol}^{-1}$$

$$\text{No. of moles of } \text{C}_6\text{H}_6 = \frac{80}{78} = 1.026 \text{ mole}$$

$$\text{No. of moles of } \text{C}_6\text{H}_5\text{CH}_3 = \frac{100}{92} = 1.087 \text{ mole}$$

Mole fraction of C_6H_6 ,

$$x_B = \frac{1.026}{1.026 + 1.087} = 0.486$$

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

2.36. The air is a mixture of a number of gases. The major components are oxygen and nitrogen with an approximate proportion of 20% is to 79% by volume at 298 K. The water is in equilibrium with air at a pressure of 10 atm. At 298 K if Henry's law constants for oxygen and nitrogen are 3.30×10^7 mm and 6.51×10^7 mm respectively, calculate the composition of these gases in water.

Solution:

Air containing 20% oxygen and 79% nitrogen by volume means

$$\begin{aligned} \text{Partial pressure of O}_2 (P_{\text{O}_2}) &= \frac{20}{100} \times 10 = 2 \text{ atm} \\ &= 2 \times 760 \text{ mm} = 1520 \text{ mm} \end{aligned}$$

$$\begin{aligned} \text{Partial pressure of N}_2 (P_{\text{N}_2}) &= \frac{79 \times 10}{100} = 7.9 \text{ atm} \\ &= 7.9 \times 760 \text{ mm} = 6004 \text{ mm} \end{aligned}$$

2.37. Determine the amount of CaCl_2 ($i = 2.47$) dissolved in 2.5 litre of water such that its osmotic pressure is 0.75 atm at 27°C .

Solution:

Using relation, $\pi = iCRT = i \frac{n}{V} RT$

$$n = \frac{\pi V}{i RT} = \frac{0.75 \times 2.5}{2.47 \times 0.0821 \times 300} = 0.0308 \text{ mole}$$

Using relation, $\pi = iCRT = i \frac{n}{V} RT$


$$n = \frac{\pi V}{i RT} = \frac{0.75 \times 2.5}{2.47 \times 0.0821 \times 300} = 0.0308 \text{ mole}$$

2.38. Determine the osmotic pressure of a solution prepared by dissolving 25 mg of K_2SO_4 in 2 litre of water at $25^\circ C$, assuming that it is completely dissociated. (C.B.S.E. 2013)

Solution:

Step I. Calculation of Van't Hoff factor (i)

K_2SO_4 dissociates in water as :



$$\alpha(\text{for complete dissociation}) = 1, n = 3 ; 1 = \frac{i-1}{3-1} \text{ or } i = 2 + 1 = 3$$

Step II. Calculation of osmotic pressure (π)

$$\text{Osmotic pressure } (\pi) = i C R T = \frac{i W_B R T}{M_B \times V}$$

$$i = 3 ; W_B = 25 \text{ mg} = 0.025 \text{ g} ; M_B = 2 \times 39 + 32 + 4 \times 16 = 174 \text{ g mol}^{-1} ; V = 2 \text{ L} ; T = 25^\circ C = 298 \text{ K} ; R = 0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}$$

$$\begin{aligned} \pi &= \frac{(3) \times (0.025 \text{ g}) \times (0.0821 \text{ L atm K}^{-1} \text{ mol}^{-1}) \times (298 \text{ K})}{(174 \text{ g mol}^{-1}) \times (2 \text{ L})} \\ &= 5.27 \times 10^{-3} \text{ atm.} \end{aligned}$$



Chapterwise NCERT Solutions for Class 12 Chemistry:

- Chapter 1 : The Solid State
- Chapter 2 : Solutions
- Chapter 3
Electrochemistry
- Chapter 4 : Chemical
Kinetics
- Chapter 5 : Surface
chemistry
- Chapter 6 : General
Principles and Processes of
Isolation of Elements
- Chapter 7 : The p Block
Elements
- Chapter 8 : The d and f
Block Elements
- Chapter 9 : Coordination
Compounds
- Chapter 10 : Haloalkanes
and Haloarenes
- Chapter 11 : Alcohols
Phenols and Ether
- Chapter 12 : Aldehydes
Ketones and Carboxylic
Acids
- Chapter 13 : Amines
- Chapter 14 : Biomolecules
- Chapter 15 : Polymers
- Chapter 16 : Chemistry in
Everyday Life

<https://www.indcareer.com/schools/ncert-solutions-for-12th-class-chemistry-chapter-2-solutions/>

About NCERT

The National Council of Educational Research and Training is an autonomous organization of the Government of India which was established in 1961 as a literary, scientific, and charitable Society under the Societies Registration Act. The major objectives of NCERT and its constituent units are to: undertake, promote and coordinate research in areas related to school education; prepare and publish model textbooks, supplementary material, newsletters, journals and develop educational kits, multimedia digital materials, etc.

Organise pre-service and in-service training of teachers; develop and disseminate innovative educational techniques and practices; collaborate and network with state educational departments, universities, NGOs and other educational institutions; act as a clearing house for ideas and information in matters related to school education; and act as a nodal agency for achieving the goals of Universalisation of Elementary Education.

Its headquarters are located at Sri Aurobindo Marg in New Delhi.

[Visit the Official NCERT website](https://www.ncert.nic.in/) to learn more.