


Class 12 - Chapter 15 Mean Value Theorems



RD Sharma Solutions for Class 12 Maths Chapter 15–Mean Value Theorems

Class 12: Maths Chapter 15 solutions. Complete Class 12 Maths Chapter 15 Notes.

RD Sharma Solutions for Class 12 Maths Chapter 15–Mean Value Theorems

RD Sharma 12th Maths Chapter 15, Class 12 Maths Chapter 15 solutions

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>



Exercise 15.1 Page No: 15.9

1. Discuss the applicability of Rolle's Theorem for the following functions on the indicated intervals:

(i) $f(x) = 3 + (x - 2)^{\frac{2}{3}}$ on $[1, 3]$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is

$$\Rightarrow f(x) = 3 + (x - 2)^{\frac{2}{3}} \text{ on } [1, 3]$$

Let us check the differentiability of the function $f(x)$.

Now we have to find the derivative of $f(x)$,

$$\Rightarrow f'(x) = \frac{d}{dx} \left(3 + (x - 2)^{\frac{2}{3}} \right)$$

$$\Rightarrow f'(x) = \frac{d(3)}{dx} + \frac{d\left((x-2)^{\frac{2}{3}}\right)}{dx}$$

$$\Rightarrow f'(x) = 0 + \frac{2}{3} (x - 2)^{\frac{2}{3} - 1}$$

$$\Rightarrow f'(x) = \frac{2}{3} (x - 2)^{-\frac{1}{3}}$$

$$\Rightarrow f'(x) = \frac{2}{3(x-2)^{\frac{1}{3}}}$$

Now we have to check differentiability at the value of $x = 2$

$$\Rightarrow \lim_{x \rightarrow 2} f'(x) = \lim_{x \rightarrow 2} \frac{2}{3(x-2)^{\frac{1}{3}}}$$

$$\Rightarrow \lim_{x \rightarrow 2} f'(x) = \frac{2}{3(2-2)^{\frac{1}{3}}}$$

$$\Rightarrow \lim_{x \rightarrow 2} f'(x) = \frac{2}{3(0)}$$

$$\Rightarrow \lim_{x \rightarrow 2} f'(x) = \text{undefined}$$

\therefore f is not differentiable at $x = 2$, so it is not differentiable in the closed interval $(1, 3)$.

So, Rolle's theorem is not applicable for the function f on the interval $[1, 3]$.

(ii) $f(x) = [x]$ for $-1 \leq x \leq 1$, where $[x]$ denotes the greatest integer not exceeding x

Solution:

Given function is $f(x) = [x]$, $-1 \leq x \leq 1$ where $[x]$ denotes the greatest integer not exceeding x .

Let us check the continuity of the function f .

Here in the interval $x \in [-1, 1]$, the function has to be Right continuous at $x = 1$ and left continuous at $x = -1$.

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} [x]$$

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+ + h} [x] \text{ Where } h > 0.$$

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = \lim_{h \rightarrow 0} 1$$

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = 1 \text{ (1)}$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} [x]$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^- - h} [x], \text{ where } h > 0$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = \lim_{h \rightarrow 0} 0$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = 0 \text{ (2)}$$

From (1) and (2), we can see that the limits are not the same so, the function is not continuous in the interval $[-1, 1]$.

\therefore Rolle's Theorem is not applicable for the function f in the interval $[-1, 1]$.

$$\text{(iii) } f(x) = \sin \frac{1}{x} \text{ for } -1 \leq x \leq 1$$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is $f(x) = \sin\left(\frac{1}{x}\right)$ for $-1 \leq x \leq 1$

Let us check the continuity of the function 'f' at the value of $x = 0$. We cannot directly find the value of limit at $x = 0$, as the function is not valid at $x = 0$. So, we take the limit on either sides of $x = 0$, and we check whether they are equal or not.

So consider RHL:

$$\Rightarrow \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \sin\left(\frac{1}{x}\right)$$

We assume that the limit $\lim_{h \rightarrow 0} \sin\left(\frac{1}{h}\right) = k$, $k \in [-1, 1]$.

$$\Rightarrow \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+ h} \sin\left(\frac{1}{x}\right), \text{ where } h > 0$$

$$\Rightarrow \lim_{x \rightarrow 0^+} f(x) = \lim_{h \rightarrow 0} \sin\left(\frac{1}{h+0}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^+} f(x) = \lim_{h \rightarrow 0} \sin\left(\frac{1}{h}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^+} f(x) = k \quad \dots\dots (1)$$

Now consider LHL:

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \sin\left(\frac{1}{x}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^- h} \sin\left(\frac{1}{x}\right), \text{ where } h > 0$$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0} \sin\left(\frac{1}{0-h}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0} \sin\left(\frac{1}{-h}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = \lim_{h \rightarrow 0} -\sin\left(\frac{1}{h}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = -\lim_{h \rightarrow 0} \sin\left(\frac{1}{h}\right)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} f(x) = -k \dots\dots (2)$$

From (1) and (2), we can see that the Right hand and left – hand limits are not equal, so the function 'f' is not continuous at $x = 0$.

\therefore Rolle's Theorem is not applicable to the function 'f' in the interval $[-1, 1]$.

(iv) $f(x) = 2x^2 - 5x + 3$ on $[1, 3]$

Solution:

Given function is $f(x) = 2x^2 - 5x + 3$ on $[1, 3]$

Since given function f is a polynomial. So, it is continuous and differentiable everywhere.

Now, we find the values of function at the extreme values.

$$\Rightarrow f(1) = 2(1)^2 - 5(1) + 3$$

$$\Rightarrow f(1) = 2 - 5 + 3$$

$$\Rightarrow f(1) = 0 \dots\dots (1)$$

$$\Rightarrow f(3) = 2(3)^2 - 5(3) + 3$$

$$\Rightarrow f(3) = 2(9) - 15 + 3$$

$$\Rightarrow f(3) = 18 - 12$$

$$\Rightarrow f(3) = 6 \dots\dots (2)$$

From (1) and (2), we can say that, $f(1) \neq f(3)$

\therefore Rolle's Theorem is not applicable for the function f in interval $[1, 3]$.

(v) $f(x) = x^{2/3}$ on $[-1, 1]$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Solution:

Given function is $f(x) = x^{\frac{2}{3}}$ on $[-1, 1]$

Now we have to find the derivative of the given function:

$$\Rightarrow f'(x) = \frac{d\left(x^{\frac{2}{3}}\right)}{dx}$$

$$\Rightarrow f'(x) = \frac{2}{3}x^{\frac{2}{3}-1}$$

$$\Rightarrow f'(x) = \frac{2}{3}x^{\frac{2}{3}-1}$$

$$\Rightarrow f'(x) = \frac{2}{3}x^{-\frac{1}{3}}$$

$$\Rightarrow f'(x) = \frac{2}{3x^{\frac{1}{3}}}$$

Now we have to check the differentiability of the function at $x = 0$.

$$\Rightarrow \lim_{x \rightarrow 0} f'(x) = \lim_{x \rightarrow 0} \frac{2}{3x^{\frac{1}{3}}}$$

$$\Rightarrow \lim_{x \rightarrow 0} f'(x) = \frac{2}{3(0)^{\frac{1}{3}}}$$

$$\Rightarrow \lim_{x \rightarrow 0} f'(x) = \text{undefined}$$

Since the limit for the derivative is undefined at $x = 0$, we can say that f is not differentiable at $x = 0$.

\therefore Rolle's Theorem is not applicable to the function ' f ' on $[-1, 1]$.

$$(vi) f(x) = \begin{cases} -4x + 5, & 0 \leq x \leq 1 \\ 2x - 3, & 1 < x \leq 2 \end{cases}$$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is $f(x) = \begin{cases} -4x + 5, & 0 \leq x \leq 1 \\ 2x - 3, & 1 < x \leq 2 \end{cases}$

Now we have to check the continuity at $x = 1$ as the equation of function changes.

Consider LHL:

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} -4x + 5$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = -4(1) + 5$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = 1 \quad \dots\dots (1)$$

$$\Rightarrow \lim_{x \rightarrow 1^-} f(x) = 1 \quad \dots\dots (1)$$

Now consider RHL:

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} 2x - 3$$

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = 2(1) - 3$$

$$\Rightarrow \lim_{x \rightarrow 1^+} f(x) = -1 \quad \dots\dots (2)$$

From (1) and (2), we can see that the values of both side limits are not equal. So, the function 'f' is not continuous at $x = 1$.

\therefore Rolle's Theorem is not applicable to the function 'f' in the interval $[0, 2]$.

2. Verify the Rolle's Theorem for each of the following functions on the indicated intervals:

(i) $f(x) = x^2 - 8x + 12$ on $[2, 6]$

Solution:

Given function is $f(x) = x^2 - 8x + 12$ on $[2, 6]$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Since, given function f is a polynomial it is continuous and differentiable everywhere i.e., on \mathbb{R} .

Let us find the values at extremes:

$$\Rightarrow f(2) = 2^2 - 8(2) + 12$$

$$\Rightarrow f(2) = 4 - 16 + 12$$

$$\Rightarrow f(2) = 0$$

$$\Rightarrow f(6) = 6^2 - 8(6) + 12$$

$$\Rightarrow f(6) = 36 - 48 + 12$$

$$\Rightarrow f(6) = 0$$

$\therefore f(2) = f(6)$, Rolle's theorem applicable for function f on $[2,6]$.

Now we have to find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(x^2 - 8x + 12)}{dx}$$

$$\Rightarrow f'(x) = \frac{d(x^2)}{dx} - \frac{d(8x)}{dx} + \frac{d(12)}{dx}$$

$$\Rightarrow f'(x) = 2x - 8 + 0$$

$$\Rightarrow f'(x) = 2x - 8 + 0$$

$$\Rightarrow f'(x) = 2x - 8$$

We have $f'(c) = 0 \in [2, 6]$, from the above definition

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 2c - 8 = 0$$

$$\Rightarrow 2c = 8$$

$$\Rightarrow c = \frac{8}{2}$$

$$\Rightarrow c = 4 \in [2, 6]$$

\therefore Rolle's Theorem is verified.

(ii) $f(x) = x^2 - 4x + 3$ on $[1, 3]$

Solution:

Given function is $f(x) = x^2 - 4x + 3$ on $[1, 3]$

Since, given function f is a polynomial it is continuous and differentiable everywhere i.e., on \mathbb{R} .
Let us find the values at extremes:

$$\Rightarrow f(1) = 1^2 - 4(1) + 3$$

$$\Rightarrow f(1) = 1 - 4 + 3$$

$$\Rightarrow f(1) = 0$$

$$\Rightarrow f(3) = 3^2 - 4(3) + 3$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow f(3) = 9 - 12 + 3$$

$$\Rightarrow f(3) = 0$$

$\therefore f(1) = f(3)$, Rolle's theorem applicable for function 'f' on [1,3].

Let's find the derivative of f(x)

$$\Rightarrow f'(x) = \frac{d(x^2 - 4x + 3)}{dx}$$

$$\Rightarrow f'(x) = \frac{d(x^2)}{dx} - \frac{d(4x)}{dx} + \frac{d(3)}{dx}$$

$$\Rightarrow f'(x) = 2x - 4 + 0$$

$$\Rightarrow f'(x) = 2x - 4$$

We have $f'(c) = 0$, $c \in (1, 3)$, from the definition of Rolle's Theorem.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 2c - 4 = 0$$

$$\Rightarrow 2c = 4$$

$$\Rightarrow c = \frac{4}{2}$$

$$\Rightarrow c = 2 \in (1, 3)$$

\therefore Rolle's Theorem is verified.

$$\Rightarrow f'(x) = 2x - 4$$

We have $f'(c) = 0$, $c \in (1, 3)$, from the definition of Rolle's Theorem.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 2c - 4 = 0$$

$$\Rightarrow 2c = 4$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow c = 4/2$$

$$\Rightarrow C = 2 \in (1, 3)$$

\therefore Rolle's Theorem is verified.

(iii) $f(x) = (x - 1)(x - 2)^2$ on $[1, 2]$

Solution:

Given function is $f(x) = (x - 1)(x - 2)^2$ on $[1, 2]$

Since, given function f is a polynomial it is continuous and differentiable everywhere that is on \mathbb{R} .

Let us find the values at extremes:

$$\Rightarrow f(1) = (1 - 1)(1 - 2)^2$$

$$\Rightarrow f(1) = 0(1)^2$$

$$\Rightarrow f(1) = 0$$

$$\Rightarrow f(2) = (2 - 1)(2 - 2)^2$$

$$\Rightarrow f(2) = 0^2$$

$$\Rightarrow f(2) = 0$$

$\therefore f(1) = f(2)$, Rolle's Theorem applicable for function ' f ' on $[1, 2]$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d((x-1)(x-2)^2)}{dx}$$

Differentiating by using product rule, we get

$$\Rightarrow f'(x) = (x-2)^2 \times \frac{d(x-1)}{dx} + (x-1) \times \frac{d((x-2)^2)}{dx}$$

$$\Rightarrow f'(x) = ((x-2)^2 \times 1) + ((x-1) \times 2 \times (x-2))$$

$$\Rightarrow f'(x) = x^2 - 4x + 4 + 2(x^2 - 3x + 2)$$

$$\Rightarrow f'(x) = 3x^2 - 10x + 8$$

We have $f'(c) = 0$ $c \in (1, 2)$, from the definition of Rolle's Theorem.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 3c^2 - 10c + 8 = 0$$

$$\Rightarrow c = \frac{10 \pm \sqrt{(-10)^2 - (4 \times 3 \times 8)}}{2 \times 3}$$

$$\Rightarrow c = \frac{10 \pm \sqrt{100 - 96}}{6}$$

$$\Rightarrow c = \frac{10 \pm 2}{6}$$

$$\Rightarrow 3c^2 - 10c + 8 = 0$$

$$\Rightarrow c = \frac{10 \pm \sqrt{(-10)^2 - (4 \times 3 \times 8)}}{2 \times 3}$$

$$\Rightarrow c = \frac{10 \pm \sqrt{100 - 96}}{6}$$

$$\Rightarrow c = \frac{10 \pm 2}{6}$$

$$\Rightarrow c = \frac{12}{6} \text{ or } c = \frac{8}{6}$$

$$\Rightarrow c = \frac{4}{3} \in (1, 2) \text{ (neglecting the value 2)}$$

\therefore Rolle's Theorem is verified.

(iv) $f(x) = x(x - 1)^2$ on $[0, 1]$

Solution:

Given function is $f(x) = x(x - 1)^2$ on $[0, 1]$

Since, given function f is a polynomial it is continuous and differentiable everywhere that is, on \mathbb{R} .

Let us find the values at extremes

$$\Rightarrow f(0) = 0(0 - 1)^2$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(1) = 1(1 - 1)^2$$

$$\Rightarrow f(1) = 0^2$$

$$\Rightarrow f(1) = 0$$

$\therefore f(0) = f(1)$, Rolle's theorem applicable for function ' f ' on $[0, 1]$.

Let's find the derivative of $f(x)$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow f'(x) = \frac{d(x(x-1)^2)}{dx}$$

Differentiating using product rule:

$$\Rightarrow f'(x) = (x-1)^2 \times \frac{d(x)}{dx} + x \frac{d((x-1)^2)}{dx}$$

$$\Rightarrow f'(x) = ((x-1)^2 \times 1) + (x \times 2 \times (x-1))$$

$$\Rightarrow f'(x) = (x-1)^2 + 2(x^2 - x)$$

$$\Rightarrow f'(x) = x^2 - 2x + 1 + 2x^2 - 2x$$

$$\Rightarrow f'(x) = 3x^2 - 4x + 1$$

We have $f'(c) = 0$ $c \in (0, 1)$, from the definition given above.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow f'(x) = (x - 1)^2 + 2(x^2 - x)$$

$$\Rightarrow f'(x) = x^2 - 2x + 1 + 2x^2 - 2x$$

$$\Rightarrow f'(x) = 3x^2 - 4x + 1$$

We have $f'(c) = 0$ $c \in (0, 1)$, from the definition given above.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 3c^2 - 4c + 1 = 0$$

$$\Rightarrow c = \frac{4 \pm \sqrt{(-4)^2 - (4 \times 3 \times 1)}}{2 \times 3}$$

$$\Rightarrow c = \frac{4 \pm \sqrt{16 - 12}}{6}$$

$$\Rightarrow c = \frac{4 \pm \sqrt{4}}{6}$$

$$\Rightarrow c = \frac{6}{6} \text{ or } c = \frac{2}{6}$$

$$\Rightarrow c = \frac{1}{3} \in (0, 1)$$

\therefore Rolle's Theorem is verified.

(v) $f(x) = (x^2 - 1)(x - 2)$ on $[-1, 2]$

Solution:

Given function is $f(x) = (x^2 - 1)(x - 2)$ on $[-1, 2]$

Since, given function f is a polynomial it is continuous and differentiable everywhere that is on \mathbb{R} .

Let us find the values at extremes:

$$\Rightarrow f(-1) = ((-1)^2 - 1)(-1 - 2)$$

$$\Rightarrow f(-1) = (1 - 1)(-3)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow f(-1) = (0)(-3)$$

$$\Rightarrow f(-1) = 0$$

$$\Rightarrow f(2) = (2^2 - 1)(2 - 2)$$

$$\Rightarrow f(2) = (4 - 1)(0)$$

$$\Rightarrow f(2) = 0$$

$\therefore f(-1) = f(2)$, Rolle's theorem applicable for function f on $[-1, 2]$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d((x^2-1)(x-2))}{dx}$$

Differentiating using product rule,

$$\Rightarrow f'(x) = (x-2) \times \frac{d(x^2-1)}{dx} + (x^2-1) \frac{d(x-2)}{dx}$$

$$\Rightarrow f'(x) = ((x-2) \times 2x) + ((x^2-1) \times 1)$$

$$\Rightarrow f'(x) = 2x^2 - 4x + x^2 - 1$$

$$\Rightarrow f'(x) = 2x^2 - 4x - 1$$

We have $f'(c) = 0$ $c \in (-1, 2)$, from the definition of Rolle's Theorem.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 2c^2 - 4c - 1 = 0$$

$$\Rightarrow c = \frac{4 \pm \sqrt{(-4)^2 - (4 \times 2 \times -1)}}{2 \times 2}$$

$$\Rightarrow c = \frac{4 \pm \sqrt{16 + 8}}{4}$$

$$\Rightarrow c = \frac{4 \pm \sqrt{24}}{4}$$

$$\Rightarrow c = \frac{4 + 2\sqrt{6}}{4} \text{ or } c = \frac{4 - 2\sqrt{6}}{4}$$

$$\Rightarrow c = 1 + \frac{\sqrt{6}}{2} \text{ or } c = 1 - \frac{\sqrt{6}}{2}$$

$$f'(x) = 3x^2 - 4x - 1$$

We have $f'(c) = 0$ $c \in (-1, 2)$, from the definition of Rolle's Theorem

$$f'(c) = 0$$

$$3c^2 - 4c - 1 = 0$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$c = 4 \pm \sqrt{(-4)^2 - (4 \times 3 \times -1)} / (2 \times 3) \text{ [Using the Quadratic Formula]}$$

$$c = 4 \pm \sqrt{16 + 12} / 6$$

$$c = (4 \pm \sqrt{28}) / 6$$

$$c = (4 \pm 2\sqrt{7}) / 6$$

$$c = (2 \pm \sqrt{7}) / 3 = 1.5 \pm \sqrt{7}/3$$

$$c = 1.5 + \sqrt{7}/3 \text{ or } 1.5 - \sqrt{7}/3$$

So,

$$c = 1.5 - \sqrt{7}/3 \text{ since } c \in (-1, 2)$$

\therefore Rolle's Theorem is verified.

(vi) $f(x) = x(x - 4)^2$ on $[0, 4]$

Solution:

Given function is $f(x) = x(x - 4)^2$ on $[0, 4]$

Since, given function f is a polynomial it is continuous and differentiable everywhere i.e., on \mathbb{R} .

Let us find the values at extremes:

$$\Rightarrow f(0) = 0(0 - 4)^2$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(4) = 4(4 - 4)^2$$

$$\Rightarrow f(4) = 4(0)^2$$

$$\Rightarrow f(4) = 0$$

$\therefore f(0) = f(4)$, Rolle's theorem applicable for function ' f ' on $[0, 4]$.

Let's find the derivative of $f(x)$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow f'(x) = \frac{d(x(x-4)^2)}{dx}$$

Differentiating using product rule

$$\Rightarrow f'(x) = (x-4)^2 \times \frac{d(x)}{dx} + x \frac{d((x-4)^2)}{dx}$$

$$\Rightarrow f'(x) = ((x-4)^2 \times 1) + (x \times 2 \times (x-4))$$

$$\Rightarrow f'(x) = (x-4)^2 + 2(x^2 - 4x)$$

$$\Rightarrow f'(x) = x^2 - 8x + 16 + 2x^2 - 8x$$

$$\Rightarrow f'(x) = 3x^2 - 16x + 16$$

We have $f'(c) = 0$ $c \in (0, 4)$, from the definition of Rolle's Theorem.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 3c^2 - 16c + 16 = 0$$

$$\Rightarrow c = \frac{16 \pm \sqrt{(-16)^2 - (4 \times 3 \times 16)}}{2 \times 3}$$

$$\Rightarrow c = \frac{16 \pm \sqrt{256 - 192}}{6}$$

$$\Rightarrow c = \frac{16 \pm \sqrt{64}}{6}$$

$$\Rightarrow c = \frac{8}{6} \text{ or } c = \frac{24}{6}$$

$$\Rightarrow c = \frac{8}{6} \in (0, 4)$$

\therefore Rolle's Theorem is verified.

(vii) $f(x) = x(x-2)^2$ on $[0, 2]$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is $f(x) = x(x - 2)^2$ on $[0, 2]$

Since, given function f is a polynomial it is continuous and differentiable everywhere that is on \mathbb{R} .

Let us find the values at extremes:

$$\Rightarrow f(0) = 0(0 - 2)^2$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(2) = 2(2 - 2)^2$$

$$\Rightarrow f(2) = 2(0)^2$$

$$\Rightarrow f(2) = 0$$

$f(0) = f(2)$, Rolle's theorem applicable for function f on $[0, 2]$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(x(x-2)^2)}{dx}$$

Differentiating using UV rule,

$$\Rightarrow f'(x) = (x-2)^2 \times \frac{d(x)}{dx} + x \frac{d((x-2)^2)}{dx}$$

$$\Rightarrow f'(x) = ((x-2)^2 \times 1) + (x \times 2 \times (x-2))$$

$$\Rightarrow f'(x) = (x-2)^2 + 2(x^2 - 2x)$$

$$\Rightarrow f'(x) = x^2 - 4x + 4 + 2x^2 - 4x$$

$$\Rightarrow f'(x) = 3x^2 - 8x + 4$$

We have $f'(c) = 0$ $c \in (0, 1)$, from the definition of Rolle's Theorem.

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 3c^2 - 8c + 4 = 0$$

$$\Rightarrow c = \frac{8 \pm \sqrt{(-8)^2 - (4 \times 3 \times 4)}}{2 \times 3}$$

$$\Rightarrow 3c^2 - 8c + 4 = 0$$

$$\Rightarrow c = \frac{8 \pm \sqrt{(-8)^2 - (4 \times 3 \times 4)}}{2 \times 3}$$

$$\Rightarrow c = \frac{8 \pm \sqrt{64 - 48}}{6}$$

$$\Rightarrow c = \frac{8 \pm \sqrt{16}}{6}$$

$$\Rightarrow c = \frac{12}{6} \text{ or } c = \frac{6}{6}$$

$$\Rightarrow c = 1 \in (0, 2)$$

\therefore Rolle's Theorem is verified.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$c = 12/6 \text{ or } 4/6$$

$$c = 2 \text{ or } 2/3$$

So,

$$c = 2/3 \text{ since } c \in (0, 2)$$

\therefore Rolle's Theorem is verified.

(viii) $f(x) = x^2 + 5x + 6$ on $[-3, -2]$

Solution:

Given function is $f(x) = x^2 + 5x + 6$ on $[-3, -2]$

Since, given function f is a polynomial it is continuous and differentiable everywhere i.e., on \mathbb{R} .
Let us find the values at extremes:

$$\Rightarrow f(-3) = (-3)^2 + 5(-3) + 6$$

$$\Rightarrow f(-3) = 9 - 15 + 6$$

$$\Rightarrow f(-3) = 0$$

$$\Rightarrow f(-2) = (-2)^2 + 5(-2) + 6$$

$$\Rightarrow f(-2) = 4 - 10 + 6$$

$$\Rightarrow f(-2) = 0$$

$\therefore f(-3) = f(-2)$, Rolle's theorem applicable for function f on $[-3, -2]$.

Let's find the derivative of $f(x)$:

$$\Rightarrow f'(x) = \frac{d(x^2 + 5x + 6)}{dx}$$

$$\Rightarrow f'(x) = \frac{d(x^2)}{dx} + \frac{d(5x)}{dx} + \frac{d(6)}{dx}$$

$$\Rightarrow f'(x) = 2x + 5 + 0$$

$$\Rightarrow f'(x) = 2x + 5$$

We have $f'(c) = 0$ $c \in (-3, -2)$, from the definition of Rolle's Theorem

$$\Rightarrow f'(c) = 0$$

We have $f'(c) = 0$ $c \in (-3, -2)$, from the definition of Rolle's Theorem

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 2c + 5 = 0$$

$$\Rightarrow 2c = -5$$

$$\Rightarrow c = -\frac{5}{2}$$

$$\Rightarrow c = -2.5 \in (-3, -2)$$

\therefore Rolle's Theorem is verified.

3. Verify the Rolle's Theorem for each of the following functions on the indicated intervals:

(i) $f(x) = \cos 2(x - \pi/4)$ on $[0, \pi/2]$

Solution:

Given function is $f(x) = \cos 2\left(x - \frac{\pi}{4}\right)$ on $\left[0, \frac{\pi}{2}\right]$

We know that cosine function is continuous and differentiable on \mathbb{R} .

Let's find the values of the function at an extreme,

$$\Rightarrow f(0) = \cos 2\left(0 - \frac{\pi}{4}\right)$$

$$\Rightarrow f(0) = \cos 2\left(-\frac{\pi}{4}\right)$$

$$\Rightarrow f(0) = \cos\left(-\frac{\pi}{2}\right)$$

We know that $\cos(-x) = \cos x$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \cos 2\left(\frac{\pi}{2} - \frac{\pi}{4}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \cos 2\left(\frac{\pi}{4}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \cos\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 0$$

We get $f(0) = f\left(\frac{\pi}{2}\right)$, so there exist a $c \in \left(0, \frac{\pi}{2}\right)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d\left(\cos 2\left(x - \frac{\pi}{4}\right)\right)}{dx}$$

$$\Rightarrow f'(x) = -\sin\left(2\left(x - \frac{\pi}{4}\right)\right) \frac{d\left(2\left(x - \frac{\pi}{4}\right)\right)}{dx}$$

$$\Rightarrow f'(x) = -2 \sin 2\left(x - \frac{\pi}{4}\right)$$

We have $f'(c) = 0$,

$$\Rightarrow -2 \sin 2\left(c - \frac{\pi}{4}\right) = 0$$

$$\Rightarrow c - \frac{\pi}{4} = 0$$

$$\Rightarrow c = \frac{\pi}{4} \in \left(0, \frac{\pi}{2}\right)$$

\therefore Rolle's Theorem is verified.

(ii) $f(x) = \sin 2x$ on $[0, \pi/2]$

Solution:

Given function is $f(x) = \sin 2x$ on $\left[0, \frac{\pi}{2}\right]$

We know that sine function is continuous and differentiable on \mathbb{R} . Let's find the values of function at extreme,

$$\Rightarrow f(0) = \sin 2(0)$$

$$\Rightarrow f(0) = \sin 0$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \sin 2\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f(0) = \sin 0$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \sin 2\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \sin(\pi)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 0$$

We have $f(0) = f\left(\frac{\pi}{2}\right)$, so there exist a $c \in \left(0, \frac{\pi}{2}\right)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(\sin 2x)}{dx}$$

$$\Rightarrow f'(x) = \cos 2x \frac{d(2x)}{dx}$$

$$\Rightarrow f'(x) = 2\cos 2x$$

We have $f'(c) = 0$,

$$\Rightarrow 2 \cos 2c = 0$$

$$\Rightarrow 2c = \frac{\pi}{2}$$

$$\Rightarrow c = \frac{\pi}{4} \in \left(0, \frac{\pi}{2}\right)$$

\therefore Rolle's Theorem is verified.

(iii) $f(x) = \cos 2x$ on $[-\pi/4, \pi/4]$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is $\cos 2x$ on $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$

We know that cosine function is continuous and differentiable on \mathbb{R} . Let's find the values of the function at an extreme,

$$\Rightarrow f\left(-\frac{\pi}{4}\right) = \cos 2\left(-\frac{\pi}{4}\right)$$

$$\Rightarrow f(0) = \cos\left(-\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(-\frac{\pi}{4}\right) = \cos 2\left(-\frac{\pi}{4}\right)$$

$$\Rightarrow f(0) = \cos\left(-\frac{\pi}{2}\right)$$

We know that $\cos(-x) = \cos x$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f\left(\frac{\pi}{4}\right) = \cos 2\left(\frac{\pi}{4}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \cos\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 0$$

We have $f\left(-\frac{\pi}{4}\right) = f\left(\frac{\pi}{4}\right)$, so there exist a $c \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(\cos 2x)}{dx}$$

$$\Rightarrow f'(x) = -\sin 2x \frac{d(2x)}{dx}$$

$$\Rightarrow f'(x) = -2\sin 2x$$

We have $f'(c) = 0$,

$$\Rightarrow -2\sin 2c = 0$$

$$\sin 2c = 0$$

$$\Rightarrow 2c = 0$$

So,

$$c = 0 \text{ as } c \in \left(-\frac{\pi}{4}, \frac{\pi}{4}\right)$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

∴ Rolle's Theorem is verified.

(iv) $f(x) = e^x \sin x$ on $[0, \pi]$

Solution:

Given function is $f(x) = e^x \sin x$ on $[0, \pi]$

We know that exponential and sine functions are continuous and differentiable on \mathbb{R} . Let's find the values of the function at an extreme,

$$\Rightarrow f(0) = e^0 \sin(0)$$

$$\Rightarrow f(0) = 1 \times 0$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(\pi) = e^\pi \sin(\pi)$$

$$\Rightarrow f(\pi) = e^\pi \times 0$$

$$\Rightarrow f(\pi) = 0$$

We have $f(0) = f(\pi)$, so there exist a $c \in (0, \pi)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(e^x \sin x)}{dx}$$

$$\Rightarrow f'(x) = \sin x \frac{d(e^x)}{dx} + e^x \frac{d(\sin x)}{dx}$$

$$\Rightarrow f'(x) = e^x (\sin x + \cos x)$$

We have $f'(c) = 0$,

$$\Rightarrow e^c (\sin c + \cos c) = 0$$

$$\Rightarrow \sin c + \cos c = 0$$

$$\Rightarrow \frac{1}{\sqrt{2}} \sin c + \frac{1}{\sqrt{2}} \cos c = 0$$

$$\Rightarrow \sin\left(\frac{\pi}{4}\right) \sin c + \cos\left(\frac{\pi}{4}\right) \cos c = 0$$

$$\Rightarrow \cos\left(c - \frac{\pi}{4}\right) = 0$$

$$\Rightarrow c - \frac{\pi}{4} = \frac{\pi}{2}$$

$$\Rightarrow c = \frac{3\pi}{4} \in (0, \pi)$$

\therefore Rolle's Theorem is verified.

(v) $f(x) = e^x \cos x$ on $[-\pi/2, \pi/2]$

Solution:

Given function is $f(x) = e^x \cos x$ on $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$

We know that exponential and cosine functions are continuous and differentiable on \mathbb{R} . Let's find the values of the function at an extreme,

$$\Rightarrow f\left(-\frac{\pi}{2}\right) = e^{-\frac{\pi}{2}} \cos\left(-\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(-\frac{\pi}{2}\right) = e^{-\frac{\pi}{2}} \times 0$$

$$\Rightarrow f\left(-\frac{\pi}{2}\right) = 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = e^{\frac{\pi}{2}} \cos\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = e^{\frac{\pi}{2}} \times 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 0$$

We have $f\left(-\frac{\pi}{2}\right) = f\left(\frac{\pi}{2}\right)$, so there exist a $c \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(e^x \cos x)}{dx}$$

$$\Rightarrow f'(x) = \cos x \frac{d(e^x)}{dx} + e^x \frac{d(\cos x)}{dx}$$

$$\Rightarrow f'(x) = e^x (-\sin x + \cos x)$$

We have $f'(c) = 0$,

$$\Rightarrow e^c (-\sin c + \cos c) = 0 \qquad \Rightarrow -\sin\left(\frac{\pi}{4}\right)\sin c + \cos\left(\frac{\pi}{4}\right)\cos c = 0$$

$$\Rightarrow -\sin c + \cos c = 0 \qquad \Rightarrow \cos\left(c + \frac{\pi}{4}\right) = 0$$

$$\Rightarrow \frac{-1}{\sqrt{2}}\sin c + \frac{1}{\sqrt{2}}\cos c = 0 \qquad \Rightarrow c + \frac{\pi}{4} = \frac{\pi}{2}$$

$$\Rightarrow -\sin\left(\frac{\pi}{4}\right)\sin c + \cos\left(\frac{\pi}{4}\right)\cos c = 0 \qquad \Rightarrow c = \frac{\pi}{4} \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$

$$\Rightarrow \cos\left(c + \frac{\pi}{4}\right) = 0 \qquad \therefore \text{Rolle's Theorem is verified.}$$

(vi) $f(x) = \cos 2x$ on $[0, \pi]$

Solution:

Given function is $f(x) = \cos 2x$ on $[0, \pi]$

We know that cosine function is continuous and differentiable on \mathbb{R} . Let's find the values of function at extreme,

$$\Rightarrow f(0) = \cos 2(0)$$

$$\Rightarrow f(0) = \cos(0)$$

$$\Rightarrow f(0) = 1$$

$$\Rightarrow f(\pi) = \cos 2(\pi)$$

$$\pi)$$

$$\Rightarrow f(\pi) = \cos(2\pi)$$

$$\Rightarrow f(\pi) = 1$$

We have $f(0) = f(\pi)$, so there exist a c belongs to $(0, \pi)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow f'(x) = \frac{d(\cos 2x)}{dx}$$

$$\Rightarrow f'(x) = -\sin 2x \frac{d(2x)}{dx}$$

$$\Rightarrow f'(x) = -2\sin 2x$$

We have $f'(c) = 0$,

$$\Rightarrow -2\sin 2c = 0$$

$$\Rightarrow 2c = 0$$

$$\Rightarrow c = \frac{\pi}{4} \in (0, \pi)$$

\therefore Rolle's Theorem is verified.

$$\sin 2c = 0$$

So, $2c = 0$ or π

$$c = 0 \text{ or } \pi/2$$

But,

$$c = \pi/2 \text{ as } c \in (0, \pi)$$

Hence, Rolle's Theorem is verified.

$$\text{(vii) } f(x) = \frac{\sin x}{e^x} \text{ on } 0 \leq x \leq \pi$$

Solution:

Given function is $f(x) = \frac{\sin x}{e^x}$ on $[0, \pi]$

This can be written as

$$\Rightarrow f(x) = e^{-x} \sin x \text{ on } [0, \pi]$$

We know that exponential and sine functions are continuous and differentiable on \mathbb{R} . Let's find the values of the function at an extreme,

$$\Rightarrow f(0) = e^{-0} \sin(0)$$

$$\Rightarrow f(0) = 1 \times 0$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(\pi) = e^{-\pi} \sin(\pi)$$

$$\Rightarrow f(\pi) = e^{-\pi} \times 0$$

$$\Rightarrow f(\pi) = 0$$

We have $f(0) = f(\pi)$, so there exist a c belongs to $(0, \pi)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(e^{-x} \sin x)}{dx}$$

$$\Rightarrow f'(x) = \sin x \frac{d(e^{-x})}{dx} + e^{-x} \frac{d(\sin x)}{dx}$$

$$\Rightarrow f'(x) = \sin x (-e^{-x}) + e^{-x} (\cos x)$$

$$\Rightarrow f'(x) = e^{-x} (-\sin x + \cos x)$$

We have $f'(c) = 0$,

$$\Rightarrow e^{-c} (-\sin c + \cos c) = 0$$

$$\Rightarrow -\sin c + \cos c = 0$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

(viii) $f(x) = \sin 3x$ on $[0, \pi]$

Solution:

Given function is $f(x) = \sin 3x$ on $[0, \pi]$

We know that sine function is continuous and differentiable on \mathbb{R} . Let's find the values of function at extreme,

$$\Rightarrow f(0) = \sin 3(0)$$

$$\Rightarrow f(0) = \sin 0$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(\pi) = \sin 3(\pi)$$

$$\Rightarrow f(\pi) = \sin(3\pi)$$

$$\Rightarrow f(\pi) = 0$$

We have $f(0) = f(\pi)$, so there exist a c belongs to $(0, \pi)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$

$$\Rightarrow f'(x) = \frac{d(\sin 3x)}{dx}$$

$$\Rightarrow f'(x) = \cos 3x \frac{d(3x)}{dx}$$

$$\Rightarrow f'(x) = 3\cos 3x$$

$$\text{We have } f'(c) = 0,$$

$$\Rightarrow 3\cos 3c = 0$$

$$\Rightarrow 3c = \frac{\pi}{2}$$

$$\Rightarrow c = \frac{\pi}{6} \in (0, \pi)$$

\therefore Rolle's Theorem is verified. \therefore Rolle's Theorem is verified.

(ix) $f(x) = e^{1-x^2}$ on $[-1, 1]$

Solution:

Given function is $f(x) = e^{1-x^2}$ on $[-1, 1]$

We know that exponential function is continuous and differentiable over \mathbb{R} .
Let's find the value of function f at extremes,

$$\Rightarrow f(-1) = e^{1-(-1)^2}$$

$$\Rightarrow f(-1) = e^{1-1}$$

$$\Rightarrow f(-1) = e^0$$

$$\Rightarrow f(-1) = 1$$

$$\Rightarrow f(1) = e^{1-1^2}$$

$$\Rightarrow f(1) = e^{1-1}$$

$$\Rightarrow f(1) = e^0$$

$$\Rightarrow f(1) = 1$$

We got $f(-1) = f(1)$ so, there exists a $c \in (-1, 1)$ such that $f'(c) = 0$.

Let's find the derivative of the function f :

$$\Rightarrow f'(x) = \frac{d(e^{1-x^2})}{dx}$$

$$\Rightarrow f'(x) = e^{1-x^2} \frac{d(1-x^2)}{dx}$$

$$\Rightarrow f'(x) = e^{1-x^2} (-2x)$$

$$\Rightarrow f'(x) = e^{1-x^2} \frac{d(1-x^2)}{dx}$$

$$\Rightarrow f'(x) = e^{1-x^2} (-2x)$$

We have $f'(c) = 0$

$$\Rightarrow e^{1-c^2} (-2c) = 0$$

$$\Rightarrow 2c = 0$$

$$\Rightarrow c = 0 \in [-1, 1]$$

\therefore Rolle's Theorem is verified.

(x) $f(x) = \log(x^2 + 2) - \log 3$ on $[-1, 1]$

Solution:

Given function is $f(x) = \log(x^2 + 2) - \log 3$ on $[-1, 1]$

We know that logarithmic function is continuous and differentiable in its own domain. We check the values of the function at the extreme,

$$\Rightarrow f(-1) = \log((-1)^2 + 2) - \log 3$$

$$\Rightarrow f(-1) = \log(1 + 2) - \log 3$$

$$\Rightarrow f(-1) = \log 3 - \log 3$$

$$\Rightarrow f(-1) = 0$$

$$\Rightarrow f(1) = \log(1^2 + 2) - \log 3$$

$$\Rightarrow f(1) = \log(1 + 2) - \log 3$$

$$\Rightarrow f(1) = \log 3 - \log 3$$

$$\Rightarrow f(1) = 0$$

We have got $f(-1) = f(1)$. So, there exists a c such that $c \in (-1, 1)$ such that $f'(c) = 0$.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Let's find the derivative of the function f,

We have $f'(c) = 0$

$$\Rightarrow f'(x) = \frac{d(\log(x^2 + 2) - \log 3)}{dx} \Rightarrow \frac{2x}{x^2 + 2} = 0$$

$$\Rightarrow f'(x) = \frac{1}{x^2 + 2} \frac{d(x^2 + 2)}{dx} - 0 \Rightarrow 2x = 0$$

$$\Rightarrow x = 0 \in (-1, 1)$$

$$\Rightarrow f'(x) = \frac{2x}{x^2 + 2}$$

\therefore Rolle's Theorem is verified.

(xi) $f(x) = \sin x + \cos x$ on $[0, \pi/2]$

Solution:

Given function is $f(x) = \sin x + \cos x$ on $\left[0, \frac{\pi}{2}\right]$

We know that sine and cosine functions are continuous and differentiable on \mathbb{R} . Let's the value of function f at extremes:

$$\Rightarrow f(0) = \sin(0) + \cos(0)$$

$$\Rightarrow f(0) = 0 + 1$$

$$\Rightarrow f(0) = 1$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \sin\left(\frac{\pi}{2}\right) + \cos\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 1 + 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 1$$

We have $f(0) = f\left(\frac{\pi}{2}\right)$. So, there exists a $c \in \left(0, \frac{\pi}{2}\right)$ such that $f'(c) = 0$.

Let's find the derivative of the function f .

$$\Rightarrow f'(x) = \frac{d(\sin x + \cos x)}{dx}$$

$$\Rightarrow f'(x) = \cos x - \sin x$$

We have $f'(c) = 0$

$$\Rightarrow \cos c - \sin c = 0$$

We have $f'(c) = 0$

$$\Rightarrow \cos c - \sin c = 0$$

$$\Rightarrow \frac{1}{\sqrt{2}} \cos c - \frac{1}{\sqrt{2}} \sin c = 0$$

$$\Rightarrow \sin\left(\frac{\pi}{4}\right) \cos c - \cos\left(\frac{\pi}{4}\right) \sin c = 0$$

$$\Rightarrow \sin\left(\frac{\pi}{4} - c\right) = 0$$

$$\Rightarrow \frac{\pi}{4} - c = 0$$

$$\Rightarrow c = \frac{\pi}{4} \in \left(0, \frac{\pi}{2}\right)$$

\therefore Rolle's Theorem is verified.

(xii) $f(x) = 2 \sin x + \sin 2x$ on $[0, \pi]$

Solution:

Given function is $f(x) = 2\sin x + \sin 2x$ on $[0, \pi]$

We know that sine function continuous and differentiable over \mathbb{R} .

Let's check the values of function f at the extremes

$$\Rightarrow f(0) = 2\sin(0) + \sin 2(0)$$

$$\Rightarrow f(0) = 2(0) + 0$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(\pi) = 2\sin(\pi) + \sin 2(\pi)$$

$$\Rightarrow f(\pi) = 2(0) + 0$$

$$\Rightarrow f(\pi) = 0$$

We have $f(0) = f(\pi)$, so there exist a c belongs to $(0, \pi)$ such that $f'(c) = 0$.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Let's find the derivative of function f.

$$\Rightarrow f'(x) = \frac{d(2\sin x + \sin 2x)}{dx}$$

$$\Rightarrow f'(x) = 2\cos x + \cos 2x \frac{d(2x)}{dx}$$

$$\Rightarrow f'(x) = 2\cos x + 2\cos 2x$$

$$\Rightarrow f'(x) = 2\cos x + 2(2\cos^2 x - 1)$$

$$\Rightarrow f'(x) = 4\cos^2 x + 2\cos x - 2$$

We have $f'(c) = 0$,

$$\Rightarrow 4\cos^2 c + 2\cos c - 2 = 0$$

$$\Rightarrow 2\cos^2 c + \cos c - 1 = 0$$

$$\Rightarrow 2\cos^2 c + 2\cos c - \cos c - 1 = 0$$

$$\Rightarrow 2\cos c (\cos c + 1) - 1(\cos c + 1) = 0$$

$$\Rightarrow (2\cos c - 1)(\cos c + 1) = 0$$

$$\Rightarrow \cos c = \frac{1}{2} \text{ or } \cos c = -1$$

$$\Rightarrow c = \frac{\pi}{3} \in (0, \pi)$$

\therefore Rolle's Theorem is verified. (xiii) $f(x) = \frac{x}{2} - \sin \frac{\pi x}{6}$ on $[-1, 0]$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is $f(x) = \frac{x}{2} - \sin\left(\frac{\pi x}{6}\right)$ on $[-1, 0]$

We know that sine function is continuous and differentiable over \mathbb{R} .

Now we have to check the values of 'f' at an extreme

$$\Rightarrow f(-1) = \frac{-1}{2} - \sin\left(\frac{\pi(-1)}{6}\right)$$

$$\Rightarrow f(-1) = -\frac{1}{2} - \sin\left(\frac{-\pi}{6}\right)$$

$$\Rightarrow f(-1) = -\frac{1}{2} - \left(-\frac{1}{2}\right)$$

$$\Rightarrow f(-1) = 0$$

$$\Rightarrow f(0) = \frac{0}{2} - \sin\left(\frac{\pi(0)}{6}\right)$$

$$\Rightarrow f(0) = 0 - \sin(0)$$

$$\Rightarrow f(0) = 0 - 0$$

$$\Rightarrow f(0) = 0$$

We have got $f(-1) = f(0)$. So, there exists a $c \in (-1, 0)$ such that $f'(c) = 0$.

Now we have to find the derivative of the function 'f'

$$\Rightarrow f'(x) = \frac{d\left(\frac{x}{2} - \sin\left(\frac{\pi x}{6}\right)\right)}{dx}$$

$$\Rightarrow f(0) = 0$$

We have got $f(-1) = f(0)$. So, there exists a $c \in (-1, 0)$ such that $f'(c) = 0$.

Now we have to find the derivative of the function 'f'

$$\Rightarrow f'(x) = \frac{d\left(\frac{x}{2} - \sin\left(\frac{\pi x}{6}\right)\right)}{dx}$$

$$\Rightarrow f'(x) = \frac{1}{2} - \cos\left(\frac{\pi x}{6}\right) \frac{d\left(\frac{\pi x}{6}\right)}{dx}$$

$$\Rightarrow f'(x) = \frac{1}{2} - \frac{\pi}{6} \cos\left(\frac{\pi x}{6}\right)$$

We have $f'(c) = 0$

$$\Rightarrow \frac{1}{2} - \frac{\pi}{6} \cos\left(\frac{\pi c}{6}\right) = 0$$

$$\Rightarrow \frac{\pi}{6} \cos\left(\frac{\pi c}{6}\right) = \frac{1}{2}$$

$$\Rightarrow \cos\left(\frac{\pi c}{6}\right) = \frac{1}{2} \times \frac{6}{\pi}$$

$$\Rightarrow \cos\left(\frac{\pi c}{6}\right) = \frac{3}{\pi}$$

$$\Rightarrow \frac{\pi c}{6} = \cos^{-1}\left(\frac{3}{\pi}\right)$$

$$\Rightarrow c = \frac{6}{\pi} \cos^{-1}\left(\frac{3}{\pi}\right)$$

Cosine is positive between $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$, for our convenience we take the interval to be $-\frac{\pi}{2} \leq \theta \leq 0$, since the values of the cosine repeats.

We know that $\frac{3}{\pi}$ value is nearly equal to 1. So, the value of the c nearly equal to 0.

So, we can clearly say that $c \in (-1, 0)$.

\therefore Rolle's Theorem is verified.

(xiv). $f(x) = \frac{6x}{\pi} - 4 \sin^2 x$ on $\left[0, \frac{\pi}{6}\right]$

Solution:

$$\Rightarrow f'(x) = \frac{6}{\pi} - 4(2\sin x \cos x)$$

$$\Rightarrow f'(x) = \frac{6}{\pi} - 4\sin 2x$$

We have $f'(c) = 0$

$$\Rightarrow \frac{6}{\pi} - 4\sin 2c = 0$$

$$\Rightarrow 4\sin 2c = \frac{6}{\pi}$$

$$\Rightarrow \sin 2c = \frac{6}{4\pi}$$

We know $\frac{6}{4\pi} < \frac{1}{2}$

$$\Rightarrow \sin 2c < \frac{1}{2}$$

$$\Rightarrow 2c < \sin^{-1}\left(\frac{1}{2}\right)$$

$$\Rightarrow 2c < \frac{\pi}{6}$$

$$\Rightarrow f'(x) = \frac{6}{\pi} - 4 \times 2\sin x \times \frac{d(\sin x)}{dx}$$

$$\Rightarrow f'(x) = \frac{6}{\pi} - 8\sin x(\cos x)$$

$$\Rightarrow c < \frac{\pi}{12} \in \left(0, \frac{\pi}{6}\right)$$

\therefore Rolle's Theorem is verified.

(xv) $f(x) = 4^{\sin x}$ on $[0, \pi]$

Solution:

Given function is $f(x) = 4^{\sin x}$ on $[0, \pi]$

We that sine function is continuous and differentiable over \mathbb{R} .

Now we have to check the values of function 'f' at extremes

$$\Rightarrow f(0) = 4^{\sin(0)}$$

$$\Rightarrow f(0) = 4^0$$

$$\Rightarrow f(0) = 1$$

$$\Rightarrow f(\pi) = 4^{\sin \pi}$$

$$\Rightarrow f(\pi) = 4^0$$

$$\Rightarrow f(\pi) = 1$$

$$\Rightarrow f(\pi) = 4^{\sin \pi}$$

$$\Rightarrow f(\pi) = 4^0$$

$$\Rightarrow f(\pi) = 1$$

We have $f(0) = f(\pi)$. So, there exists a $c \in (0, \pi)$ such that $f'(c) = 0$.

Now we have to find the derivative of 'f'

$$\Rightarrow f'(x) = \frac{d(4^{\sin x})}{dx}$$

$$\Rightarrow f'(x) = 4^{\sin x} \log 4 \frac{d(\sin x)}{dx}$$

$$\Rightarrow f'(x) = 4^{\sin x} \log 4 \cos x$$

We have $f'(c) = 0$

$$\Rightarrow 4^{\sin c} \log 4 \cos c = 0$$

$$\Rightarrow \cos c = 0$$

$$\Rightarrow c = \frac{\pi}{2} \in (0, \pi)$$

\therefore Rolle's Theorem is verified.

(xvi) $f(x) = x^2 - 5x + 4$ on $[0, \pi/6]$

Solution:

Given function is $f(x) = x^2 - 5x + 4$ on $[1, 4]$

Since, given function f is a polynomial it is continuous and differentiable everywhere i.e., on \mathbb{R} .

Let us find the values at extremes

$$\Rightarrow f(1) = 1^2 - 5(1) + 4$$

$$\Rightarrow f(1) = 1 - 5 + 4$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow f(1) = 0$$

$$\Rightarrow f(4) = 4^2 - 5(4) + 4$$

$$\Rightarrow f(4) = 16 - 20 + 4$$

$$\Rightarrow f(4) = 0$$

We have $f(1) = f(4)$. So, there exists a $c \in (1, 4)$ such that $f'(c) = 0$.

Let's find the derivative of $f(x)$:

$$\Rightarrow f'(x) = \frac{d(x^2 - 5x + 4)}{dx}$$

$$\Rightarrow f'(x) = \frac{d(x^2)}{dx} - \frac{d(5x)}{dx} + \frac{d(4)}{dx}$$

$$\Rightarrow f'(x) = 2x - 5 + 0$$

$$\Rightarrow f'(x) = 2x - 5$$

We have $f'(c) = 0$

$$\Rightarrow f'(c) = 0$$

$$\Rightarrow 2c - 5 = 0$$

$$\Rightarrow 2c = 5$$

$$\Rightarrow c = \frac{5}{2}$$

$$\Rightarrow c = 2.5 \in (1, 4)$$

\therefore Rolle's Theorem is verified.

(xvii) $f(x) = \sin^4 x + \cos^4 x$ on $[0, \pi/2]$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given function is $f(x) = \sin^4 x + \cos^4 x$ on $\left[0, \frac{\pi}{2}\right]$

We know that sine and cosine functions are continuous and differentiable functions over \mathbb{R} .

Now we have to find the value of function 'f' at extremes

$$\Rightarrow f(0) = \sin^4(0) + \cos^4(0)$$

$$\Rightarrow f(0) = (0)^4 + (1)^4$$

$$\Rightarrow f(0) = 0 + 1$$

$$\Rightarrow f(0) = 1$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \sin^4\left(\frac{\pi}{2}\right) + \cos^4\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 1^4 + 0^4$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = \sin^4\left(\frac{\pi}{2}\right) + \cos^4\left(\frac{\pi}{2}\right)$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 1^4 + 0^4$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 1 + 0$$

$$\Rightarrow f\left(\frac{\pi}{2}\right) = 1$$

We have $f(0) = f\left(\frac{\pi}{2}\right)$. So, there exists a $c \in \left(0, \frac{\pi}{2}\right)$ such that $f'(c) = 0$.

Now we have to find the derivative of the function 'f'.

$$\Rightarrow f'(x) = \frac{d(\sin^4 x + \cos^4 x)}{dx}$$

$$\Rightarrow f'(x) = 4 \sin^3 x \frac{d(\sin x)}{dx} + 4 \cos^3 x \frac{d(\cos x)}{dx}$$

$$\Rightarrow f'(x) = 4 \sin^3 x \cos x - 4 \cos^3 x \sin x$$

$$\Rightarrow f'(x) = 4 \sin x \cos x (\sin^2 x - \cos^2 x)$$

$$\Rightarrow f'(x) = 2(2 \sin x \cos x) (-\cos 2x)$$

$$\Rightarrow f'(x) = -2(\sin 2x) (\cos 2x)$$

$$\Rightarrow f'(x) = -\sin 4x$$

We have $f'(c) = 0$

$$\Rightarrow -\sin 4c = 0$$

$$\Rightarrow \sin 4c = 0$$

$$\Rightarrow 4c = 0 \text{ or } \pi$$

$$\Rightarrow c = \frac{\pi}{4} \in \left(0, \frac{\pi}{2}\right)$$

\therefore Rolle's Theorem is verified.

(xviii) $f(x) = \sin x - \sin 2x$ on $[0, \pi]$

Solution:

Given function is $f(x) = \sin x - \sin 2x$ on $[0, \pi]$

We know that sine function is continuous and differentiable over \mathbb{R} .

Now we have to check the values of the function 'f' at the extremes.

$$\Rightarrow f(0) = \sin(0) - \sin 2(0)$$

$$\Rightarrow f(0) = 0 - \sin(0)$$

$$\Rightarrow f(0) = 0$$

$$\Rightarrow f(\pi) = \sin(\pi) - \sin 2(\pi)$$

$$\Rightarrow f(\pi) = 0 - \sin(2\pi)$$

$$\Rightarrow f(\pi) = 0$$

We have $f(0) = f(\pi)$. So, there exists a $c \in (0, \pi)$ such that $f'(c) = 0$.

Now we have to find the derivative of the function 'f'

$$\Rightarrow f'(x) = \frac{d(\sin x - \sin 2x)}{dx}$$

$$\Rightarrow f'(x) = \cos x - \cos 2x \frac{d(2x)}{dx}$$

$$\Rightarrow f'(x) = \cos x - 2\cos 2x$$

$$\Rightarrow f'(x) = \cos x - 2(2\cos^2 x - 1)$$

$$\Rightarrow f'(x) = \cos x - 4\cos^2 x + 2$$

We have $f'(c) = 0$

$$\Rightarrow \cos c - 4\cos^2 c + 2 = 0$$

$$\Rightarrow \cos c = \frac{-1 \pm \sqrt{(1)^2 - (4 \times -4 \times 2)}}{2 \times -4}$$

$$\Rightarrow \cos c = \frac{-1 \pm \sqrt{1 + 33}}{-8}$$

$$\Rightarrow c = \cos^{-1}\left(\frac{-1 \pm \sqrt{33}}{-8}\right)$$

We can see that $c \in (0, \pi)$

\therefore Rolle's Theorem is verified.

4. Using Rolle's Theorem, find points on the curve $y = 16 - x^2$, $x \in [-1, 1]$, where tangent is parallel to x - axis.

Solution:

Given function is $y = 16 - x^2$, $x \in [-1, 1]$

We know that polynomial function is continuous and differentiable over \mathbb{R} .

Let us check the values of 'y' at extremes

$$\Rightarrow y(-1) = 16 - (-1)^2$$

$$\Rightarrow y(-1) = 16 - 1$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow y(-1) = 15$$

$$\Rightarrow y(1) = 16 - (1)^2$$

$$\Rightarrow y(1) = 16 - 1$$

$$\Rightarrow y(1) = 15$$

We have $y(-1) = y(1)$. So, there exists a $c \in (-1, 1)$ such that $f'(c) = 0$.

We know that for a curve g , the value of the slope of the tangent at a point r is given by $g'(r)$.

Now we have to find the derivative of curve y

\Rightarrow

$$y' = \frac{d(16-x^2)}{dx}$$

$$\Rightarrow y' = -2x$$

We have $y'(c) = 0$

$$\Rightarrow -2c = 0$$

$$\Rightarrow c = 0 \in (-1, 1)$$

Value of y at $x = 1$ is

$$\Rightarrow y = 16 - 0^2$$

$$\Rightarrow y = 16$$

\therefore The point at which the curve y has a tangent parallel to x - axis (since the slope of x - axis is 0) is $(0, 16)$.

Exercise 15.2 Page No: 15.17

1. Verify Lagrange's mean value theorem for the following functions on the indicated intervals. In each case find a point 'c' in the indicated interval as stated by the Lagrange's mean value theorem:

(i) $f(x) = x^2 - 1$ on $[2, 3]$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given $f(x) = x^2 - 1$ on $[2, 3]$

We know that every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[2, 3]$ and differentiable in $(2, 3)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (2, 3)$ such that:

$$f'(c) = \frac{f(3) - f(2)}{3 - 2}$$
$$\Rightarrow f'(c) = \frac{f(3) - f(2)}{1}$$

$$f(x) = x^2 - 1$$

Differentiating with respect to x

$$f'(x) = 2x$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = 2c$$

For $f(3)$, put the value of $x=3$ in $f(x)$:

$$f(3) = (3)^2 - 1$$

$$= 9 - 1$$

$$= 8$$

For $f(2)$, put the value of $x=2$ in $f(x)$:

$$f(2) = (2)^2 - 1$$

$$= 4 - 1$$

$$= 3$$

$$\therefore f'(c) = f(3) - f(2)$$

$$\Rightarrow 2c = 8 - 3$$

$$\Rightarrow 2c = 5$$

$$\Rightarrow c = \frac{5}{2} \in (2, 3)$$

Hence, Lagrange's mean value theorem is verified.

(ii) $f(x) = x^3 - 2x^2 - x + 3$ on $[0, 1]$

Solution:

Given $f(x) = x^3 - 2x^2 - x + 3$ on $[0, 1]$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[0, 1]$ and differentiable in $(0, 1)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (0, 1)$ such that:

$$f'(c) = \frac{f(1) - f(0)}{1 - 0}$$

$$\Rightarrow f'(c) = \frac{f(1) - f(0)}{1}$$

$$f(x) = x^3 - 2x^2 - x + 3$$

Differentiating with respect to x

$$f'(x) = 3x^2 - 2(2x) - 1$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$= 3x^2 - 4x - 1$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$

$$f'(c) = 3c^2 - 4c - 1$$

For $f(1)$, put the value of $x = 1$ in $f(x)$

$$f(1) = (1)^3 - 2(1)^2 - (1) + 3$$

$$= 1 - 2 - 1 + 3$$

$$= 1$$

For $f(0)$, put the value of $x=0$ in $f(x)$

$$f(0) = (0)^3 - 2(0)^2 - (0) + 3$$

$$= 0 - 0 - 0 + 3$$

$$= 3$$

$$\therefore f'(c) = f(1) - f(0)$$

$$\Rightarrow 3c^2 - 4c - 1 = 1 - 3$$

$$\Rightarrow 3c^2 - 4c = 1 + 1 - 3$$

$$\Rightarrow 3c^2 - 4c = -1$$

$$\Rightarrow 3c^2 - 4c + 1 = 0$$

$$\Rightarrow 3c^2 - 3c - c + 1 = 0$$

$$\Rightarrow 3c(c - 1) - 1(c - 1) = 0$$

$$\Rightarrow (3c - 1)(c - 1) = 0$$

$$\Rightarrow c = \frac{1}{3}, 1$$

$$\Rightarrow c = \frac{1}{3} \in (0, 1)$$

Hence, Lagrange's mean value theorem is verified.

(iii) $f(x) = x(x - 1)$ on $[1, 2]$

Solution:

Given $f(x) = x(x - 1)$ on $[1, 2]$

$$= x^2 - x$$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[1, 2]$ and differentiable in $(1, 2)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (1, 2)$ such that:

$$f'(c) = \frac{f(2) - f(1)}{2 - 1}$$
$$\Rightarrow f'(c) = \frac{f(2) - f(1)}{1}$$

$$f(x) = x^2 - x$$

Differentiating with respect to x

$$f'(x) = 2x - 1$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = 2c - 1$$

For $f(2)$, put the value of $x = 2$ in $f(x)$

$$f(2) = (2)^2 - 2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$= 4 - 2$$

$$= 2$$

For $f(1)$, put the value of $x = 1$ in $f(x)$:

$$f(1) = (1)^2 - 1$$

$$= 1 - 1$$

$$= 0$$

$$\therefore f'(c) = f(2) - f(1)$$

$$\Rightarrow 2c - 1 = 2 - 0$$

$$\Rightarrow 2c = 2 + 1$$

$$\Rightarrow 2c = 3$$

$$\Rightarrow c = \frac{3}{2} \in (1, 2)$$

Hence, Lagrange's mean value theorem is verified.

(iv) $f(x) = x^2 - 3x + 2$ on $[-1, 2]$

Solution:

Given $f(x) = x^2 - 3x + 2$ on $[-1, 2]$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[-1, 2]$ and differentiable in $(-1, 2)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (-1, 2)$ such that:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$f'(c) = \frac{f(2) - f(-1)}{2 - (-1)}$$

$$\Rightarrow f'(c) = \frac{f(2) - f(-1)}{2 + 1}$$

$$\Rightarrow f'(c) = \frac{f(2) - f(-1)}{3}$$

$$f(x) = x^2 - 3x + 2$$

Differentiating with respect to x

$$f'(x) = 2x - 3$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = 2c - 3$$

For $f(2)$, put the value of $x = 2$ in $f(x)$

$$f(2) = (2)^2 - 3(2) + 2$$

$$= 4 - 6 + 2$$

$$= 0$$

For $f(-1)$, put the value of $x = -1$ in $f(x)$:

$$f(-1) = (-1)^2 - 3(-1) + 2$$

$$= 1 + 3 + 2$$

$$= 6$$

$$f'(c) = \frac{f(2) - f(-1)}{3}$$

$$\Rightarrow 2c - 3 = \frac{0 - 6}{3}$$

$$\Rightarrow 2c = \frac{-6}{3} + 3$$

$$\Rightarrow 2c = -2 + 3$$

$$\Rightarrow 2c = 1$$

$$\Rightarrow c = \frac{1}{2} \in (-1, 2)$$

Hence, Lagrange's mean value theorem is verified.

$$\Rightarrow 2c = 1$$

$$\Rightarrow c = \frac{1}{2} \in (-1, 2)$$

Hence, Lagrange's mean value theorem is verified.

(v) $f(x) = 2x^2 - 3x + 1$ on $[1, 3]$

Solution:

Given $f(x) = 2x^2 - 3x + 1$ on $[1, 3]$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[1, 3]$ and differentiable in $(1, 3)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (1, 3)$ such that:

$$f'(c) = \frac{f(3) - f(1)}{3 - 1}$$

$$\Rightarrow f'(c) = \frac{f(3) - f(1)}{2}$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$f(x) = 2x^2 - 3x + 1$$

Differentiating with respect to x

$$f'(x) = 2(2x) - 3$$

$$= 4x - 3$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = 4c - 3$$

For $f(3)$, put the value of $x = 3$ in $f(x)$:

$$f(3) = 2(3)^2 - 3(3) + 1$$

$$= 2(9) - 9 + 1$$

$$= 18 - 9 = 9$$

For $f(1)$, put the value of $x = 1$ in $f(x)$:

$$f(1) = 2(1)^2 - 3(1) + 1$$

$$= 2(1) - 3 + 1$$

$$= 2 - 2 = 0$$

$$f'(c) = \frac{f(3) - f(1)}{2}$$

$$\Rightarrow 4c - 3 = \frac{10 - 0}{2}$$

$$\Rightarrow 4c = \frac{10}{2} + 3$$

$$\Rightarrow 4c = 5 + 3$$

$$\Rightarrow 4c = 8$$

$$\Rightarrow c = \frac{8}{4} = 2 \in (1, 3)$$

Hence, Lagrange's mean value theorem is verified.

(vi) $f(x) = x^2 - 2x + 4$ on $[1, 5]$

Solution:

Given $f(x) = x^2 - 2x + 4$ on $[1, 5]$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[1, 5]$ and differentiable in $(1, 5)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (1, 5)$ such that:

$$f'(c) = \frac{f(5) - f(1)}{5 - 1}$$

$$\Rightarrow f'(c) = \frac{f(5) - f(1)}{4}$$

$$f(x) = x^2 - 2x + 4$$

Differentiating with respect to x :

$$f'(x) = 2x - 2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = 2c - 2$$

For $f(5)$, put the value of $x=5$ in $f(x)$:

$$f(5) = (5)^2 - 2(5) + 4$$

$$= 25 - 10 + 4$$

$$= 19$$

For $f(1)$, put the value of $x = 1$ in $f(x)$

$$f(1) = (1)^2 - 2(1) + 4$$

$$= 1 - 2 + 4$$

$$= 3$$

$$f'(c) = \frac{f(5) - f(1)}{4}$$

$$\Rightarrow 2c - 2 = \frac{19 - 3}{4}$$

$$\Rightarrow 2c = \frac{16}{4} + 2$$

$$\Rightarrow 2c = 4 + 2$$

$$\Rightarrow 2c = 6$$

$$\Rightarrow c = \frac{6}{2} = 3 \in (1, 5)$$

Hence, Lagrange's mean value theorem is verified.

(vii) $f(x) = 2x - x^2$ on $[0, 1]$

Solution:

Given $f(x) = 2x - x^2$ on $[0, 1]$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[0, 1]$ and differentiable in $(0, 1)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (0, 1)$ such that:

$$f'(c) = \frac{f(1) - f(0)}{1 - 0}$$

$$\Rightarrow f'(c) = f(1) - f(0)$$

$$f(x) = 2x - x^2$$

Differentiating with respect to x :

$$f'(x) = 2 - 2x$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = 2 - 2c$$

For $f(1)$, put the value of $x = 1$ in $f(x)$:

$$f(1) = 2(1) - (1)^2$$

$$= 2 - 1$$

$$= 1$$

For $f(0)$, put the value of $x = 0$ in $f(x)$:

$$f(0) = 2(0) - (0)^2$$

$$= 0 - 0$$

$$= 0$$

$$f'(c) = f(1) - f(0)$$

$$\Rightarrow 2 - 2c = 1 - 0$$

$$\Rightarrow -2c = 1 - 2$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow -2c = -1$$

$$\Rightarrow c = \frac{-1}{-2} = \frac{1}{2} \in (0, 1)$$

Hence, Lagrange's mean value theorem is verified.

(viii) $f(x) = (x - 1)(x - 2)(x - 3)$

Solution:

Given $f(x) = (x - 1)(x - 2)(x - 3)$ on $[0, 4]$

$$= (x^2 - x - 2x + 2)(x - 3)$$

$$= (x^2 - 3x + 2)(x - 3)$$

$$= x^3 - 3x^2 + 2x - 3x^2 + 9x - 6$$

$$= x^3 - 6x^2 + 11x - 6 \text{ on } [0, 4]$$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[0, 4]$ and differentiable in $(0, 4)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (0, 4)$ such that:

$$f'(c) = \frac{f(4) - f(0)}{4 - 0}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(0)}{4}$$

$$f(x) = x^3 - 6x^2 + 11x - 6$$

Differentiating with respect to x :

$$f'(x) = 3x^2 - 6(2x) + 11$$

$$= 3x^2 - 12x + 11$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$f'(c) = 3c^2 - 12c + 11$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f(4) = (4)^3 - 6(4)^2 + 11(4) - 6$$

$$= 64 - 96 + 44 - 6$$

$$= 6$$

For $f(0)$, put the value of $x = 0$ in $f(x)$:

$$f(0) = (0)^3 - 6(0)^2 + 11(0) - 6$$

$$= 0 - 0 + 0 - 6$$

$$= -6$$

$$f'(c) = \frac{f(4) - f(0)}{4}$$

$$\Rightarrow 3c^2 - 12c + 12 = \frac{7 - (-9)}{4}$$

$$\Rightarrow 3c^2 - 12c + 12 = \frac{7 + 9}{4}$$

$$\Rightarrow 3c^2 - 12c + 12 = \frac{16}{4}$$

$$\Rightarrow 3c^2 - 12c + 12 = 4$$

$$\Rightarrow 3c^2 - 12c + 8 = 0$$

We know that for quadratic equation, $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow c = \frac{-(-12) \pm \sqrt{(-12)^2 - 4 \times 3 \times 8}}{2 \times 3}$$

$$\Rightarrow c = \frac{12 \pm \sqrt{144 - 96}}{6}$$

$$\Rightarrow c = \frac{12 \pm \sqrt{48}}{6}$$

$$3c^2 - 12c + 11 = [6 - (-6)]/4$$

$$3c^2 - 12c + 11 = 12/4$$

$$3c^2 - 12c + 11 = 3$$

$$3c^2 - 12c + 8 = 0$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow 3c^2 - 12c + 12 = 4$$

$$\Rightarrow 3c^2 - 12c + 8 = 0$$

We know that for quadratic equation, $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow c = \frac{-(-12) \pm \sqrt{(-12)^2 - 4 \times 3 \times 8}}{2 \times 3}$$

$$\Rightarrow c = \frac{12 \pm \sqrt{144 - 96}}{6}$$

$$\Rightarrow c = \frac{12 \pm \sqrt{48}}{6}$$

$$\Rightarrow c = \frac{12 \pm 4\sqrt{3}}{6}$$

$$\Rightarrow c = \frac{12}{6} \pm \frac{4\sqrt{3}}{6}$$

$$\Rightarrow c = 2 \pm \frac{2\sqrt{3}}{3}$$

$$\Rightarrow c = 2 + \frac{2\sqrt{3}}{3}, 2 - \frac{2\sqrt{3}}{3} \in c$$

Hence, Lagrange's mean value theorem is verified.

(ix). $f(x) = \sqrt{25 - x^2}$ on $[-3, 4]$

Solution:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Given

$$f(x) = \sqrt{25 - x^2} \text{ on } [-3, 4]$$

$$\text{Here, } \sqrt{25 - x^2} > 0$$

$$\Rightarrow 25 - x^2 > 0$$

$$\Rightarrow x^2 < 25$$

$$\Rightarrow -5 < x < 5$$

$$\Rightarrow \sqrt{25 - x^2} \text{ has unique values for all } x \in (-5, 5)$$

$\therefore f(x)$ is continuous in $[-3, 4]$

$$f(x) = (25 - x^2)^{\frac{1}{2}}$$

Differentiating with respect to x :

$$f'(x) = \frac{1}{2} (25 - x^2)^{\left(\frac{1}{2} - 1\right)} \frac{d(25 - x^2)}{dx}$$

$$\Rightarrow f'(x) = \frac{1}{2} (25 - x^2)^{-\frac{1}{2}} (-2x)$$

$$\Rightarrow f'(x) = \frac{-2x}{2(25 - x^2)^{\frac{1}{2}}}$$

$$\Rightarrow 25 - x^2 > 0$$

$$\Rightarrow x^2 < 25$$

$$\Rightarrow -5 < x < 5$$

$\Rightarrow \sqrt{25 - x^2}$ has unique values for all $x \in (-5, 5)$

$\therefore f(x)$ is continuous in $[-3, 4]$

$$f(x) = (25 - x^2)^{\frac{1}{2}}$$

Differentiating with respect to x :

$$f'(x) = \frac{1}{2} (25 - x^2)^{\left(\frac{1}{2} - 1\right)} \frac{d(25 - x^2)}{dx}$$

$$\Rightarrow f'(x) = \frac{1}{2} (25 - x^2)^{-\frac{1}{2}} (-2x)$$

$$\Rightarrow f'(x) = \frac{-2x}{2 (25 - x^2)^{\frac{1}{2}}}$$

$$\Rightarrow f'(x) = \frac{-2x}{2 (25 - x^2)^{\frac{1}{2}}}$$

$$\Rightarrow f'(x) = \frac{-x}{\sqrt{25 - x^2}}$$

Here also,

$$\sqrt{25 - x^2} > 0$$

$$\Rightarrow -5 < x < 5$$

$\therefore f(x)$ is differentiable in $(-3, 4)$

So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (-3, 4)$ such that:

$$f'(c) = \frac{f(4) - f(-3)}{4 - (-3)}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(-3)}{4 + 3}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(-3)}{7}$$

$$f(x) = (25 - x^2)^{\frac{1}{2}}$$

On differentiating with respect to x :

$$f'(x) = \frac{-x}{\sqrt{25 - x^2}}$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = \frac{-c}{\sqrt{25 - c^2}}$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$\Rightarrow f'(c) = \frac{f(4) - f(-3)}{4 + 3}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(-3)}{7}$$

$$f(x) = (25 - x^2)^{\frac{1}{2}}$$

On differentiating with respect to x:

$$f'(x) = \frac{-x}{\sqrt{25 - x^2}}$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = \frac{-c}{\sqrt{25 - c^2}}$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f(4) = (25 - 4^2)^{\frac{1}{2}}$$

$$\Rightarrow f(4) = (25 - 16)^{\frac{1}{2}}$$

$$\Rightarrow f(4) = (9)^{\frac{1}{2}}$$

$$\Rightarrow f(4) = 3$$

For $f(-3)$, put the value of $x = -3$ in $f(x)$:

$$f(-3) = (25 - (-3)^2)^{\frac{1}{2}}$$

$$\Rightarrow f(-3) = (25 - 9)^{\frac{1}{2}}$$

$$\Rightarrow f(-3) = (16)^{\frac{1}{2}}$$

$$\Rightarrow f(-3) = 4$$

$$f'(c) = \frac{f(4) - f(-3)}{7}$$

$$\Rightarrow \frac{-c}{\sqrt{25-c^2}} = \frac{3-4}{7}$$

$$\Rightarrow \frac{-c}{\sqrt{25-c^2}} = \frac{-1}{7}$$

$$\Rightarrow -7c = -\sqrt{25-c^2}$$

Squaring on both sides:

$$\Rightarrow (-7c)^2 = (-\sqrt{25-c^2})^2$$

$$\Rightarrow 49c^2 = 25 - c^2$$

$$\Rightarrow 50c^2 = 25$$

$$\Rightarrow c^2 = \frac{25}{50}$$

$$\Rightarrow c^2 = \frac{1}{2}$$

$$\Rightarrow c = \pm \frac{1}{\sqrt{2}} \in (-3, 4)$$

Hence, Lagrange's mean value theorem is verified.

(x) $f(x) = \tan^{-1}x$ on $[0, 1]$

Solution:

Given $f(x) = \tan^{-1}x$ on $[0, 1]$

$\tan^{-1}x$ has unique value for all x between 0 and 1.

$\therefore f(x)$ is continuous in $[0, 1]$

$f(x) = \tan^{-1}x$

Differentiating with respect to x :

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$f(x) = \frac{1}{1+x^2}$$

x^2 always has value greater than 0.

$$\Rightarrow 1 + x^2 > 0$$

$\therefore f(x)$ is differentiable in $(0, 1)$

So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (0, 1)$ such that:

$$f'(c) = \frac{f(1) - f(0)}{1 - 0}$$

$$\Rightarrow f'(c) = f(1) - f(0)$$

$$f(x) = \tan^{-1} x$$

Differentiating with respect to x:

$$f'(x) = \frac{1}{1+x^2}$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = \frac{1}{1+c^2} \quad \Rightarrow \frac{1}{1+c^2} = \frac{\pi}{4} - 0$$

$$\text{For } f(1), \text{ put the value of } x=1 \text{ in } f(x): \quad \Rightarrow \frac{1}{1+c^2} = \frac{\pi}{4}$$

$$f(1) = \tan^{-1} 1 \quad \Rightarrow 4 = \pi(1+c^2)$$

$$\Rightarrow f(1) = \frac{\pi}{4} \quad \Rightarrow 4 = \pi + \pi c^2$$

$$\text{For } f(0), \text{ put the value of } x=0 \text{ in } f(x): \quad \Rightarrow -\pi c^2 = \pi - 4$$

$$f(0) = \tan^{-1} 0 \quad \Rightarrow c^2 = \frac{\pi - 4}{-\pi}$$

$$\Rightarrow f(0) = 0$$

$$f'(c) = f(1) - f(0) \quad \Rightarrow c^2 = \frac{4 - \pi}{\pi}$$

$$\Rightarrow c^2 = \frac{\pi - 4}{-\pi}$$

$$\Rightarrow c^2 = \frac{4 - \pi}{\pi}$$

$$\Rightarrow c = \sqrt{\frac{4}{\pi} - 1} \approx 0.52 \in (0, 1)$$

Hence, Lagrange's mean value theorem is verified. (xi) $f(x) = x + \frac{1}{x}$ on $[1, 3]$

Solution:

Given

$$f(x) = x + \frac{1}{x} \text{ on } [1, 3]$$

$f(x)$ has unique values for all $x \in (1, 3)$

$\therefore f(x)$ is continuous in $[1, 3]$

$$f(x) = x + \frac{1}{x} \text{ on } [1, 3]$$

Differentiating with respect to x

$$f'(x) = 1 + (-1)(x)^{-2}$$

$$\Rightarrow f'(x) = 1 - \frac{1}{x^2}$$

$$\Rightarrow f'(x) = \frac{x^2 - 1}{x^2}$$

Here, $x^2 \neq 0$

$\Rightarrow f'(x)$ exists for all values except 0

$\therefore f(x)$ is differentiable in $(1, 3)$

So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (1, 3)$ such that:

$$f'(c) = \frac{f(3) - f(1)}{3 - 1}$$

So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (1, 3)$ such that:

$$f'(c) = \frac{f(3) - f(1)}{3 - 1}$$

$$\Rightarrow f'(c) = \frac{f(3) - f(1)}{2}$$

$$f(x) = x + \frac{1}{x}$$

On differentiating with respect to x :

$$f'(x) = \frac{x^2 - 1}{x^2}$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = \frac{c^2 - 1}{c^2}$$

For $f(3)$, put the value of $x = 3$ in $f(x)$:

$$f(3) = 3 + \frac{1}{3}$$

$$\Rightarrow f(3) = \frac{9+1}{3}$$

$$\Rightarrow f(3) = \frac{10}{3}$$

For $f(1)$, put the value of $x = 1$ in $f(x)$:

$$f(1) = 1 + \frac{1}{1}$$

$$\Rightarrow f(1) = 2$$

$$\Rightarrow f'(c) = \frac{f(3) - f(1)}{2}$$

$$\Rightarrow \frac{c^2 - 1}{c^2} = \frac{\frac{10}{3} - 2}{2}$$

$$\Rightarrow 2(c^2 - 1) = c^2 \left(\frac{10}{3} - 2 \right)$$

$$\Rightarrow 2(c^2 - 1) = c^2 \left(\frac{10}{3} - 2 \right)$$

$$\Rightarrow 2(c^2 - 1) = c^2 \left(\frac{10 - 6}{3} \right)$$

$$\Rightarrow 2(c^2 - 1) = c^2 \left(\frac{4}{3} \right)$$

$$\Rightarrow 6(c^2 - 1) = 4c^2$$

$$\Rightarrow 6c^2 - 6 = 4c^2$$

$$\Rightarrow 6c^2 - 4c^2 = 6$$

$$\Rightarrow 2c^2 = 6$$

$$\Rightarrow c^2 = \frac{6}{2}$$

$$\Rightarrow c^2 = 3$$

$$\Rightarrow c = \pm \sqrt{3} \in (-3, 4)$$

Hence, Lagrange's mean value theorem is verified.

(xii) $f(x) = x(x + 4)^2$ on $[0, 4]$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

Solution:

Given $f(x) = x(x+4)^2$ on $[0, 4]$

$$= x[(x)^2 + 2(4)(x) + (4)^2]$$

$$= x(x^2 + 8x + 16)$$

$$= x^3 + 8x^2 + 16x \text{ on } [0, 4]$$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[0, 4]$ and differentiable in $(0, 4)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (0, 4)$ such that:

$$f'(c) = \frac{f(4) - f(0)}{4 - 0}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(0)}{4}$$

$$f(x) = x^3 + 8x^2 + 16x$$

Differentiating with respect to x :

$$f'(x) = 3x^2 + 8(2x) + 16$$

$$= 3x^2 + 16x + 16$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = 3c^2 + 16c + 16$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f(4) = (4)^3 + 8(4)^2 + 16(4)$$

$$= 64 + 128 + 64$$

$$= 256$$

For $f(0)$, put the value of $x = 0$ in $f(x)$:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$f(0) = (0)^3 + 8(0)^2 + 16(0)$$

$$= 0 + 0 + 0$$

$$= 0$$

$$f'(c) = \frac{f(4) - f(0)}{4}$$

$$\Rightarrow 3c^2 + 16c + 16 = \frac{256 - 0}{4}$$

$$\Rightarrow 3c^2 + 16c + 16 = \frac{256}{4}$$

$$\Rightarrow 3c^2 + 16c + 16 = 64$$

$$\Rightarrow 3c^2 + 16c + 16 - 64 = 0$$

$$\Rightarrow 3c^2 + 16c - 48 = 0$$

For quadratic equation, $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow c = \frac{-(16) \pm \sqrt{(16)^2 - 4 \times 3 \times (-48)}}{2 \times 3}$$

$$\Rightarrow c = \frac{-16 \pm \sqrt{256 + 576}}{6}$$

$$\Rightarrow c = \frac{-16 \pm \sqrt{832}}{6}$$

$$\Rightarrow c = \frac{-16 \pm 8\sqrt{13}}{6}$$

$$\Rightarrow c = \frac{-16 \pm \sqrt{832}}{6}$$

$$\Rightarrow c = \frac{-16 \pm 8\sqrt{13}}{6}$$

$$\Rightarrow c = \frac{-16}{6} \pm \frac{8\sqrt{13}}{6}$$

$$\Rightarrow c = \frac{-8}{3} \pm \frac{4\sqrt{13}}{3}$$

$$\Rightarrow c = \frac{-8}{3} + \frac{4\sqrt{13}}{3}, \frac{-8}{3} - \frac{4\sqrt{13}}{3} \in c$$

Hence, Lagrange's mean value theorem is verified.

(xiii) $f(x) = \sqrt{x^2 - 4}$ on $[2, 4]$

Solution:

$$\Rightarrow f'(x) = \frac{1}{2}(x^2 - 4)^{-\frac{1}{2}}(2x)$$

$$\Rightarrow f'(x) = \frac{2x}{2(x^2 - 4)^{\frac{1}{2}}}$$

$$\Rightarrow f'(x) = \frac{x}{\sqrt{x^2 - 4}}$$

Here also, $\sqrt{x^2 - 4} > 0$

$\Rightarrow f'(x)$ exists for all values of x except $(2, -2)$

$\therefore f(x)$ is differentiable in $(2, 4)$

So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (2, 4)$ such that:

$$f'(c) = \frac{f(4) - f(2)}{4 - 2}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(2)}{2}$$

$$f(x) = \sqrt{x^2 - 4}$$

On differentiating with respect to x :

$$f'(x) = \frac{x}{\sqrt{x^2 - 4}}$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$: For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = \frac{c}{\sqrt{c^2 - 4}}$$

$$f'(c) = \frac{c}{\sqrt{c^2 - 4}}$$

For $f(4)$, put the value of $x = 4$ in $f(x)$: For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f(4) = \sqrt{4^2 - 4}$$

$$f(4) = \sqrt{4^2 - 4}$$

$$\Rightarrow f(4) = (16 - 4)^{\frac{1}{2}}$$

$$\Rightarrow f(4) = (16 - 4)^{\frac{1}{2}}$$

$$\Rightarrow f(4) = \sqrt{12}$$

$$\Rightarrow f(4) = \sqrt{12}$$

$$\Rightarrow c^2 = 3c^2 - 12$$

$$\Rightarrow -2c^2 = -12$$

$$\Rightarrow c^2 = \frac{-12}{-2}$$

$$\Rightarrow c^2 = 6$$

$$\Rightarrow c = \pm\sqrt{6}$$

$$\Rightarrow c = \sqrt{6} \in (2, 4)$$

Hence, Lagrange's mean value theorem is verified.

$$\Rightarrow c = \sqrt{6} \in (2, 4)$$

Hence, Lagrange's mean value theorem is verified.

(xiv) $f(x) = x^2 + x - 1$ on $[0, 4]$

Solution:

Given $f(x) = x^2 + x - 1$ on $[0, 4]$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[0, 4]$ and differentiable in <https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

(0, 4). So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (0, 4)$ such that:

$$f'(c) = \frac{f(4) - f(0)}{4 - 0}$$
$$\Rightarrow f'(c) = \frac{f(4) - f(0)}{4}$$

$$f(x) = x^2 + x - 1$$

Differentiating with respect to x:

$$f'(x) = 2x + 1$$

For $f'(c)$, put the value of $x = c$ in $f'(x)$:

$$f'(c) = 2c + 1$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f(4) = (4)^2 + 4 - 1$$

$$= 16 + 4 - 1$$

$$= 19$$

For $f(0)$, put the value of $x = 0$ in $f(x)$:

$$f(0) = (0)^2 + 0 - 1$$

$$= 0 + 0 - 1$$

$$= -1$$

$$f'(c) = \frac{f(4) - f(0)}{4}$$

$$\Rightarrow 2c + 1 = \frac{19 - (-1)}{4}$$

$$\Rightarrow 2c + 1 = \frac{20}{4}$$

$$\Rightarrow 2c + 1 = 5$$

$$\Rightarrow 2c = 5 - 1$$

$$\Rightarrow 2c = 4$$

$$\Rightarrow c = \frac{4}{2} = 2 \in (0, 4)$$

Hence, Lagrange's mean value theorem is verified.

$$\Rightarrow 2c + 1 = 5$$

$$\Rightarrow 2c = 5 - 1$$

$$\Rightarrow 2c = 4$$

$$\Rightarrow c = \frac{4}{2} = 2 \in (0, 4)$$

Hence, Lagrange's mean value theorem is verified.

(xv) $f(x) = \sin x - \sin 2x - x$ on $[0, \pi]$

Solution:

Given $f(x) = \sin x - \sin 2x - x$ on $[0, \pi]$

$\sin x$ and $\cos x$ functions are continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (0, \pi)$ such that:

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$f'(c) = \frac{f(\pi) - f(0)}{\pi - 0}$$

$$\Rightarrow f'(c) = \frac{f(\pi) - f(0)}{\pi}$$

$$f(x) = \sin x - \sin 2x - x$$

Differentiating with respect to x:

$$f(x) = \sin x - \sin 2x - x$$

$$\Rightarrow f'(x) = \cos x - \cos 2x \frac{d(2x)}{dx} - 1$$

$$\Rightarrow f'(x) = \cos x - 2\cos 2x - 1$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = \cos c - 2\cos 2c - 1$$

For $f(\pi)$, put the value of $x = \pi$ in $f(x)$:

$$f(\pi) = \sin \pi - \sin 2\pi - \pi$$

$$= 0 - 0 - \pi$$

$$= -\pi$$

For $f(0)$, put the value of $x=0$ in $f(x)$:

$$f(0) = \sin 0 - \sin 2(0) - 0$$

$$= -\pi$$

For $f(0)$, put the value of $x=0$ in $f(x)$:

$$f(0) = \sin 0 - \sin 2(0) - 0$$

$$= \sin 0 - \sin 0 - 0$$

$$= 0 - 0 - 0$$

$$= 0$$

$$f'(c) = \frac{f(\pi) - f(0)}{\pi}$$

$$\Rightarrow \cos c - 2\cos 2c - 1 = \frac{-\pi - 0}{\pi}$$

$$\Rightarrow \cos c - 2\cos 2c - 1 = -1$$

$$\Rightarrow \cos c - 2(2\cos^2 c - 1) = -1 + 1$$

$$\Rightarrow \cos c - 4\cos^2 c + 2 = 0$$

$$\Rightarrow 4\cos^2 c - \cos c - 2 = 0$$

For quadratic equation, $ax^2 + bx + c = 0$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$\Rightarrow \cos c = \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \times 4 \times (-2)}}{2 \times 4}$$

$$\Rightarrow \cos c = \frac{1 \pm \sqrt{1 + 32}}{8}$$

$$\Rightarrow \cos c = \frac{1 \pm \sqrt{33}}{8}$$

$$\Rightarrow c = \cos^{-1} \left(\frac{1 \pm \sqrt{33}}{8} \right) \in (0, \pi)$$

Hence, Lagrange's mean value theorem is verified.

(xvi) $f(x) = x^3 - 5x^2 - 3x$ on $[1, 3]$

Solution:

Given $f(x) = x^3 - 5x^2 - 3x$ on $[1, 3]$

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[1, 3]$ and differentiable in $(1, 3)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (1, 3)$ such that:

$$f'(c) = \frac{f(3) - f(1)}{3 - 1}$$

$$\Rightarrow f'(c) = \frac{f(3) - f(1)}{2}$$

$$f(x) = x^3 - 5x^2 - 3x$$

Differentiating with respect to x :

$$f'(x) = 3x^2 - 5(2x) - 3$$

$$= 3x^2 - 10x - 3$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = 3c^2 - 10c - 3$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

For $f(3)$, put the value of $x = 3$ in $f(x)$:

$$f(3) = (3)^3 - 5(3)^2 - 3(3)$$

$$= 27 - 45 - 9$$

$$= -27$$

For $f(1)$, put the value of $x = 1$ in $f(x)$:

$$f(1) = (1)^3 - 5(1)^2 - 3(1)$$

$$= 1 - 5 - 3$$

$$= -7$$

$$\begin{aligned}f'(c) &= \frac{f(3) - f(1)}{2} \\ \Rightarrow 3c^2 - 10c - 3 &= \frac{(-27) - (-7)}{2} \\ \Rightarrow 3c^2 - 10c - 3 &= \frac{-27+7}{2} \\ \Rightarrow 3c^2 - 10c - 3 &= \frac{-20}{2} \\ \Rightarrow 3c^2 - 10c - 3 &= -10 \\ \Rightarrow 3c^2 - 10c - 3 + 10 &= 0 \\ \Rightarrow 3c^2 - 10c + 7 &= 0 \\ \Rightarrow 3c^2 - 7c - 3c + 7 &= 0 \\ \Rightarrow c(3c - 7) - 1(3c - 7) &= 0 \\ \Rightarrow (3c - 7)(c - 1) &= 0 \\ \Rightarrow c = \frac{7}{3}, 1 \\ \Rightarrow c = \frac{7}{3} \in (1, 3)\end{aligned}$$

Hence, Lagrange's mean value theorem is verified.

$$\begin{aligned}\Rightarrow 3c^2 - 10c - 3 &= -10 \\ \Rightarrow 3c^2 - 10c - 3 + 10 &= 0 \\ \Rightarrow 3c^2 - 10c + 7 &= 0 \\ \Rightarrow 3c^2 - 7c - 3c + 7 &= 0 \\ \Rightarrow c(3c - 7) - 1(3c - 7) &= 0 \\ \Rightarrow (3c - 7)(c - 1) &= 0\end{aligned}$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow c = \frac{7}{3}, 1$$

$$\Rightarrow c = \frac{7}{3} \in (1, 3)$$

Hence, Lagrange's mean value theorem is verified.

2. Discuss the applicability of Lagrange's mean value theorem for the function $f(x) = |x|$ on $[-1, 1]$.

Solution:

Given $f(x) = |x|$ on $[-1, 1]$

So $f(x)$ can be defined as $= \begin{cases} -x, & x < 0 \\ x, & x \geq 0 \end{cases}$

For differentiability at $x = 0$,

$$\text{LHD} = \lim_{x \rightarrow 0^-} \frac{f(0-h) - f(0)}{-h}$$

{Since $f(x) = -x, x < 0$ }

$$= \lim_{x \rightarrow 0^-} \frac{-(0-h) - 0}{-h}$$

$$= \lim_{x \rightarrow 0^-} \frac{h - 0}{-h}$$

$$= \lim_{x \rightarrow 0^-} \frac{h}{-h}$$

$$= -1$$

$$\text{RHD} = \lim_{x \rightarrow 0^+} \frac{f(0-h) - f(0)}{-h}$$

{Since $f(x) = x, x > 0$ }

{Since $f(x) = x, x > 0$ }

$$= \lim_{x \rightarrow 0^-} \frac{(0 - h) - 0}{-h}$$

$$= \lim_{x \rightarrow 0^-} \frac{-h - 0}{-h}$$

$$= \lim_{x \rightarrow 0^-} \frac{-h}{-h}$$

$$= 1$$

LHD \neq RHD

$\Rightarrow f(x)$ is not differential at $x=0$

\therefore Lagrange's mean value theorem is not applicable for the function $f(x) = |x|$ on $[-1, 1]$.

3. Show that the Lagrange's mean value theorem is not applicable to the function $f(x) = 1/x$ on $[-1, 1]$.

Solution:

Given

$$f(x) = \frac{1}{x} \text{ on } [-1, 1]$$

Here, $x \neq 0$

$\Rightarrow f(x)$ exists for all values of x except 0

$\Rightarrow f(x)$ is discontinuous at $x=0$

$\therefore f(x)$ is not continuous in $[-1, 1]$

Hence the Lagrange's mean value theorem is not applicable to the function $f(x) = 1/x$ on $[-1, 1]$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

4. Verify the hypothesis and conclusion of Lagrange's mean value theorem for the function

$$f(x) = \frac{1}{4x - 1}, 1 \leq x \leq 4.$$

Solution:

Given

$$f(x) = \frac{1}{4x - 1} \text{ on } [1, 4]$$

Where $4x - 1 > 0$

$f'(x)$ has unique values for all x except $\frac{1}{4}$

$\therefore f(x)$ is continuous in $[1, 4]$

$$f(x) = \frac{1}{4x - 1}$$

Differentiating with respect to x :

$$f'(x) = (-1)(4x - 1)^{-2}(4)$$

$$\Rightarrow f'(x) = -\frac{4}{(4x - 1)^2}$$

Here, $4x - 1 > 0$

$f'(x)$ has unique values for all x except $\frac{1}{4}$

$\therefore f(x)$ is differentiable in $(1, 4)$

So both the necessary conditions of Lagrange's mean value theorem is satisfied. Therefore, there exist a point $c \in (1, 4)$ such that:

$$f'(c) = \frac{f(4) - f(1)}{4 - 1}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(1)}{3}$$

$$f(x) = \frac{1}{4x - 1}$$

On differentiating with respect to x:

$$f'(x) = -\frac{4}{(4x - 1)^2}$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = -\frac{4}{(4c - 1)^2}$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f'(c) = -\frac{4}{(4c-1)^2}$$

For $f(4)$, put the value of $x = 4$ in $f(x)$:

$$f(4) = \frac{1}{4(4) - 1}$$

$$\Rightarrow f(4) = \frac{1}{16 - 1}$$

$$\Rightarrow f(4) = \frac{1}{15}$$

For $f(1)$, put the value of $x = 1$ in $f(x)$:

$$f(1) = \frac{1}{4(1) - 1}$$

$$\Rightarrow f(1) = \frac{1}{4 - 1}$$

$$\Rightarrow f(1) = \frac{1}{3}$$

$$\Rightarrow f'(c) = \frac{f(4) - f(1)}{3}$$

$$\Rightarrow -\frac{4}{(4c-1)^2} = \frac{\frac{1}{15} - \frac{1}{3}}{3}$$

$$\Rightarrow -3(4) = (4c-1)^2 \left(\frac{1}{15} - \frac{1}{3} \right)$$

$$\Rightarrow -12 = (4c-1)^2 \left(\frac{3-15}{45} \right)$$

$$\Rightarrow -12 = (4c-1)^2 \left(\frac{-12}{45} \right)$$

$$\Rightarrow -12 \times \frac{45}{-12} = (4c-1)^2$$

$$\Rightarrow -12 \times \frac{45}{-12} = (4c - 1)^2$$

$$\Rightarrow (4c - 1)^2 = 45$$

$$\Rightarrow (4c - 1) = \pm\sqrt{45}$$

$$\Rightarrow (4c - 1) = \pm 3\sqrt{5}$$

$$\Rightarrow c = \frac{\pm 3\sqrt{5} + 1}{4}$$

$$\Rightarrow c = \frac{3\sqrt{5} + 1}{4} \approx 1.92 \in (1, 4)$$

Hence, Lagrange's mean value theorem is verified.

5. Find a point on the parabola $y = (x - 4)^2$, where the tangent is parallel to the chord joining (4, 0) and (5, 1).

Solution:

Given $f(x) = (x - 4)^2$ on $[4, 5]$

This interval $[a, b]$ is obtained by x - coordinates of the points of the chord.

Every polynomial function is continuous everywhere on $(-\infty, \infty)$ and differentiable for all arguments. Here, $f(x)$ is a polynomial function. So it is continuous in $[4, 5]$ and differentiable in $(4, 5)$. So both the necessary conditions of Lagrange's mean value theorem is satisfied.

Therefore, there exist a point $c \in (4, 5)$ such that:

$$f'(c) = \frac{f(5) - f(4)}{5 - 4}$$

$$\Rightarrow f'(c) = \frac{f(5) - f(4)}{1}$$

$$f(x) = (x - 4)^2$$

Differentiating with respect to x:

$$f'(x) = 2(x - 4) \frac{d(x - 4)}{dx}$$

$$\Rightarrow f'(x) = 2(x - 4)(1)$$

$$\Rightarrow f'(x) = 2(x - 4)$$

For $f'(c)$, put the value of $x=c$ in $f'(x)$:

$$f'(c) = 2(c - 4)$$

For $f(5)$, put the value of $x=5$ in $f(x)$:

$$f(5) = (5 - 4)^2$$

$$= (1)^2$$

$$= 1$$

For $f(4)$, put the value of $x=4$ in $f(x)$:

$$f(4) = (4 - 4)^2$$

$$= (0)^2$$

$$= 0$$

$$f'(c) = f(5) - f(4)$$

$$\Rightarrow 2(c - 4) = 1 - 0$$

$$\Rightarrow 2c - 8 = 1$$

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

$$\Rightarrow 2c = 1 + 8$$

$$\Rightarrow c = \frac{9}{2} = 4.5 \in (4, 5)$$

We know that, the value of c obtained in Lagrange's Mean value Theorem is nothing but the value of x – coordinate of the point of the contact of the tangent to the curve which is parallel to the chord joining the points $(4, 0)$ and $(5, 1)$.

Now, put this value of x in $f(x)$ to obtain y :

$$y = (x - 4)^2$$

$$\Rightarrow y = \left(\frac{9}{2} - 4\right)^2$$

$$\Rightarrow y = \left(\frac{9 - 8}{2}\right)^2$$

$$\Rightarrow y = \left(\frac{1}{2}\right)^2$$

$$\Rightarrow y = \frac{1}{4}$$

Hence, the required point is $\left(\frac{9}{2}, \frac{1}{4}\right)$



Chapterwise RD Sharma Solutions for Class 12 Maths :

- Chapter 1–Relation
- Chapter 2–Functions
- Chapter 3–Binary Operations
- Chapter 4–Inverse Trigonometric Functions
- Chapter 5–Algebra of Matrices
- Chapter 6–Determinants
- Chapter 7–Adjoint and Inverse of a Matrix
- Chapter 8–Solution of Simultaneous Linear Equations
- Chapter 9–Continuity
- Chapter 10–Differentiability
- Chapter 11–Differentiation
- Chapter 12–Higher Order Derivatives
- Chapter 13–Derivatives as a Rate Measurer
- Chapter 14–Differentials, Errors and Approximations
- Chapter 15–Mean Value Theorems
- Chapter 16–Tangents and Normals
- Chapter 17–Increasing and Decreasing Functions
- Chapter 18–Maxima and Minima
- Chapter 19–Indefinite Integrals

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-value-theorems/>

About RD Sharma

RD Sharma isn't the kind of author you'd bump into at lit fests. But his bestselling books have helped many CBSE students lose their dread of maths. Sunday Times profiles the tutor turned internet star

He dreams of algorithms that would give most people nightmares. And, spends every waking hour thinking of ways to explain concepts like 'series solution of linear differential equations'. Meet Dr Ravi Dutt Sharma — mathematics teacher and author of 25 reference books — whose name evokes as much awe as the subject he teaches. And though students have used his thick tomes for the last 31 years to ace the dreaded maths exam, it's only recently that a spoof video turned the tutor into a YouTube star.

R D Sharma had a good laugh but said he shared little with his on-screen persona except for the love for maths. "I like to spend all my time thinking and writing about maths problems. I find it relaxing," he says. When he is not writing books explaining mathematical concepts for classes 6 to 12 and engineering students, Sharma is busy dispensing his duty as vice-principal and head of department of science and humanities at Delhi government's Guru Nanak Dev Institute of Technology.

<https://www.indcareer.com/schools/rd-sharma-solutions-for-class-12-maths-chapter-15-mean-variance-theorems/>