


# Maharashtra Board Solutions Class 12-Arts & Science Maths (Part 2): Chapter 2- Applications of Derivatives

## Class 12 - Chapter 2 Applications of Derivatives



For any clarifications or questions you can write to [info@indcareer.com](mailto:info@indcareer.com)

### Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015  
Maharashtra, India

WhatsApp: +91 9561 204 888, Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>



# Maharashtra Board Solutions Class 12-Arts & Science Maths (Part 2): Chapter 2- Applications of Derivatives

Class 12: Maths Chapter 2 solutions. Complete Class 12 Maths Chapter 2 Notes.

## Maharashtra Board Solutions Class 12-Arts & Science Maths (Part 2): Chapter 2- Applications of Derivatives

Maharashtra Board 12th Maths Chapter 2, Class 12 Maths Chapter 2 solutions

**Ex 2.1**

**Question 1.**

**Find the equations of tangents and normals to the curve at the point on it.**

**(i)  $y = x^2 + 2e^x + 2$  at  $(0, 4)$**

**Solution:**

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-part-2-chapter-2-applications-of-derivatives/>

$$y = x^2 + 2e^x + 2$$

$$\therefore \frac{dy}{dx} = \frac{d}{dx}(x^2 + 2e^x + 2)$$

$$= 2x + 2 \times e^x + 0 = 2x + 2e^x$$

$$\therefore \left(\frac{dy}{dx}\right)_{\text{at } (0, 4)} = 2(0) + 2e^0 = 2$$

= slope of the tangent at (0, 4)

$\therefore$  the equation of the tangent at (0, 4) is

$$y - 4 = 2(x - 0)$$

$$\therefore y - 4 = 2x$$

$$\therefore 2x - y + 4 = 0$$

The slope of the normal at (0, 4)

$$= \frac{-1}{\left(\frac{dy}{dx}\right)_{\text{at } (0, 4)}} = -\frac{1}{2}$$

$\therefore$  the equation of the normal at (0, 4) is

$$y - 4 = -\frac{1}{2}(x - 0)$$

$$\therefore 2y - 8 = -x \quad \therefore x + 2y - 8 = 0$$

Hence, the equations of tangent and normal are

$2x - y + 4 = 0$  and  $x + 2y - 8 = 0$  respectively.

(ii)  $x^3 + y^3 - 9xy = 0$  at  $(2, 4)$

Solution:

$$x^3 + y^3 - 9xy = 0$$

Differentiating both sides w.r.t.  $x$ , we get

$$3x^2 + 3y^2 \frac{dy}{dx} - 9 \left[ x \frac{dy}{dx} + y \cdot \frac{d}{dx}(x) \right] = 0$$

$$\therefore 3x^2 + 3y^2 \frac{dy}{dx} - 9x \frac{dy}{dx} - 9y \times 1 = 0$$

$$\therefore (3y^2 - 9x) \frac{dy}{dx} = 9y - 3x^2$$

$$\therefore \frac{dy}{dx} = \frac{9y - 3x^2}{3y^2 - 9x}$$

$$\begin{aligned} \therefore \left( \frac{dy}{dx} \right)_{\text{at}(2,4)} &= \frac{9(4) - 3(2)^2}{3(4)^2 - 9(2)} \\ &= \frac{36 - 12}{48 - 18} = \frac{24}{30} = \frac{4}{5} \end{aligned}$$

= slope of the tangent at  $(2, 4)$

$\therefore$  the equation of the tangent at  $(2, 4)$  is

$$y - 4 = \frac{4}{5}(x - 2)$$

$$\therefore 5y - 20 = 4x - 8$$

$$\therefore 4x - 5y + 12 = 0$$

$$\text{The slope of normal at } (2, 4) = \frac{-1}{\left(\frac{dy}{dx}\right)_{\text{at } (2, 4)}} = -\frac{5}{4}$$

$\therefore$  the equation of the normal at  $(2, 4)$  is

$$y - 4 = -\frac{5}{4}(x - 2)$$

$$\therefore 4y - 16 = -5x + 10$$

$$\therefore 5x + 4y - 26 = 0$$

Hence, the equations of tangent and normal are  $4x - 5y + 12 = 0$  and  $5x + 4y - 26 = 0$  respectively.

(iii)  $x^2 - \sqrt{3}xy + 2y^2 = 5$  at  $(\sqrt{3}, 2)$

Solution:

$$x^2 - \sqrt{3}xy + 2y^2 = 5$$

Differentiating both sides w.r.t.  $x$ , we get

$$2x - \sqrt{3} \left[ x \frac{dy}{dx} + y \cdot \frac{d}{dx}(x) \right] + 2 \times 2y \frac{dy}{dx} = 0$$

$$\therefore 2x - \sqrt{3}x \frac{dy}{dx} - \sqrt{3}y \times 1 + 4y \frac{dy}{dx} = 0$$

$$\therefore (4y - \sqrt{3}x) \frac{dy}{dx} = \sqrt{3}y - 2x$$

$$\therefore \frac{dy}{dx} = \frac{\sqrt{3}y - 2x}{4y - \sqrt{3}x} = \frac{2x - \sqrt{3}x}{\sqrt{3}x - 4y}$$

$$\therefore \left( \frac{dy}{dx} \right)_{\text{at}(\sqrt{3}, 2)} = \frac{2\sqrt{3} - \sqrt{3}(2)}{\sqrt{3}(\sqrt{3}) - 4(2)} = 0$$

= slope of the tangent at  $(\sqrt{3}, 2)$

$\therefore$  the equation of the tangent at  $(\sqrt{3}, 2)$  is

$$y - 2 = 0(x - \sqrt{3})$$

$$\therefore y - 2 = 0 \quad \therefore y = 2$$

The slope of normal at  $(\sqrt{3}, 2)$

$$= \frac{-1}{\left( \frac{dy}{dx} \right)_{\text{at}(\sqrt{3}, 2)}} \quad \text{where} \quad \left( \frac{dy}{dx} \right)_{\text{at}(\sqrt{3}, 2)} = 0$$

the slope of normal at  $(\sqrt{3}, 2)$  does not exist.

normal is parallel to Y-axis.

equation of the normal is of the form  $x = k$

Since, it passes through the point  $(\sqrt{3}, 2)$ ,  $k = \sqrt{3}$

equation of the normal is  $x = \sqrt{3}$ .

Hence, the equations of tangent and normal are  $y = 2$  and  $x = \sqrt{3}$  respectively.

---

(iv)  $2xy + \pi \sin y = 2\pi$  at  $(1, \frac{\pi}{2})$

Solution:

$$2xy + \pi \sin y = 2\pi$$

Differentiating both sides w.r.t.  $x$ , we get

$$2 \left[ x \frac{dy}{dx} + y \cdot \frac{d}{dx}(x) \right] + \pi \cos y \frac{dy}{dx} = 0$$

$$\therefore 2x \frac{dy}{dx} + 2y \times 1 + \pi \cos y \frac{dy}{dx} = 0$$

$$\therefore (2x + \pi \cos y) \frac{dy}{dx} = -2y$$

$$\therefore \frac{dy}{dx} = \frac{-2y}{2x + \pi \cos y}$$

$$\therefore \left( \frac{dy}{dx} \right)_{\text{at} \left( 1, \frac{\pi}{2} \right)} = \frac{-2 \left( \frac{\pi}{2} \right)}{2(1) + \pi \cos \frac{\pi}{2}}$$

$$= \frac{-\pi}{2 + \pi(0)} = -\frac{\pi}{2}$$

$$= \text{slope of the tangent at} \left( 1, \frac{\pi}{2} \right)$$

$\therefore$  the equation of the tangent at  $\left( 1, \frac{\pi}{2} \right)$  is

$$y - \frac{\pi}{2} = -\frac{\pi}{2}(x - 1)$$

$$\therefore 2y - \pi = -\pi x + \pi$$

$$\therefore \pi x + 2y - 2\pi = 0$$

The slope of normal at  $\left(1, \frac{\pi}{2}\right)$

$$= \frac{-1}{\left(\frac{dy}{dx}\right)_{\text{at}\left(1, \frac{\pi}{2}\right)}} = \frac{-1}{\left(-\frac{\pi}{2}\right)} = \frac{2}{\pi}$$

$\therefore$  the equation of the normal at  $\left(1, \frac{\pi}{2}\right)$  is

$$y - \frac{\pi}{2} = \frac{2}{\pi}(x - 1)$$

$$\therefore \pi y - \frac{\pi^2}{2} = 2x - 2$$

$$\therefore 2\pi y - \pi^2 = 4x - 4$$

$$\therefore 4x - 2\pi y + \pi^2 - 4 = 0$$

Hence, the equations of tangent and normal are  $\pi x + 2y - 2\pi = 0$  and  $4x - 2\pi y + \pi^2 - 4 = 0$  respectively.

(v)  $x \sin 2y = y \cos 2x$  at  $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

Solution:

$$x \sin 2y = y \cos 2x$$

Differentiating both sides w.r.t.  $x$ , we get

$$x \frac{d}{dx}(\sin 2y) + \sin 2y \cdot \frac{d}{dx}(x) = y \cdot \frac{d}{dx}(\cos 2x) + \cos 2x \cdot \frac{dy}{dx}$$

$$\therefore x \cdot \cos 2y \cdot \frac{d}{dx}(2y) + \sin 2y \times 1$$

$$= y \cdot (-\sin 2x) \cdot \frac{d}{dx}(2x) + \cos 2x \cdot \frac{dy}{dx}$$

$$\therefore x \cos 2y \times 2 \frac{dy}{dx} + \sin 2y = -y \sin 2x \times 2 + \cos 2x \cdot \frac{dy}{dx}$$

$$\therefore (2x \cos 2y - \cos 2x) \frac{dy}{dx} = -2y \sin 2x - \sin 2y$$

$$\therefore \frac{dy}{dx} = \frac{-2y \sin 2x - \sin 2y}{2x \cos 2y - \cos 2x}$$

$$\begin{aligned} \therefore \left(\frac{dy}{dx}\right)_{\text{at}\left(\frac{\pi}{4}, \frac{\pi}{2}\right)} &= \frac{-2\left(\frac{\pi}{2}\right)\sin\frac{\pi}{2} - \sin\pi}{2\left(\frac{\pi}{4}\right)\cos\pi - \cos\frac{\pi}{2}} \\ &= \frac{-\pi(1) - 0}{\frac{\pi}{2}(-1) - 0} = \frac{-\pi}{\left(\frac{-\pi}{2}\right)} = 2 \end{aligned}$$

= slope of the tangent at  $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

∴ the equation of the tangent at  $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$  is

$$y - \frac{\pi}{2} = 2\left(x - \frac{\pi}{4}\right)$$

$$\therefore y - \frac{\pi}{2} = 2x - \frac{\pi}{2}$$

$$\therefore 2x - y = 0$$

The slope of normal at  $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$

$$= \frac{-1}{\left(\frac{dy}{dx}\right)_{\text{at}\left(\frac{\pi}{4}, \frac{\pi}{2}\right)}} = \frac{-1}{2}$$

∴ the equation of the normal at  $\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$  is

$$y - \frac{\pi}{2} = -\frac{1}{2}\left(x - \frac{\pi}{4}\right)$$

$$\therefore 2y - \pi = -x + \frac{\pi}{4}$$

$$\therefore 8y - 4\pi = -4x + \pi$$

$$\therefore 4x + 8y - 5\pi = 0$$

---

Hence, the equations of the tangent and normal are  $2x - y = 0$  and  $4x + 8y - 5n = 0$  respectively.

(vi)  $x = \sin \theta$  and  $y = \cos 2\theta$  at  $\theta = \frac{\pi}{6}$

Solution:

When  $\theta = \frac{\pi}{6}$ ,  $x = \sin \frac{\pi}{6}$  and  $y = \cos \frac{\pi}{3}$

$\therefore x = \frac{1}{2}$  and  $y = \frac{1}{2}$

Hence, the point at which we want to find the equations of tangent and normal is  $(\frac{1}{2}, \frac{1}{2})$

Now,  $x = \sin \theta$ ,  $y = \cos 2\theta$

Differentiating  $x$  and  $y$  w.r.t.  $\theta$ , we get

$$\frac{dx}{d\theta} = \frac{d}{d\theta}(\sin \theta) = \cos \theta$$

$$\begin{aligned}\text{and } \frac{dy}{d\theta} &= \frac{d}{d\theta}(\cos 2\theta) = -\sin 2\theta \cdot \frac{d}{d\theta}(2\theta) \\ &= -\sin 2\theta \times 2 = -2 \sin 2\theta\end{aligned}$$

$$\therefore \frac{dy}{dx} = \frac{(dy/d\theta)}{(dx/d\theta)} = \frac{-2 \sin 2\theta}{\cos \theta}$$

$$\therefore \left(\frac{dy}{dx}\right)_{\text{at } \theta = \frac{\pi}{6}} = \frac{-2 \sin \frac{\pi}{3}}{\cos \frac{\pi}{6}} = \frac{-2\left(\frac{\sqrt{3}}{2}\right)}{\left(\frac{\sqrt{3}}{2}\right)} = -2$$

= slope of the tangent at  $\theta = \frac{\pi}{6}$

$\therefore$  the equation of the tangent at  $\theta = \frac{\pi}{6}$  i.e. at  $\left(\frac{1}{2}, \frac{1}{2}\right)$  is

$$y - \frac{1}{2} = -2\left(x - \frac{1}{2}\right)$$

$$\therefore y - \frac{1}{2} = -2x + 1$$

$$\therefore 2y - 1 = -4x + 2$$

$$\therefore 4x + 2y - 3 = 0$$

The slope of normal at  $\theta = \frac{\pi}{6}$

$$= -\frac{1}{\left(\frac{dy}{dx}\right)_{\text{at } \theta = \frac{\pi}{6}}} = \frac{-1}{-2} = \frac{1}{2}$$

$\therefore$  equation of the normal at  $\theta = \frac{\pi}{6}$ , i.e. at  $\left(\frac{1}{2}, \frac{1}{2}\right)$  is

$$y - \frac{1}{2} = \frac{1}{2}\left(x - \frac{1}{2}\right)$$

$$2y - 1 = x - \frac{1}{2}$$

$$4y - 2 = 2x - 1$$

$$2x - 4y + 1 = 0$$

---

$$2x - 4y + 1 = 0$$

Hence, equations of the tangent and normal are  $4x + 2y - 3 = 0$  and  $2x - 4y + 1 = 0$  respectively.

$$(vii) x = \sqrt{t}, y = t - \frac{1}{\sqrt{t}}, \text{ at } t = 4.$$

Solution:

$$\text{When } t = 4, x = \sqrt{4} \text{ and } y = 4 - \frac{1}{\sqrt{4}}$$

$$\therefore x = 2 \text{ and } y = 4 - \frac{1}{2} = \frac{7}{2}$$

Hence, the point at which we want to find the equations of tangent and normal is  $(2, \frac{7}{2})$ .

$$\text{Now, } x = \sqrt{t}, y = t - \frac{1}{\sqrt{t}}$$

Differentiating  $x$  and  $y$  w.r.t.  $t$ , we get

$$\frac{dx}{dt} = \frac{d}{dt}(\sqrt{t}) = \frac{1}{2\sqrt{t}}$$

$$\text{and } \frac{dy}{dt} = \frac{d}{dt}\left(t - \frac{1}{\sqrt{t}}\right) = 1 - \left(-\frac{1}{2}\right)t^{-\frac{3}{2}} = 1 + \frac{1}{2t^{\frac{3}{2}}}$$

$$= \frac{2t^{\frac{3}{2}} + 1}{2t^{\frac{3}{2}}}$$

$$\therefore \frac{dy}{dx} = \frac{(dy/dt)}{(dx/dt)} = \frac{\left(\frac{2t^{\frac{3}{2}} + 1}{2t^{\frac{3}{2}}}\right)}{\left(\frac{1}{2\sqrt{t}}\right)}$$

$$= \frac{2t^{\frac{3}{2}} + 1}{2t^{\frac{3}{2}}} \times 2\sqrt{t} = \frac{2t^{\frac{3}{2}} + 1}{t}$$

$$\therefore \left(\frac{dy}{dx}\right)_{\text{at } t=4} = \frac{2(4)^{\frac{3}{2}} + 1}{4} = \frac{2 \times 8 + 1}{4} = \frac{17}{4}$$

= slope of the tangent at  $t = 4$

$\therefore$  the equation of the tangent at  $t = 4$ , i.e. at  $\left(2, \frac{7}{2}\right)$  is

$$y - \frac{7}{2} = \frac{17}{4}(x - 2)$$

$$\therefore 4y - 14 = 17x - 34$$

$$\therefore 17x - 4y - 20 = 0$$

The slope of normal at  $t = 4$

$$= \frac{-1}{\left(\frac{dy}{dx}\right)_{\text{at } t=4}} = \frac{-1}{\left(\frac{17}{4}\right)} = -\frac{4}{17}$$

$\therefore$  the equation of the normal at  $t = 4$ , i.e. at  $\left(2, \frac{7}{2}\right)$  is

$$y - \frac{7}{2} = -\frac{4}{17}(x - 2)$$

---

$$\therefore 34y - 119 = -8x + 16$$

$$\therefore 8x + 34y - 135 = 0$$

Hence, the equations of tangent and normal are  $17x - 4y - 20 = 0$  and  $8x + 34y - 135 = 0$  respectively.

Question 2.

Find the point of the curve  $y = \sqrt{x-3}$  where the tangent is perpendicular to the line  $6x + 3y - 5 = 0$ .

Solution:

Let the required point on the curve  $y = \sqrt{x-3}$  be  $P(x_1, y_1)$ .

Differentiating  $y = \sqrt{x-3}$  w.r.t.  $x$ , we get

$$\frac{dy}{dx} = \frac{d}{dx}(\sqrt{x-3}) = \frac{1}{2\sqrt{x-3}} \cdot \frac{d}{dx}(x-3)$$

$$= \frac{1}{2\sqrt{x-3}} \times (1-0) = \frac{1}{2\sqrt{x-3}}$$

$\therefore$  slope of the tangent at  $(x_1, y_1)$

$$= \left(\frac{dy}{dx}\right)_{\text{at } (x_1, y_1)} = \frac{1}{2\sqrt{x_1-3}}$$

Since, this tangent is perpendicular to  $6x + 3y - 5 = 0$

whose slope is  $\frac{-6}{3} = -2$ ,

$$\text{slope of the tangent} = \frac{-1}{-2} = \frac{1}{2}$$

$$\therefore \frac{1}{2\sqrt{x_1-3}} = \frac{1}{2}$$

$$\therefore \sqrt{x_1-3} = 1$$

$$\therefore x_1 - 3 = 1 \quad \therefore x_1 = 4$$

Since,  $(x_1, y_1)$  lies on  $y = \sqrt{x-3}$ ,  $y_1 = \sqrt{x_1-3}$

When  $x_1 = 4$ ,  $y_1 = \sqrt{4-3} = \pm 1$

Hence, the required points are  $(4, 1)$  and  $(4, -1)$ .

Question 3.

Find the points on the curve  $y = x^3 - 2x^2 - x$  where the tangents are parallel to  $3x - y + 1 = 0$ .

Solution:

Let the required point on the curve  $y = x^3 - 2x^2 - x$  be  $P(x_1, y_1)$ .

Differentiating  $y = x^3 - 2x^2 - x$  w.r.t.  $x$ , we get

$$\frac{dy}{dx} = \frac{d}{dx}(x^3 - 2x^2 - x)$$

$$= 3x^2 - 2 \times 2x - 1 = 3x^2 - 4x - 1$$

$\therefore$  slope of the tangent at  $(x_1, y_1)$

$$= \left( \frac{dy}{dx} \right)_{\text{at } (x_1, y_1)} = 3x_1^2 - 4x_1 - 1$$

Since this tangent is parallel to  $3x - y + 1 = 0$  whose slope

$$\text{is } \frac{-3}{-1} = 3,$$

slope of the tangent = 3

$$\therefore 3x_1^2 - 4x_1 - 1 = 3$$

$$\therefore 3x_1^2 - 4x_1 - 4 = 0$$

$$\therefore 3x_1^2 - 6x_1 + 2x_1 - 4 = 0$$

$$\therefore 3x_1(x_1 - 2) + 2(x_1 - 2) = 0$$

$$\therefore (x_1 - 2)(3x_1 + 2) = 0$$

$$\therefore x_1 - 2 = 0 \quad \text{or} \quad 3x_1 + 2 = 0$$

$$\therefore x_1 = 2 \quad \text{or} \quad x_1 = -\frac{2}{3}$$

Since,  $(x_1, y_1)$  lies on  $y = x^3 - 2x^2 - x$ ,  $y_1 = x_1^3 - 2x_1^2 - x_1$

$$\text{When } x_1 = 2, y_1 = (2)^3 - 2(2)^2 - 2 = 8 - 8 - 2 = -2$$

$$\begin{aligned} \text{When } x_1 = -\frac{2}{3}, y_1 &= \left(-\frac{2}{3}\right)^3 - 2\left(-\frac{2}{3}\right)^2 + \frac{2}{3} \\ &= \frac{-8}{27} - \frac{8}{9} + \frac{2}{3} = \frac{-14}{27} \end{aligned}$$

Hence, the required points are  $(2, -2)$  and  $\left(-\frac{2}{3}, \frac{-14}{27}\right)$ .

Question 4.

Find the equations of the tangents to the curve  $x^2 + y^2 - 2x - 4y + 1 = 0$  which are parallel to the X-axis.

Solution:

Let P  $(x_1, y_1)$  be the point on the curve  $x^2 + y^2 - 2x - 4y + 1 = 0$  where the tangent is parallel to X-axis.

Differentiating  $x^2 + y^2 - 2x - 4y + 1 = 0$  w.r.t.  $x$ , we get

$$2x + 2y \frac{dy}{dx} - 2 \times 1 - 4 \frac{dy}{dx} + 0 = 0$$

$$\therefore (2y - 4) \frac{dy}{dx} = 2 - 2x$$

$$\therefore \frac{dy}{dx} = \frac{2 - 2x}{2y - 4} = \frac{1 - x}{y - 2}$$

$$\therefore \left( \frac{dy}{dx} \right)_{\text{at } (x_1, y_1)} = \frac{1 - x_1}{y_1 - 2}$$

= slope of the tangent at  $(x_1, y_1)$

Since, the tangent is parallel to X-axis,

slope of the tangent = 0.

$$\therefore \frac{1 - x_1}{y_1 - 2} = 0$$

$$\therefore 1 - x_1 = 0 \quad \therefore x_1 = 1$$

Since,  $(x_1, y_1)$  lies on  $x^2 + y^2 - 2x - 4y + 1 = 0$ ,

$$x_1^2 + y_1^2 - 2x_1 - 4y_1 + 1 = 0$$

When  $x_1 = 1$ ,  $(1)^2 + y_1^2 - 2(1) - 4y_1 + 1 = 0$

$$\therefore 1 + y_1^2 - 2 - 4y_1 + 1 = 0$$

$$\therefore y_1^2 - 4y_1 = 0$$

$$\therefore y_1(y_1 - 4) = 0$$

$$\therefore y_1 = 0 \text{ or } y_1 = 4$$

the coordinates of the points are  $(1, 0)$  or  $(1, 4)$

Since the tangents are parallel to X-axis, their equations are of the form  $y = k$

If it passes through the point  $(1, 0)$ ,  $k = 0$ , and if it passes through the point  $(1, 4)$ ,  $k = 4$

Hence, the equations of the tangents are  $y = 0$  and  $y = 4$ .

Question 5.

Find the equations of the normals to the curve  $3x^2 - y^2 = 8$ , which are parallel to the line  $x + 3y = 4$ .

Solution:

Let  $P(x_1, y_1)$  be the foot of the required normal to the curve  $3x^2 - y^2 = 8$ .

Differentiating  $3x^2 - y^2 = 8$  w.r.t.  $x$ , we get

$$3 \times 2x - 2y \frac{dy}{dx} = 0$$

$$\therefore -2y \frac{dy}{dx} = -6x$$

$$\therefore \frac{dy}{dx} = \frac{3x}{y}$$

$$\therefore \left( \frac{dy}{dx} \right)_{\text{at } (x_1, y_1)} = \frac{3x_1}{y_1}$$

= slope of the tangent at  $(x_1, y_1)$

$\therefore$  slope of the normal at  $P(x_1, y_1)$

$$= m_1 = \frac{-1}{\left( \frac{dy}{dx} \right)_{\text{at } (x_1, y_1)}} = -\frac{y_1}{3x_1}$$

The slope of line  $x + 3y = 4$  is  $m_2 = \frac{-1}{3}$

Since, the normal at  $P(x_1, y_1)$  is parallel to the line

$$x + 3y = 4, m_1 = m_2$$

$$\therefore -\frac{y_1}{3x_1} = -\frac{1}{3} \quad \therefore y_1 = x_1$$

Since,  $(x_1, y_1)$  lies on the curve  $3x^2 - y^2 = 8$ ,

$$3x_1^2 - y_1^2 = 8$$

$$\therefore 3x_1^2 - x_1^2 = 8 \quad \dots [\because y_1 = x_1]$$

$$\therefore 2x_1^2 = 8 \quad \therefore x_1^2 = 4$$

$$\therefore x_1 = \pm 2$$

When  $x_1 = 2, y_1 = 2$

When  $x_1 = -2, y_1 = -2$

$\therefore$  the coordinates of the point P are  $(2, 2)$  or  $(-2, -2)$

and the slope of the normal is  $m_1 = m_2 = -\frac{1}{3}$

$\therefore$  the equation of the normal at (2, 2) is

$$y - 2 = -\frac{1}{3}(x - 2)$$

$$\therefore 3y - 6 = -x + 2$$

$$\therefore x + 3y - 8 = 0$$

and the equation of the normal at (-2, -2) is

$$y + 2 = -\frac{1}{3}(x + 2)$$

$$\therefore 3y - 6 = -x + 2$$

$$\therefore x + 3y + 8 = 0$$

Hence, the equations of the normals are  $x + 3y - 8 = 0$  and  $x + 3y + 8 = 0$ .

Question 6.

If the line  $y = 4x - 5$  touches the curve  $y^2 = ax^3 + b$  at the point (2, 3), find a and b.

Solution:

$$y^2 = ax^3 + b$$

Differentiating both sides w.r.t. x, we get

$$2y \frac{dy}{dx} = a \times 3x^2 + 0$$

$$\therefore \frac{dy}{dx} = \frac{3ax^2}{2y}$$

$$\therefore \left(\frac{dy}{dx}\right)_{\text{at } (2, 3)} = \frac{3a(2)^2}{2(3)} = 2a$$

= slope of the tangent at (2, 3)

Since, the line  $y = 4x - 5$  touches the curve at the point (2, 3), slope of the tangent at (2, 3) is 4.

$$2a = 4 \Rightarrow a = 2$$

Since (2, 3) lies on the curve  $y^2 = ax^3 + b$

$$(3)^2 = a(2)^3 + b$$

$$9 = 8a + b$$

$$9 = 8(2) + b \dots\dots [\because a = 2]$$

$$b = -7$$

Hence,  $a = 2$  and  $b = -7$ .

Question 7.

A particle moves along the curve  $6y = x^3 + 2$ . Find the points on the curve at which y-coordinate is changing 8 times as fast as the x-coordinate.

Solution:

Let  $P(x_1, y_1)$  be the point on the curve  $6y = x^3 + 2$  whose y-coordinate is changing 8 times as fast as the x-coordinate.

$$\text{Then } \left(\frac{dy}{dt}\right)_{\text{at } (x_1, y_1)} = 8 \left(\frac{dx}{dt}\right)_{\text{at } (x_1, y_1)} \quad \dots (1)$$

Differentiating  $6y = x^3 + 2$  w.r.t.  $t$ , we get

$$6 \frac{dy}{dt} = \frac{d}{dt}(x^3 + 2) = 3x^2 \frac{dx}{dt} + 0$$

$$\therefore 2 \frac{dy}{dt} = x^2 \frac{dx}{dt}$$

$$\therefore 2 \left( \frac{dy}{dt} \right)_{\text{at } (x_1, y_1)} = x_1^2 \cdot \left( \frac{dx}{dt} \right)_{\text{at } (x_1, y_1)}$$

$$\therefore 2 \times 8 \left( \frac{dx}{dt} \right)_{\text{at } (x_1, y_1)} = x_1^2 \cdot \left( \frac{dx}{dt} \right)_{\text{at } (x_1, y_1)} \quad \dots \text{ [By (1)]}$$

$$\therefore x_1^2 = 16 \quad \therefore x_1 = \pm 4$$

Now,  $(x_1, y_1)$  lies on the curve  $6y = x^3 + 2$ .

$$\therefore 6y_1 = x_1^3 + 2$$

$$\text{When } x_1 = 4, 6y_1 = (4)^3 + 2 = 66 \quad \therefore y_1 = 11$$

$$\text{When } x_1 = -4, 6y_1 = (-4)^3 + 2 = -62 \quad \therefore y_1 = -\frac{31}{3}$$

Hence, the required points on the curve are  $(4, 11)$  and

$$\left( -4, -\frac{31}{3} \right).$$

Question 8.

A spherical soap bubble is expanding so that its radius is increasing at the rate of 0.02 cm/sec. At what rate is the surface area increasing, when its radius is 5 cm?

Solution:

Let  $r$  be the radius and  $S$  be the surface area of the soap bubble at any time  $t$ .

$$\text{Then } S = 4\pi r^2$$

Differentiating w.r.t.  $t$ , we get

$$\frac{dS}{dt} = 4\pi \times 2r \frac{dr}{dt}$$

$$\therefore \frac{dS}{dt} = 8\pi r \frac{dr}{dt} \quad \dots\dots(1)$$

$$\text{Now, } \frac{dr}{dt} = 0.02 \text{ cm/sec and } r = 5 \text{ cm}$$

$$\begin{aligned} \therefore (1) \text{ gives, } \frac{dS}{dt} &= 8\pi(5)(0.02) \\ &= 0.8\pi \end{aligned}$$

Hence, the surface area of the soap bubble is increasing at the rate of  $0.87c \text{ cm}^2 / \text{sec}$ .

Question 9.

The surface area of a spherical balloon is increasing at the rate of  $2 \text{ cm}^2/\text{sec}$ . At what rate is the volume of the balloon increasing, when the radius of the balloon is  $6 \text{ cm}$ ?

Solution:

Let  $r$  be the radius,  $S$  be the surface area and  $V$  be the volume of the spherical balloon at any time  $t$ .

$$\text{Then } S = 4\pi r^2 \text{ and } V = \frac{4}{3}\pi r^3$$

Differentiating w.r.t.  $t$ , we get

$$\frac{dS}{dt} = 4\pi \times 2r \frac{dr}{dt} = 8\pi r \frac{dr}{dt} \quad \dots (1)$$

$$\text{and } \frac{dV}{dt} = \frac{4}{3}\pi \times 3r^2 \frac{dr}{dt} = 4\pi r^2 \frac{dr}{dt}$$

$$\text{From (1), } \frac{dr}{dt} = \frac{1}{8\pi r} \cdot \frac{dS}{dt}$$

$$\therefore \frac{dV}{dt} = 4\pi r^2 \times \frac{1}{8\pi r} \cdot \frac{dS}{dt}$$

$$\therefore \frac{dV}{dt} = \frac{r}{2} \cdot \frac{dS}{dt} \quad \dots (2)$$

$$\text{Now, } \frac{dS}{dt} = 2 \text{ cm}^2/\text{sec} \text{ and } r = 6 \text{ cm}$$

$$\therefore (2) \text{ gives, } \frac{dV}{dt} = \frac{6}{2} \times 2 = 6$$

Hence, the volume of the spherical balloon is increasing at the rate of  $6 \text{ cm}^3/\text{sec}$ .

Question 10.

If each side of an equilateral triangle increases at the rate of  $\sqrt{2} \text{ cm/sec}$ , find the rate of increase of its area when its side of length is 3 cm.

Solution:

If  $x \text{ cm}$  is the side of the equilateral triangle and  $A$  is its area, then  $A = \frac{\sqrt{3}}{4} x^2$

Differentiating w.r.t.  $t$ , we get

$$\frac{dA}{dt} = \frac{\sqrt{3}}{4} \times 2x \frac{dx}{dt} = \frac{\sqrt{3}}{2} \cdot x \frac{dx}{dt} \quad \dots (1)$$

Now,  $\frac{dx}{dt} = \sqrt{2}$  cm/sec and  $x = 3$  cm

$$\begin{aligned} \therefore (1) \text{ gives, } \frac{dA}{dt} &= \frac{\sqrt{3}}{2} \times 3 \times \sqrt{2} \\ &= \frac{3\sqrt{6}}{2} \text{ cm}^2/\text{sec} \end{aligned}$$

Hence, rate of increase of the area of equilateral triangle =  $\frac{3\sqrt{6}}{2}$  cm<sup>2</sup>/sec.

Question 11.

The volume of a sphere increases at the rate of 20 cm<sup>3</sup>/sec. Find the rate of change of its surface area, when its radius is 5 cm.

Solution:

Let  $r$  be the radius,  $S$  be the surface area and  $V$  be the volume of the sphere at any time  $t$ .

Then  $S = 4\pi r^2$  and  $V = \frac{4}{3}\pi r^3$

Differentiating w.r.t.  $t$ , we get

$$\frac{dS}{dt} = 4\pi \times 2r \frac{dr}{dt} = 8\pi r \frac{dr}{dt}$$

$$\text{and } \frac{dV}{dt} = \frac{4\pi}{3} \times 3r^2 \frac{dr}{dt} = 4\pi r^2 \frac{dr}{dt} \quad \dots (1)$$

From (1),  $\frac{dr}{dt} = \frac{1}{4\pi r^2} \frac{dV}{dt}$

$$\therefore \frac{dS}{dt} = 8\pi r \times \frac{1}{4\pi r^2} \frac{dV}{dt}$$

$$\therefore \frac{dS}{dt} = \frac{2}{r} \cdot \frac{dV}{dt} \quad \dots (2)$$

Now,  $\frac{dV}{dt} = 20 \text{ cm}^3/\text{sec}$  and  $r = 5 \text{ cm}$

$$\therefore (2) \text{ gives, } \frac{dS}{dt} = \frac{2}{5} \times 20 = 8$$

Hence, the surface area of the sphere is changing at the rate of  $8 \text{ cm}^2/\text{sec}$ .

Question 12.

The edge of a cube is decreasing at the rate of  $0.6 \text{ cm}/\text{sec}$ . Find the rate at which its volume is decreasing, when the edge of the cube is  $2 \text{ cm}$ .

Solution:

Let  $x$  be the edge of the cube and  $V$  be its volume at any time  $t$ .

$$\text{Then } V = x^3$$

Differentiating both sides w.r.t.  $t$ , we get

$$\frac{dV}{dt} = 3x^2 \frac{dx}{dt}$$

Now,  $\frac{dx}{dt} = 0.6 \text{ cm}/\text{sec}$  and  $x = 2 \text{ cm}$

$$\begin{aligned}\therefore \frac{dV}{dt} &= 3(2)^2(0.6) \\ &= 7.2\end{aligned}$$


Hence, the volume of the cube is decreasing at the rate of  $7.2 \text{ cm}^3/\text{sec}$ .

Question 13.

A man of height 2 meters walks at a uniform speed of 6 km/hr away from a lamp post of 6 meters high. Find the rate at which the length of the shadow is increasing.

Solution:

Let OA be the lamp post, MN the man, MB = x, his shadow, and OM = y, the distance of the man from the lamp post at time t.



Then  $\frac{dy}{dt} = 6 \text{ km/hr}$  is the rate at which the man is moving away from the lamp post.

$\frac{dx}{dt}$  is the rate at which his shadow is increasing.

From the figure,

$$\begin{aligned}\frac{x}{2} &= \frac{x+y}{6} \\ 6x &= 2x + 2y\end{aligned}$$

$$4x = 2y$$

$$x = \frac{1}{2} y$$

$$\frac{dx}{dt} = \frac{1}{2} \frac{dy}{dt} = \frac{1}{2} \times 6 = 3 \text{ km/hr}$$

Hence, the length of the shadow is increasing at the rate of 3 km/hr.

Question 14.


A man of height 1.5 meters walks towards a lamp post of height 4.5 meters, at the rate of  $\left(\frac{3}{4}\right)$  meter/sec.

Find the rate at which

(i) his shadow is shortening

(ii) the tip of the shadow is moving.

Solution:



Let OA be the lamp post, MN the man, MB = x his shadow and OM = y the distance of the man from lamp post at time t.

Then  $\frac{dy}{dt} = \frac{3}{4}$  is the rate at which the man is moving towards the lamp post.

$\frac{dx}{dt}$  is the rate at which his shadow is shortening.

B is the tip of the shadow and it is at a distance of  $x + y$  from the post.

$\frac{d}{dt}(x + y) = \frac{dx}{dt} + \frac{dy}{dt}$  is the rate at which the tip of the shadow is moving.

From the figure,

$$\frac{x}{1.5} = \frac{x+y}{4.5}$$

$$45x = 15x + 15y$$

$$30x = 15y$$

$$x = \frac{1}{2}y$$

$$\frac{dx}{dt} = \frac{1}{2} \cdot \frac{dy}{dt} = \frac{1}{2} \left( \frac{3}{4} \right) = \left( \frac{3}{8} \right) \text{ metre/sec}$$

$$\text{and } \frac{dx}{dt} + \frac{dy}{dt} = \frac{3}{8} + \frac{3}{4} = \left( \frac{9}{8} \right) \text{ metres /sec}$$


Hence (i) the shadow is shortening at the rate of  $\left( \frac{3}{8} \right)$  metre/sec, and

(ii) the tip of shadow is moving at the rate of  $\left( \frac{9}{8} \right)$  metres/sec.

Question 15.

A ladder 10 metres long is leaning against a vertical wall. If the bottom of the ladder is pulled horizontally away from the wall at the rate of 1.2 metres per second, find how fast the top of the ladder is sliding down the wall, when the bottom is 6 metres away from the wall.

Solution:



Let AB be the ladder, where  $AB = 10$  metres.

Let at time  $t$  seconds, the end A of the ladder be  $x$  metres from the wall and the end B be  $y$  metres from the ground.

Since, OAB is a right angled triangle, by Pythagoras' theorem

$$x^2 + y^2 = 10^2 \text{ i.e. } y^2 = 100 - x^2$$

Differentiating w.r.t.  $t$ , we get

$$2y \frac{dy}{dt} = 0 - 2x \frac{dx}{dt}$$

$$\therefore \frac{dy}{dt} = -\frac{x}{y} \cdot \frac{dx}{dt} \dots\dots(1)$$

Now,  $\frac{dx}{dt} = 1.2$  metres/sec is the rate at which the bottom at of the ladder is pulled horizontally and  $\frac{dy}{dt}$  is the rate at which the top of ladder B is sliding.

$$\text{When } x = 6, y^2 = 100 - 36 = 64$$

$$y = 8$$


$$\begin{aligned} (1) \text{ gives } \frac{dy}{dt} &= -\frac{6}{8}(1.2) = -\frac{6}{8} \times \frac{12}{10} \\ &= -\frac{9}{10} = -0.9 \end{aligned}$$

Hence, the top of the ladder is sliding down the wall, at the rate of 0.9 metre/sec.

Question 16.

If water is poured into an inverted hollow cone whose semi-vertical angle is  $30^\circ$  so that its depth (measured along the axis) increases at the rate of 1 cm/sec. Find the rate at which the volume of water increases when the depth is 2 cm.

Solution:



Let  $r$  be the radius,  $h$  be the height,  $\theta$  be the semi-vertical angle and  $V$  be the volume of the water at any time  $t$ .

$$\text{Given : } \frac{dh}{dt} = 1 \text{ cm/sec, } \theta = 30^\circ$$

$$\text{Now, } V = \frac{1}{3} \pi r^2 h$$

---

But,  $\tan 30^\circ = \frac{r}{h}$

$$\therefore \frac{1}{\sqrt{3}} = \frac{r}{h} \quad \therefore r = \frac{h}{\sqrt{3}}$$

$$\therefore V = \frac{1}{3} \pi \left( \frac{h}{\sqrt{3}} \right)^2 h = \frac{\pi}{9} h^3$$

Differentiating w.r.t.  $t$ , we get,

$$\frac{dV}{dt} = \frac{\pi}{9} \times 3h^2 \frac{dh}{dt} = \frac{\pi}{3} h^2 \frac{dh}{dt}$$

When  $h = 2$  cm, then

$$\frac{dV}{dt} = \frac{\pi}{3} \times (2)^2 \times 1 = \frac{4\pi}{3}$$

Hence, the volume of water is increasing at the rate of  $\left(\frac{4\pi}{3}\right)$  cm<sup>3</sup>/sec.

## Ex 2.2

### Question 1.

Find the approximate value of given functions, at required points.

(i)  $\sqrt{8.95}$

Solution:

$$\text{Let } f(x) = \sqrt{x}. \text{ Then } f'(x) = \frac{1}{2\sqrt{x}}.$$

Take  $a = 9$  and  $h = -0.05$ . Then  $f(a) = f(9) = \sqrt{9} = 3$  and

$$f'(a) = f'(9) = \frac{1}{2\sqrt{9}} = \frac{1}{6}.$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore \sqrt{8.95} = f(9 - 0.05)$$

$$\doteq f(9) - (0.05)f'(9)$$

$$\doteq 3 - 0.05 \times \frac{1}{6}$$

$$\doteq 3 - 0.0083 = 2.9917$$

$$\sqrt{8.95} = 2.9917$$

(ii)  $\sqrt[3]{28}$

Solution:

$$\text{Let } f(x) = \sqrt[3]{x}$$

$$\text{Then } f'(x) = \frac{d}{dx}(x^{\frac{1}{3}}) = \frac{1}{3}x^{-\frac{2}{3}} = \frac{1}{3x^{\frac{2}{3}}}$$

Take  $a = 27$  and  $h = 1$ .

$$\text{Then } f(a) = f(27) = \sqrt[3]{27} = 3$$

$$\text{and } f'(a) = f'(27) = \frac{1}{3(27)^{\frac{2}{3}}} = \frac{1}{3 \times 9} = \frac{1}{27} = 0.03704$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore \sqrt[3]{28} = f(28) = f(27+1)$$

$$\doteq f(27) + 1 \cdot f'(27)$$

$$\doteq 3 + 1 \times 0.03704 = 3.03704$$

$$\therefore \sqrt[3]{28} \doteq 3.03704.$$

(iii)  $\sqrt[5]{31.98}$

Solution:

$$\text{Let } f(x) = \sqrt[5]{x}$$

$$\text{Then } f'(x) = \frac{d}{dx}(x^{\frac{1}{5}}) = \frac{1}{5}x^{-\frac{4}{5}} = \frac{1}{5x^{\frac{4}{5}}}$$

Take  $a = 32$  and  $h = -0.02$ .

Then  $f(a) = f(32) = \sqrt[5]{32} = 2$

$$f'(a) = f'(32) = \frac{1}{5(32)^{\frac{4}{5}}} = \frac{1}{5 \times 16} = \frac{1}{80} = 0.0125$$

The formula for approximation is

$$f(a + h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned} \therefore \sqrt[5]{31.98} &= f(31.98) = f(32 - 0.02) \\ &\doteq f(32) - 0.02 \cdot f'(32) \\ &\doteq 2 - 0.02 \times 0.0125 \\ &\doteq 2 - 0.000250 = 1.99975 \\ \therefore \sqrt[5]{31.98} &\doteq 1.99975. \end{aligned}$$

(iv)  $(3.97)^4$

Solution:

$$\text{Let } f(x) = x^4$$

$$\text{Then } f'(x) = \frac{d}{dx}(x^4) = 4x^3$$

Take  $a = 4$  and  $h = -0.03$ .

Then  $f(a) = f(4) = (4)^4 = 256$  and

$$f'(a) = f'(4) = 4(4)^3 = 256$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned}\therefore (3.97)^4 &= f(3.97) = f(4 - 0.03) \\ &\doteq f(4) - (0.03) f'(4) \\ &\doteq 256 - 0.03 \times 256 \\ &\doteq 256 - 7.68 = 248.32 \quad \therefore (3.97)^4 \doteq 248.32.\end{aligned}$$

(v)  $(4.01)^3$

Solution:

Let  $f(x) = x^3$ . Then,  $f'(x) = 3x^2$

Take  $a = 4$  and  $h = 0.01$ . Then

$$f(a) = f(4) = 4^3 = 64 \text{ and}$$

$$f'(a) = f'(4) = 3 \times 4^2 = 48.$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned}\therefore (4.01)^3 &= f(4 + 0.01) \\ &\doteq f(4) + (0.01) f'(4) \\ &\doteq 64 + 0.01 \times 48 \\ &\doteq 64 + 0.48 = 64.48 \quad \therefore (4.01)^3 \doteq 64.48.\end{aligned}$$

Question 2.

Find the approximate values of:

(i)  $\sin 61^\circ$ , given that  $1^\circ = 0.0174^c$ ,  $\sqrt{3} = 1.732$ .

Solution:

$$\text{Let } f(x) = \sin x$$

$$\text{Then } f'(x) = \frac{d}{dx}(\sin x) = \cos x$$

$$\text{Take } a = 60^\circ = \frac{\pi}{3} \text{ and } h = 1^\circ = 0.0174^c$$

$$\text{Then } f(a) = f\left(\frac{\pi}{3}\right) = \sin \frac{\pi}{3} = \frac{\sqrt{3}}{2} = \frac{1.732}{2} = 0.866$$

$$\text{and } f'(a) = f'\left(\frac{\pi}{3}\right) = \cos \frac{\pi}{3} = \frac{1}{2} = 0.5$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned} \therefore \sin 61^\circ &\doteq f(61^\circ) = f\left(\frac{\pi}{3} + 0.0174\right) \\ &\doteq f\left(\frac{\pi}{3}\right) + 0.0174 \cdot f'\left(\frac{\pi}{3}\right) \\ &\doteq 0.866 + 0.0174 \times 0.5 \\ &\doteq 0.866 + 0.00870 = 0.8747 \\ \therefore \sin 61^\circ &\doteq 0.8747. \end{aligned}$$

(ii)  $\sin(29^\circ 30')$ , given that  $1^\circ = 0.0175^c$ ,  $\sqrt{3} = 1.732$ .

Solution:

$$\text{Let } f(x) = \sin x$$

$$\text{Then } f'(x) = \frac{d}{dx}(\sin x) = \cos x$$

$$\text{Take } a = 30^\circ = \frac{\pi}{6} \text{ and}$$

$$h = -30' = -\left(\frac{1}{2}\right)^\circ = -\frac{1}{2} \times 0.0175 = -0.00875$$

$$\text{Then } f(a) = f\left(\frac{\pi}{6}\right) = \sin \frac{\pi}{6} = \frac{1}{2} = 0.5$$

$$\text{and } f'(a) = f'\left(\frac{\pi}{6}\right) = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} = \frac{1.732}{2} = 0.866$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore \sin(29^\circ 30') = f(29^\circ 30') = f\left(\frac{\pi}{6} + 0.00875\right)$$

$$\doteq f\left(\frac{\pi}{6}\right) + (0.00875) \cdot f'\left(\frac{\pi}{6}\right)$$

$$\doteq 0.5 + 0.00875 \times 0.866$$

$$\doteq 0.5 + 0.0075775 = 0.4924$$

$$\therefore \sin(29^\circ 30') \doteq 0.4924.$$

(iii)  $\cos(60^\circ 30')$ , given that  $1^\circ = 0.0175^c$ ,  $\sqrt{3} = 1.732$ .

Solution:

$$\text{Let } f(x) = \cos x$$

$$\text{Then } f'(x) = \frac{d}{dx}(\cos x) = -\sin x$$

$$\text{Take } a = 60^\circ = \frac{\pi}{3} \text{ and}$$

$$h = 30' = \left(\frac{1}{2}\right)^\circ = \left(\frac{1}{2} \times 0.0175\right)^c = 0.00875^c$$

$$\text{Then } f(a) = f\left(\frac{\pi}{3}\right) = \cos \frac{\pi}{3} = \frac{1}{2} = 0.5$$

$$f'(a) = f'\left(\frac{\pi}{3}\right) = -\sin \frac{\pi}{3} = -\frac{\sqrt{3}}{2} = -\frac{1.732}{2} = -0.866$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore \cos(60^\circ 30') = f(60^\circ 30') = f\left(\frac{\pi}{3} + 0.00875\right)$$

$$\doteq f\left(\frac{\pi}{3}\right) + 0.00875 \cdot f'\left(\frac{\pi}{3}\right)$$

$$\doteq 0.5 + (0.00875)(-0.8660)$$

$$\doteq 0.5 - 0.0075775 = 0.4924225$$

$$\therefore \cos(60^\circ 30') \doteq 0.4924.$$

(iv)  $\tan(45^\circ 40')$ , given that  $1^\circ = 0.0175^c$ .

Solution:

$$\text{Let } f(x) = \tan x$$

$$\text{Then } f'(x) = \frac{d}{dx}(\tan x) = \sec^2 x$$

$$\text{Take } a = 45^\circ = \frac{\pi}{4} \text{ and } h = 40' = \left(\frac{40}{60} \times 0.0175\right)^c = 0.01167^c$$

$$\text{Then } f(a) = f\left(\frac{\pi}{4}\right) = \tan \frac{\pi}{4} = 1$$

$$\text{and } f'(a) = f'\left(\frac{\pi}{4}\right) = \sec^2 \frac{\pi}{4} = (\sqrt{2})^2 = 2$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore \tan(45^\circ 40') = f(45^\circ 40') = f\left(\frac{\pi}{4} + 0.01167\right)$$

$$\doteq f\left(\frac{\pi}{4}\right) + (0.01167) \cdot f'\left(\frac{\pi}{4}\right)$$

$$\doteq 1 + 0.01167 \times 2 = 1 + 0.02334 = 1.02334$$

$$\therefore \tan(45^\circ 40') \doteq 1.02334.$$

Question 3.

---

Find the approximate values of

(i)  $\tan^{-1}(0.999)$ .

Solution:

$$\text{Let } f(x) = \tan^{-1} x$$

$$\text{Then } f'(x) = \frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$$

$$\text{Take } a = 1 \text{ and } h = -0.001$$

$$\text{Then } f(a) = f(1) = \tan^{-1} 1 = \frac{\pi}{4}$$

$$\text{and } f'(a) = f'(1) = \frac{1}{1+1^2} = \frac{1}{2}$$

The formula for approximation is

$$f(a+h) \doteq f(a) + hf'(a)$$

$$\therefore \tan^{-1}(0.999) \doteq f(0.999) = f(1 - 0.001)$$

$$\doteq f(1) - (0.001) \cdot f'(1)$$

$$\doteq \frac{\pi}{4} - 0.001 \times \frac{1}{2} = \frac{\pi}{4} - 0.0005$$

$$\therefore \tan^{-1}(0.999) \doteq \frac{\pi}{4} - 0.0005.$$

Remark : The answer can also be given as :

$$\begin{aligned}\tan^{-1}(0.999) &\doteq \frac{3.1416}{4} - 0.0005 \\ &\doteq 0.7854 - 0.0005 = 0.7849.\end{aligned}$$

(ii)  $\cot^{-1}(0.999)$ .

Solution:

$$\text{Let } f(x) = \cot^{-1} x$$

$$\therefore f'(x) = \frac{d}{dx}(\cot^{-1} x) = \frac{-1}{1+x^2}$$

Take  $a = 1$  and  $h = -0.001$

$$\text{Then } f(a) = f(1) = \cot^{-1} 1 = \frac{\pi}{4}$$

$$\text{and } f'(a) = f'(1) = \frac{-1}{1+1^2} = \frac{-1}{2}$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned}\therefore \cot^{-1}(0.999) &= f(0.999) = f(1 - 0.001) \\ &\doteq f(1) - (0.001) \cdot f'(1) \\ &\doteq \frac{\pi}{4} - (0.001) \cdot \left(\frac{-1}{2}\right) = \frac{\pi}{4} + 0.0005\end{aligned}$$

$$\therefore \cot^{-1}(0.999) \doteq \frac{\pi}{4} + 0.0005.$$

$$\begin{aligned}\cot^{-1}(0.999) &\doteq \frac{3.1416}{4} + 0.0005 \\ &\doteq 0.7854 + 0.0005 = 0.7859.\end{aligned}$$

(iii)  $\tan^{-1}(1.001)$ .

Solution:

$$\text{Let } f(x) = \tan^{-1} x$$

$$\therefore f'(x) = \frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$$

Take  $a = 1$  and  $h = 0.001$

$$\text{Then } f(a) = f(1) = \tan^{-1} 1 = \frac{\pi}{4}$$

$$\text{and } f'(a) = f'(1) = \frac{1}{1+1^2} = \frac{1}{2}$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned}\therefore \tan^{-1}(1.001) &= f(1.001) = f(1 + 0.001) \\ &\doteq f(1) + (0.001) \cdot f'(1) \\ &\doteq \frac{\pi}{4} + (0.001) \times \frac{1}{2} = \frac{\pi}{4} + 0.0005\end{aligned}$$

$$\therefore \tan^{-1}(1.001) \doteq \frac{\pi}{4} + 0.0005.$$

**Remark :** The answer can also be given as :

$$\begin{aligned}\tan^{-1}(1.001) &\doteq f(1) + (0.001) \cdot f'(1) \\ &\doteq \frac{\pi}{4} + (0.001) \times \frac{1}{2} \\ &\doteq \frac{3.1416}{4} + 0.0005 \\ &\doteq 0.7854 + 0.0005 = 0.7859.\end{aligned}$$

Question 4.

Find the approximate values of:

(i)  $e^{0.995}$ , given that  $e = 2.7183$ .

Solution:

$$\text{Let } f(x) = e^x.$$

$$\text{Then } f'(x) = \frac{d}{dx}(e^x) = e^x$$

Take  $a = 1$  and  $h = -0.005$ .

Then  $f(a) = f(1) = e = 2.7183$

and  $f'(a) = f'(1) = e = 2.7183$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore e^{0.995} = f(0.995) = f(1 - 0.005)$$

$$\doteq f(1) - (0.005) \cdot f'(1)$$

$$\doteq 2.7183 - 0.005 \times 2.7183$$

$$\doteq 2.7183 - 0.01359 = 2.70471$$

$$\therefore e^{0.995} \doteq 2.70471$$

(ii)  $e^{2.1}$ , given that  $e^2 = 7.389$ .

Solution:

$$\text{Let } f(x) = e^x$$

$$\text{Then } f'(x) = \frac{d}{dx}(e^x) = e^x$$

$$\text{Take } a = 2 \text{ and } h = 0.1$$

$$\text{Then } f(a) = f(2) = e^2 = 7.389$$

$$f'(a) = f'(2) = e^2 = 7.389$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore e^{2.1} = f(2.1) = f(2+0.1)$$

$$\doteq f(2) + (0.1) \cdot f'(2)$$

$$\doteq 7.389 + 0.1 \times 7.389$$

$$\doteq 7.389 + 0.7389 = 8.1279$$

$$\therefore e^{2.1} \doteq 8.1279.$$

(iii)  $3^{2.01}$ , given that  $\log 3 = 1.0986$ .

Solution:

$$\text{Let } f(x) = 3^x$$

$$\text{Then } f'(x) = \frac{d}{dx}(3^x) = 3^x \cdot \log 3$$

Take  $a = 2$  and  $h = 0.01$

$$\text{Then } f(a) = f(2) = 3^2 = 9$$

$$\text{and } f'(a) = f'(2) = 3^2 \cdot \log 3 = 9 \times 1.0986 = 9.8874$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore 3^{2.01} = f(2.01) = f(2 + 0.01)$$

$$\doteq f(2) + (0.01) \cdot f'(2)$$

$$\doteq 9 + 0.01 \times 9.8874$$

$$\doteq 9 + 0.098874 = 9.098874$$

$$\therefore 3^{2.01} \doteq 9.098874.$$

Question 5.

Find the approximate values of:

(i)  $\log_e(101)$ , given that  $\log_e 10 = 2.3026$ .

Solution:

$$\text{Let } f(x) = \log_e x. \text{ Then } f'(x) = \frac{1}{x}.$$

Take  $a = 100$  and  $h = 1$ . Then

$$\begin{aligned}f(a) &= f(100) = \log_e 100 = 2 \log_e 10 \\ &= 2 \times 2.3026 = 4.6052\end{aligned}$$

$$f'(a) = f'(100) = \frac{1}{100} = 0.01$$

The formula for approximation is

$$f(a+h) \doteq f(a) + h \cdot f'(a)$$

$$\therefore \log_e 101 = f(101) = f(100+1)$$

$$\doteq f(100) + 1 \cdot f'(100)$$

$$\doteq 4.6052 + 1 \times 0.01 = 4.6152$$

$$\log_e(101) \doteq 4.6152.$$

(ii)  $\log_e(9.01)$ , given that  $\log 3 = 1.0986$ .

Solution:

Let  $f(x) = \log_e x$ .

$$\text{Then } f'(x) = \frac{1}{x}$$

Take  $a = 100$  and  $h = 1$ . then

$$f(a) = f(100)$$

$$= \log_e 100$$

$$\begin{aligned} &= 3 \log_e 10 \\ &= 3 \times 1.0986 \\ &= 3.2958 \\ f'(a) &= f'(100) \\ &= \frac{1}{100} \\ &= 0.01 \end{aligned}$$

The formula for a approximation is

$$\begin{aligned} f(a + h) &\doteq f(a) + h.f'(a) \\ \therefore \log_e 9.01 &= f(9.01) \\ &= f(100 + 1) \\ &\doteq f(100) + 1.f'(100) \\ &\doteq 3.2958 + 1 \times 0.01 \\ &= 2.1983 \\ \log_e(901) &\doteq 2.1983. \end{aligned}$$

(iii)  $\log_{10}(1016)$ , given that  $\log_{10}e = 0.4343$ .

Solution:

$$\begin{aligned} \text{Let } f(x) &= \log_{10}x = \frac{\log_e x}{\log_e 10} \\ &= (\log_{10}e)(\log x) = (0.4343) \log x \end{aligned}$$

---

$$= (\log_{10} e)(\log x) = (0.4343) \log x$$

$$\text{Then } f'(x) = (0.4343) \cdot \frac{d}{dx}(\log x) = \frac{0.4343}{x}$$

Take  $a = 1000$  and  $h = 16$ . Then

$$f(a) = f(1000) = \log_{10} 1000 = \log_{10} 10^3 = 3$$

$$f'(a) = f'(1000) = \frac{0.4343}{1000}$$

The formula for approximation is

$$f(a+h) \doteq f(a) + hf'(a)$$

$$\therefore \log_{10} 1016 = f(1016) = f(1000 + 16)$$

$$\doteq f(1000) + 16 \cdot f'(1000)$$

$$\doteq 3 + 16 \times \frac{0.4343}{1000}$$

$$\doteq 3 + 0.0069488 \doteq 3.006949$$

$$\therefore \log_{10} 1016 \doteq 3.006949.$$

Question 6.

Find the approximate values of:

(i)  $f(x) = x^3 - 3x + 5$  at  $x = 1.99$

Solution:

$$f(x) = x^3 - 3x + 5$$

---

$$\begin{aligned}\therefore f'(x) &= \frac{d}{dx}(x^3 - 3x + 5) \\ &= 3x^2 - 3 \times 1 + 0 = 3x^2 - 3\end{aligned}$$

Take  $a = 2$ ,  $h = -0.01$

$$\begin{aligned}\text{Then } f(a) = f(2) &= (2)^3 - 3(2) + 5 \\ &= 8 - 6 + 5 = 7\end{aligned}$$

$$f'(a) = f'(2) = 3(2)^2 - 3 = 12 - 3 = 9$$

The formula for approximation is

$$f(a + h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned}\therefore f(1.99) &= f(2 - 0.01) \doteq f(2) - (0.01) \cdot f'(2) \\ &\doteq 7 - 0.01 \times 9 = 7 - 0.09 = 6.91\end{aligned}$$

$$\therefore f(1.99) \doteq 6.91.$$

(ii)  $f(x) = x^3 + 5x^2 - 7x + 10$  at  $x = 1.12$

Solution:

$$f(x) = x^3 + 5x^2 - 7x + 10$$

$$\begin{aligned}\therefore f'(x) &= \frac{d}{dx}(x^3 + 5x^2 - 7x + 10) \\ &= 3x^2 + 5 \times 2x - 7 \times 1 + 0 = 3x^2 + 10x - 7\end{aligned}$$

Take  $a = 1$ ,  $h = 0.12$

$$\begin{aligned}\text{Then } f(a) = f(1) &= (1)^3 + 5(1)^2 - 7(1) + 10 \\ &= 1 + 5 - 7 + 10 = 9\end{aligned}$$

$$\text{and } f'(a) = f'(1) = 3(1)^2 + 10(1) - 7 = 3 + 10 - 7 = 6$$

The formula for approximation is

$$f(a + h) \doteq f(a) + h \cdot f'(a)$$

$$\begin{aligned}\therefore f(1.12) &= f(1 + 0.12) \doteq f(1) + (0.12) \cdot f'(1) \\ &\doteq 9 + 0.12 \times 6 \doteq 9 + 0.72 = 9.72\end{aligned}$$

$$\therefore f(1.12) \doteq 9.72.$$

### Ex 2.3

#### Question 1.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>

Check the validity of the Rolle's theorem for the following functions.

(i)  $f(x) = x^2 - 4x + 3, x \in [1, 3]$

**Solution:**

The function  $f$  given as  $f(x) = x^2 - 4x + 3$  is polynomial function.

Hence, it is continuous on  $[1, 3]$  and differentiable on  $(1, 3)$ .

$$\text{Now, } f(1) = 1^2 - 4(1) + 3 = 1 - 4 + 3 = 0$$

$$\text{and } f(3) = 3^2 - 4(3) + 3 = 9 - 12 + 3 = 0$$

$$\therefore f(1) = f(3)$$

Thus, the function  $f$  satisfies all the conditions of Rolle's theorem.

(ii)  $f(x) = e^{-x} \sin x, x \in [0, \pi]$ .

**Solution:**

The functions  $e^{-x}$  and  $\sin x$  are continuous and differentiable on their domains.

$$\therefore f(x) = e^{-x} \sin x \text{ is continuous on } [0, \pi] \text{ and differentiable on } (0, \pi).$$

$$\text{Now, } f(0) = e^0 \sin 0 = 1 \times 0 = 0$$

$$\text{and } f(\pi) = e^{-\pi} \sin \pi = e^{-\pi} \times 0 = 0$$

$$\therefore f(0) = f(\pi)$$

Thus, the function  $f$  satisfies all the conditions of the Rolle's theorem.

(iii)  $f(x) = 2x^2 - 5x + 3, x \in [1, 3]$ .

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>

**Solution:**

The function  $f$  given as  $f(x) = 2x^2 - 5x + 3$  is a polynomial function.

Hence, it is continuous on  $[1, 3]$  and differentiable on  $(1, 3)$ .

$$\text{Now, } f(1) = 2(1)^2 - 5(1) + 3 = 2 - 5 + 3 = 0$$

$$\text{and } f(3) = 2(3)^2 - 5(3) + 3 = 18 - 15 + 3 = 6$$

$$\therefore f(1) \neq f(3)$$

Hence, the conditions of Rolle's theorem are not satisfied.

**(iv)  $f(x) = \sin x - \cos x + 3$ ,  $x \in [0, 2\pi]$ .**

**Solution:**

The functions  $\sin x$ ,  $\cos x$  and  $3$  are continuous and differentiable on their domains.

$\therefore f(x) = \sin x - \cos x + 3$  is continuous on  $[0, 2\pi]$  and differentiable on  $(0, 2\pi)$ .

$$\text{Now, } f(0) = \sin 0 - \cos 0 + 3 = 0 - 1 + 3 = 2$$

$$\text{and } f(2\pi) = \sin 2\pi - \cos 2\pi + 3 = 0 - 1 + 3 = 2$$

$$\therefore f(0) = f(2\pi)$$

Thus, the function  $f$  satisfies all the conditions of the Rolle's theorem.

$$(v) f(x) = x^2, \text{ if } 0 \leq x \leq 2 \\ = 6 - x, \text{ if } 2 < x \leq 6.$$

Solution:

$$f(x) = x^2, \text{ if } 0 \leq x \leq 2 \\ = 6 - x, \text{ if } 2 < x \leq 6$$

$$\therefore f'(x) = \frac{d}{dx}(x^2) = 2x, \text{ if } 0 \leq x \leq 2$$

$$= \frac{d}{dx}(6 - x) = -1, \text{ if } 2 < x \leq 6$$

$$\therefore LF'(2) = 2(2) = 4 \text{ and } RF'(2) = -1$$

$$\therefore LF'(2) \neq RF'(2)$$

$\therefore f$  is not differentiable at  $x = 2$  and  $2 \in (0, 6)$ .

$\therefore f$  is not differentiable at all the points on  $(0, 6)$ .

Hence, the conditions of Rolle's theorem are not satisfied.

$$(vi) f(x) = x^{\frac{2}{3}}, x \in [-1, 1].$$

Solution:

$$f(x) = x^{\frac{2}{3}}$$

$$\therefore f'(x) = \frac{d}{dx}\left(x^{\frac{2}{3}}\right) = \frac{2}{3}x^{-\frac{1}{3}} = \frac{2}{3\sqrt[3]{x}}$$

This does not exist at  $x = 0$  and  $0 \in (-1, 1)$

$\therefore f$  is not differentiable on the interval  $(-1, 1)$ .

Hence, the conditions of Rolle's theorem are not satisfied.

Question 2.

Given an interval  $[a, b]$  that satisfies hypothesis of Rolle's theorem for the function  $f(x) = x^4 + x^2 - 2$ . It is known that  $a = -1$ . Find the value of  $b$ .

Solution:

$$f(x) = x^4 + x^2 - 2$$

Since the hypothesis of Rolle's theorem are satisfied by  $f$  in the interval  $[a, b]$ , we have

$$f(a) = f(b), \text{ where } a = -1$$

$$\text{Now, } f(a) = f(-1) = (-1)^4 + (-1)^2 - 2 = 1 + 1 - 2 = 0$$

$$\text{and } f(b) = b^4 + b^2 - 2$$

$$\therefore f(a) = f(b) \text{ gives}$$

$$0 = b^4 + b^2 - 2 \text{ i.e. } b^4 + b^2 - 2 = 0.$$

Since,  $b = 1$  satisfies this equation,  $b = 1$  is one of the roots of this equation.

Hence,  $b = 1$ .

Question 3.

Verify Rolle's theorem for the following functions.

(i)  $f(x) = \sin x + \cos x + 7, x \in [0, 2\pi]$

Solution:

The functions  $\sin x, \cos x$  and  $7$  are continuous and differentiable on their domains.

$\therefore f(x) = \sin x + \cos x + 7$  is continuous on  $[0, 2\pi]$  and differentiable on  $(0, 2\pi)$

$$\text{Now, } f(0) = \sin 0 + \cos 0 + 7 = 0 + 1 + 7 = 8$$

$$\text{and } f(2\pi) = \sin 2\pi + \cos 2\pi + 7 = 0 + 1 + 7 = 8$$

$$\therefore f(0) = f(2\pi)$$

Thus, the function  $f$  satisfies all the conditions of Rolle's theorem.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>

Thus, the function  $f$  satisfies all the conditions of Rolle's theorem.

$\therefore$  there exists  $c \in (0, 2\pi)$  such that  $f'(c) = 0$ .

Now,  $f(x) = \sin x + \cos x + 7$

$$\therefore f'(x) = \frac{d}{dx} (\sin x + \cos x + 7)$$

$$= \cos x - \sin x + 0$$

$$= \cos x - \sin x$$

$$\therefore f'(c) = \cos c - \sin c$$

$$\therefore f'(c) = 0 \text{ gives, } \cos c - \sin c = 0$$

$$\therefore \cos c = \sin c$$

$$\therefore c = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}, \dots$$

$$\text{But } \frac{\pi}{4}, \frac{5\pi}{4} \in (0, 2\pi)$$

$$\therefore c = \frac{\pi}{4} \text{ or } \frac{5\pi}{4}$$

Hence, the Rolle's theorem is verified.

(ii)  $f(x) = \sin\left(\frac{x}{2}\right)$ ,  $x \in [0, 2\pi]$

Solution:

The function  $f(x) = \sin\left(\frac{x}{2}\right)$  is continuous on  $[0, 2\pi]$  and differentiable on  $(0, 2\pi)$ .

$$\text{Now, } f(0) = \sin 0 = 0$$

$$\text{and } f(2\pi) = \sin \pi = 0$$

$$\therefore f(0) = f(2\pi)$$

Thus, the function  $f$  satisfies all the conditions of Rolle's theorem.

$\therefore$  there exists  $c \in (0, 2\pi)$  such that  $f'(c) = 0$ .

$$\text{Now, } f(x) = \sin\left(\frac{x}{2}\right)$$

$$\therefore f'(x) = \frac{d}{dx}\left[\sin\left(\frac{x}{2}\right)\right]$$

$$= \cos\left(\frac{x}{2}\right) \cdot \frac{d}{dx}\left(\frac{x}{2}\right)$$

$$= \cos\left(\frac{x}{2}\right) \cdot \frac{1}{2} = \frac{1}{2}\cos\left(\frac{x}{2}\right)$$

$$\therefore f'(c) = \frac{1}{2}\cos\left(\frac{c}{2}\right)$$

$$\therefore f'(c) = 0 \text{ gives } \frac{1}{2}\cos\left(\frac{c}{2}\right) = 0$$

$$\therefore \cos\left(\frac{c}{2}\right) = 0$$

$$\therefore \cos\frac{c}{2} = \cos\frac{\pi}{2} = \cos\frac{3\pi}{2} = \cos\frac{5\pi}{2} = \dots$$

$$\therefore \frac{c}{2} = \frac{\pi}{2}, \frac{3\pi}{2}, \frac{5\pi}{2}, \dots$$

$$\therefore c = \pi, 3\pi, 5\pi, \dots$$

But  $\pi \in (0, 2\pi)$

$$\therefore c = \pi$$

---

Hence, Rolle's theorem is verified.

(iii)  $f(x) = x^2 - 5x + 9, x \in [1, 4]$ .

Solution:

The function  $f$  given as  $f(x) = x^2 - 5x + 9$  is a polynomial function.

Hence it is continuous on  $[1, 4]$  and differentiable on  $(1, 4)$ .

$$\text{Now, } f(1) = 1^2 - 5(1) + 9 = 1 - 5 + 9 = 5$$

$$\text{and } f(4) = 4^2 - 5(4) + 9 = 16 - 20 + 9 = 5$$

$$\therefore f(1) = f(4)$$

Thus, the function  $f$  satisfies all the conditions of the Rolle's theorem.

$\therefore$  there exists  $c \in (1, 4)$  such that  $f'(c) = 0$ .

$$\text{Now, } f(x) = x^2 - 5x + 9$$

$$\therefore f'(x) = \frac{d}{dx} (x^2 - 5x + 9)$$

$$= 2x - 5 \times 1 + 0$$

$$= 2x - 5$$

$$\therefore f'(c) = 2c - 5$$

$$\therefore f'(c) = 0 \text{ gives, } 2c - 5 = 0$$

$$\therefore c = 5/2 \in (1, 4)$$

Hence, the Rolle's theorem is verified.

Question 4.

If Rolle's theorem holds for the function  $f(x) = x^3 + px^2 + qx + 5, x \in [1, 3]$  with  $c = 2 + \frac{1}{\sqrt{3}}$ , find the values of  $p$  and  $q$ .

Solution:

The Rolle's theorem holds for the function  $f(x) = x^3 + px^2 + qx + 5$ ,  $x \in [1, 3]$

$$\therefore f(1) = f(3)$$

$$\therefore 1^3 + p(1)^2 + q(1) + 5 = 3^3 + p(3)^2 + q(3) + 5$$

$$\therefore 1 + p + q + 5 = 27 + 9p + 3q + 5$$

$$\therefore 8p + 2q = -26$$

$$\therefore 4p + q = -13 \dots\dots (1)$$

Also, there exists at least one point  $c \in (1, 3)$  such that  $f'(c) = 0$ .

$$\text{Now, } f'(x) = \frac{d}{dx} (x^3 + px^2 + qx + 5)$$

$$= 3x^2 + p \times 2x + q \times 1 + 0$$

$$= 3x^2 + 2px + q$$

$$\therefore f'(c) = 3c^2 + 2pc + q, \text{ where } c = 2 + \frac{1}{\sqrt{3}}$$

$$\therefore f'(c) = 3\left(2 + \frac{1}{\sqrt{3}}\right)^2 + 2p\left(2 + \frac{1}{\sqrt{3}}\right) + q$$

$$= 3\left(4 + \frac{4}{\sqrt{3}} + \frac{1}{3}\right) + 4p + \frac{2p}{\sqrt{3}} + q$$

$$= 12 + \frac{12}{\sqrt{3}} + 1 + 4p + \frac{2p}{\sqrt{3}} + q$$

$$= 4p + \frac{2p}{\sqrt{3}} + q + 13 + \frac{12}{\sqrt{3}}$$

But  $f'(c) = 0$

$$\therefore 4p + \frac{2p}{\sqrt{3}} + q + 13 + \frac{12}{\sqrt{3}} = 0$$

$$\therefore (4\sqrt{3} + 2)p + \sqrt{3}q + (13\sqrt{3} + 12) = 0$$

$$\therefore (4\sqrt{3} + 2)p + \sqrt{3}q = -13\sqrt{3} - 12 \dots\dots\dots (2)$$

Multiplying equation (1) by  $\sqrt{3}$ , we get

$$4\sqrt{3}p + \sqrt{3}q = -13\sqrt{3}$$

Subtracting this equation from (2), we get

$$2p = -12 \Rightarrow p = -6$$

$$\therefore \text{from (1), } 4(-6) + q = -13 \Rightarrow q = 11$$

Hence,  $p = -6$  and  $q = 11$ .

Question 5.

If Rolle's theorem holds for the function  $f(x) = (x - 2) \log x$ ,  $x \in [1, 2]$ , show that the equation  $x \log x = 2 - x$  is satisfied by at least one value of  $x$  in  $(1, 2)$ .

Solution:

The Rolle's theorem holds for the function  $f(x) = (x - 2) \log x$ ,  $x \in [1, 2]$ .

$\therefore$  there exists at least one real number  $c \in (1, 2)$  such that  $f'(c) = 0$ .

Now,  $f(x) = (x - 2) \log x$

$$\therefore f'(x) = \frac{d}{dx} [(x - 2) \log x]$$

$$= (x - 2) \cdot \frac{d}{dx} (\log x) + \log x \cdot \frac{d}{dx} (x - 2)$$

---

$$= (x-2) \times \frac{1}{x} + (\log x)(1-0)$$

$$= 1 - \frac{2}{x} + \log x$$

$$\therefore f'(c) = 1 - \frac{2}{c} + \log c$$

$$\therefore f'(c) = 0 \text{ gives } 1 - \frac{2}{c} + \log c = 0$$

$$\therefore c - 2 + c \log c = 0$$

$$\therefore c \log c = 2 - c, \text{ where } c \in (1, 2)$$

$$\therefore c \text{ satisfies the equation } x \log x = 2 - x, c \in (1, 2).$$

Hence, the equation  $x \log x = 2 - x$  is satisfied by at least one value of  $x$  in  $(1, 2)$ .

Question 6.

The function  $f(x) = x(x+3)e^{-\frac{x}{2}}$  satisfies all the conditions of Rolle's theorem on  $[-3, 0]$ . Find the value of  $c$  such that  $f'(c) = 0$ .

Solution:

The function  $f(x)$  satisfies all the conditions of Rolle's theorem, therefore there exist  $c \in (-3, 0)$  such that  $f'(c) = 0$ .

$$\text{Now, } f(x) = x(x+3)e^{-\frac{x}{2}} = (x^2 + 3x)e^{-\frac{x}{2}}$$

$$\therefore f'(x) = \frac{d}{dx} [(x^2 + 3x)e^{-\frac{x}{2}}]$$

$$\begin{aligned} &= (x^2 + 3x) \cdot \frac{d}{dx} (e^{-\frac{x}{2}}) + e^{-\frac{x}{2}} \cdot \frac{d}{dx} (x^2 + 3x) \\ &= (x^2 + 3x) \cdot e^{-\frac{x}{2}} \cdot \frac{d}{dx} \left( -\frac{x}{2} \right) + e^{-\frac{x}{2}} \times (2x + 3 \times 1) \\ &= (x^2 + 3x) \cdot e^{-\frac{x}{2}} \times -\frac{1}{2} + e^{-\frac{x}{2}} (2x + 3) \\ &= e^{-\frac{x}{2}} \left[ (2x + 3) - \frac{x^2 + 3x}{2} \right] \\ &= e^{-\frac{x}{2}} \left[ \frac{4x + 6 - x^2 - 3x}{2} \right] \\ &= \frac{e^{-\frac{x}{2}}}{2} (6 + x - x^2) \\ &= \frac{e^{-\frac{x}{2}}}{2} (3 - x)(2 + x) \end{aligned}$$

$$\therefore f'(c) = \frac{e^{-\frac{c}{2}}}{2} (3 - c)(2 + c)$$

$$\therefore f'(c) = 0 \text{ gives } \frac{e^{-\frac{c}{2}}}{2} (3 - c)(2 + c) = 0$$

$$\therefore (3 - c)(2 + c) = 0 \quad \dots \left[ \because \frac{e^{-\frac{c}{2}}}{2} \neq 0 \right]$$

$$\therefore (3 - c) = 0 \text{ or } (2 + c) = 0$$

$$\therefore c = 3 \text{ or } c = -2$$

$$\text{But } 3 \notin (-3, 0) \quad \therefore c \neq 3$$

$$\text{Hence, } c = -2.$$

Question 7.

Verify Lagrange's mean value theorem for the following functions:

(i)  $f(x) = \log x$  on  $[1, e]$ .

Solution:

The function  $f$  given as  $f(x) = \log x$  is a logarithmic function that is continuous for all positive real numbers.

Hence, it is continuous on  $[1, e]$  and differentiable on  $(1, e)$ .

Thus, the function  $f$  satisfies the conditions of Lagrange's mean value theorem.

$\therefore$  there exists  $c \in (1, e)$  such that

$$f'(c) = \frac{f(e) - f(1)}{e - 1} \quad \text{-----(1)}$$

$$\text{Now, } f(x) = \log x$$

$$\therefore f(1) = \log 1 = 0 \text{ and } f(e) = \log e = 1$$

$$\text{Also, } f'(x) = \frac{d}{dx}(\log x) = \frac{1}{x}$$

$$\therefore f'(c) = \frac{1}{c}$$

$$\therefore \text{ from (1), } \frac{1}{c} = \frac{1-0}{e-1} = \frac{1}{e-1}$$

$$\therefore c = e - 1 \in (1, e)$$

Hence, Lagrange's mean value theorem is verified.

(ii)  $f(x) = (x - 1)(x - 2)(x - 3)$  on  $[0, 4]$ .

Solution:

The function  $f$  given as

$$f(x) = (x - 1)(x - 2)(x - 3)$$

$$= (x - 1)(x^2 - 5x + 6)$$

$$= x^3 - 5x^2 + 6x - x^2 + 5x - 6$$

$$= x^3 - 6x^2 + 11x - 6 \text{ is a polynomial function.}$$

Hence, it is continuous on  $[0, 4]$  and differentiable on  $(0, 4)$ .

Thus, the function  $f$  satisfies the conditions of Lagrange's, mean value theorem.

$\therefore$  there exists  $c \in (0, 4)$  such that

$$f'(c) = \frac{f(4) - f(0)}{4 - 0} \quad \dots\dots(1)$$

$$\text{Now, } f(x) = (x - 1)(x - 2)(x - 3)$$

$$\therefore f(0) = (0 - 1)(0 - 2)(0 - 3) = (-1)(-2)(-3) = -6$$

$$\text{and } f(4) = (4 - 1)(4 - 2)(4 - 3) = (3)(2)(1) = 6$$

$$\text{Also, } f'(x) = \frac{d}{dx}(x^3 - 6x^2 + 11x - 6)$$

$$= 3x^2 - 6 \times 2x + 11 \times 1 - 0$$

$$= 3x^2 - 12x + 11$$

$$\therefore f'(c) = 3c^2 - 12c + 11$$

$$\therefore \text{from (1), } 3c^2 - 12c + 11 = \frac{6 - (-6)}{4}$$

$$\therefore 3c^2 - 12c + 11 = 3$$

$$\therefore 3c^2 - 12c + 8 = 0.$$

$$\therefore c = \frac{12 \pm \sqrt{144 - 4(3)(8)}}{2(3)}$$

$$\therefore c = \frac{12 \pm \sqrt{48}}{6} = \frac{12 \pm 4\sqrt{3}}{6}$$

$$\therefore c = 2 \pm \frac{2}{\sqrt{3}} \in (0, 4)$$

Hence, Lagrange's mean value theorem is verified.

$$(iii) f(x) = x^2 - 3x - 1, x \in \left[ \frac{-11}{7}, \frac{13}{7} \right]$$

Solution:

The function  $f$  given as  $f(x) = x^2 - 3x - 1$  is a polynomial function.

Hence, it is continuous on  $\left[ \frac{-11}{7}, \frac{13}{7} \right]$  and differentiable on  $\left( \frac{-11}{7}, \frac{13}{7} \right)$ .

Thus, the function  $f$  satisfies the conditions of LMVT.

---

$\therefore$  there exists  $c \in \left( \frac{-11}{7}, \frac{13}{7} \right)$  such that

$$f'(c) = \frac{f\left(\frac{13}{7}\right) - f\left(\frac{-11}{7}\right)}{\frac{13}{7} - \left(\frac{-11}{7}\right)} \quad \text{.....(1)}$$

$$\text{Now, } f(x) = x^2 - 3x - 1$$

$$\therefore f\left(\frac{-11}{7}\right) = \left(\frac{-11}{7}\right)^2 - 3\left(\frac{-11}{7}\right) - 1$$

$$= \frac{121}{49} + \frac{33}{7} - 1$$

$$= \frac{121 + 231 - 49}{49} = \frac{303}{49}$$

$$\text{and } f\left(\frac{13}{7}\right) = \left(\frac{13}{7}\right)^2 - 3\left(\frac{13}{7}\right) - 1 = \frac{169}{49} - \frac{39}{7} - 1$$

$$= \frac{169 - 273 - 49}{49} = \frac{-153}{49}$$

$$\text{Also, } f'(x) = \frac{d}{dx}(x^2 - 3x - 1) = 2x - 3 \times 1 - 0$$

$$= 2x - 3$$

$$\therefore f'(c) = 2c - 3$$

$$\therefore \text{ from (1), } 2c - 3 = \frac{\frac{-153}{49} - \frac{303}{49}}{\frac{13}{7} + \frac{11}{7}}$$

$$\therefore 2c - 3 = -\frac{456}{49} \times \frac{7}{24} = \frac{-57}{21}$$

$$\therefore 2c = \frac{-57}{21} + 3 = \frac{-57 + 63}{21} = \frac{6}{21} = \frac{2}{7}$$

$$\therefore c = \frac{1}{7} \in \left( \frac{-11}{7}, \frac{13}{7} \right)$$

Hence, Lagrange's mean value theorem is verified.

(iv)  $f(x) = 2x - x^2$ ,  $x \in [0, 1]$ .

Solution:

The function  $f$  given as  $f(x) = 2x - x^2$  is a polynomial function.

Hence, it is continuous on  $[0, 1]$  and differentiable on  $(0, 1)$ .

Thus, the function  $f$  satisfies the conditions of Lagrange's mean value theorem.

$\therefore$  there exists  $c \in (0, 1)$  such that

$$f'(c) = \frac{f(1) - f(0)}{1 - 0} \quad \text{.....(1)}$$

Now,  $f(x) = 2x - x^2$

$\therefore f(0) = 0 - 0 = 0$  and  $f(1) = 2(1) - 1^2 = 1$

Also,  $f'(x) = \frac{d}{dx}(2x - x^2) = 2 \times 1 - 2x$

$= 2 - 2x$

$\therefore f'(c) = 2 - 2c$

$\therefore$  from (1),  $2 - 2c = \frac{1-0}{1} = 1$

$\therefore 2c = 1 \quad \therefore c = \frac{1}{2} \in (0, 1)$

Hence, Lagrange's mean value theorem is verified.

(v)  $f(x) = \frac{x-1}{x-3}$  on  $[4, 5]$ .

Solution:

The function  $f$  given as

$f(x) = \frac{x-1}{x-3}$  is a rational function which is continuous except at  $x = 3$ .

But  $3 \notin [4, 5]$

Hence, it is continuous on  $[4, 5]$  and differentiable on  $(4, 5)$ .

Thus, the function  $f$  satisfies the conditions of Lagrange's mean value theorem.

$\therefore$  there exists  $c \in (4, 5)$  such that

$f'(c) = \frac{f(5) - f(4)}{5 - 4} \dots\dots\dots(1)$

$$\text{Now, } f(x) = \frac{x-1}{x-3}$$

$$\therefore f(4) = \frac{4-1}{4-3} = \frac{3}{1} = 3$$

$$\text{and } f(5) = \frac{5-1}{5-3} = \frac{4}{2} = 2$$

$$\text{Also, } f'(x) = \frac{d}{dx} \left( \frac{x-1}{x-3} \right)$$

$$= \frac{(x-3) \cdot \frac{d}{dx}(x-1) - (x-1) \cdot \frac{d}{dx}(x-3)}{(x-3)^2}$$

$$= \frac{(x-3) \times (1-0) - (x-1) \times (1-0)}{(x-3)^2}$$

$$= \frac{x-3-x+1}{(x-3)^2} = \frac{-2}{(x-3)^2}$$

$$\therefore f'(c) = \frac{-2}{(c-3)^2}$$

$$\therefore \text{from (1), } \frac{-2}{(c-3)^2} = \frac{2-3}{1} = -1$$

$$\therefore (c-3)^2 = 2$$

$$\therefore c-3 = \pm \sqrt{2} \quad \therefore c = 3 \pm \sqrt{2}$$

$$\text{But } (3 - \sqrt{2}) \notin (4, 5) \quad \therefore c \neq 3 - \sqrt{2}$$

$$\therefore c = 3 + \sqrt{2} \in (4, 5)$$

Hence, Lagrange's mean value theorem is verified.

#### Ex 2.4

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>

Question 1.

Test whether the following functions are increasing or decreasing.

(i)  $f(x) = x^3 - 6x^2 + 12x - 16$ ,  $x \in \mathbb{R}$ .

Solution:

$$f(x) = x^3 - 6x^2 + 12x - 16$$

$$\therefore f'(x) = \frac{d}{dx} (x^3 - 6x^2 + 12x - 16)$$

$$= 3x^2 - 6 \times 2x + 12 \times 1 - 0$$

$$= 3x^2 - 12x + 12$$

$$= 3(x^2 - 4x + 4)$$

$$= 3(x - 2)^2 \geq 0 \text{ for all } x \in \mathbb{R}$$

$$\therefore f'(x) \geq 0 \text{ for all } x \in \mathbb{R}$$

$$\therefore f \text{ is increasing for all } x \in \mathbb{R}.$$

(ii)  $f(x) = 2 - 3x + 3x^2 - x^3, x \in \mathbb{R}$ .

Solution:

$$f(x) = 2 - 3x + 3x^2 - x^3$$

$$\therefore f'(x) = \frac{d}{dx} (2 - 3x + 3x^2 - x^3)$$

$$= 0 - 3 \times 1 + 3 \times 2x - 3x^2$$

$$= -3 + 6x - 3x^2$$

$$= -3(x^2 - 2x + 1)$$

$$= -3(x - 1)^2 \leq 0 \text{ for all } x \in \mathbb{R}$$

$$\therefore f'(x) \leq 0 \text{ for all } x \in \mathbb{R}$$

$\therefore f$  is decreasing for all  $x \in \mathbb{R}$ .

(iii)  $f(x) = x - \frac{1}{x}, x \in \mathbb{R}, x \neq 0$ .

Solution:

$$f(x) = x - \frac{1}{x}$$

$$f'(x) = \frac{d}{dx} \left( x - \frac{1}{x} \right) = 1 - \left( \frac{-1}{x^2} \right)$$

$$= 1 + \frac{1}{x^2} > 0 \text{ for all } x \in \mathbb{R}, x \neq 0$$

$$\therefore f'(x) > 0 \text{ for all } x \in \mathbb{R}, \text{ where } x \neq 0$$

$\therefore f$  is increasing for all  $x \in \mathbb{R}$ , where  $x \neq 0$ .

Question 2.

Find the values of  $x$  for which the following functions are strictly increasing:

(i)  $f(x) = 2x^3 - 3x^2 - 12x + 6$

Solution:

$$f(x) = 2x^3 - 3x^2 - 12x + 6$$

$$\therefore f'(x) = \frac{d}{dx} (2x^3 - 3x^2 - 12x + 6)$$

$$= 2 \times 3x^2 - 3 \times 2x - 12 \times 1 + 0$$

$$= 6x^2 - 6x - 12$$

$$= 6(x^2 - x - 2)$$

f is strictly increasing if  $f'(x) > 0$

$$\text{i.e. if } 6(x^2 - x - 2) > 0$$

$$\text{i.e. if } x^2 - x - 2 > 0$$

$$\text{i.e. if } x^2 - x > 2$$

$$\text{i.e. if } x^2 - x + \frac{1}{4} > 2 + \frac{1}{4}$$

$$\text{i.e. if } \left(x - \frac{1}{2}\right)^2 > \frac{9}{4}$$

$$\text{i.e. if } x - \frac{1}{2} > \frac{3}{2} \text{ or } x - \frac{1}{2} < \frac{-3}{2} \text{ i.e. if } x > 2 \text{ or } x < -1$$

$\therefore$  f is strictly increasing if  $x < -1$  or  $x > 2$ .

$$\text{(ii) } f(x) = 3 + 3x - 3x^2 + x^3$$

Solution:

$$f(x) = 3 + 3x - 3x^2 + x^3$$

$$\therefore f'(x) = \frac{d}{dx} (3 + 3x - 3x^2 + x^3)$$

$$= 0 + 3 \times 1 - 3 \times 2x + 3x^2$$

$$= 3 - 6x + 3x^2$$

$$= 3(x^2 - 2x + 1)$$

f is strictly increasing if  $f'(x) > 0$

$$\text{i.e. if } 3(x^2 - 2x + 1) > 0$$

$$\text{i.e. if } x^2 - 2x + 1 > 0$$

$$\text{i.e. if } (x - 1)^2 > 0$$

This is possible if  $x \in \mathbb{R}$  and  $x \neq 1$

$$\text{i.e. } x \in \mathbb{R} - \{1\}$$

$\therefore f$  is strictly increasing if  $x \in \mathbb{R} - \{1\}$ .

(iii)  $f(x) = x^3 - 6x^2 - 36x + 7$

Solution:

$$f(x) = x^3 - 6x^2 - 36x + 7$$

$$\therefore f'(x) = \frac{d}{dx} (x^3 - 6x^2 - 36x + 7)$$

$$= 3x^2 - 6 \times 2x - 36 \times 1 + 0$$

$$= 3x^2 - 12x - 36$$

$$= 3(x^2 - 4x - 12)$$

$f$  is strictly increasing if  $f'(x) > 0$

i.e. if  $3(x^2 - 4x - 12) > 0$

i.e. if  $x^2 - 4x - 12 > 0$

i.e. if  $x^2 - 4x > 12$

i.e. if  $x^2 - 4x + 4 > 12 + 4$

i.e. if  $(x - 2)^2 > 16$

i.e. if  $x - 2 > 4$  or  $x - 2 < -4$  i.e. if  $x > 6$  or  $x < -2$

$\therefore f$  is strictly increasing if  $x < -2$  or  $x > 6$ .

Question 3.

Find the values of  $x$  for which the following functions are strictly decreasing:

(i)  $f(x) = 2x^3 - 3x^2 - 12x + 6$

Solution:

$$f(x) = 2x^3 - 3x^2 - 12x + 6$$

$$\therefore f'(x) = \frac{d}{dx} (2x^3 - 3x^2 - 12x + 6)$$

$$\therefore f'(x) = \frac{d}{dx} (2x^3 - 3x^2 - 12x + 6)$$

$$= 2 \times 3x^2 - 3 \times 2x - 12 \times 1 + 0$$

$$= 6x^2 - 6x - 12$$

$$= 6(x^2 - x - 2)$$

f is strictly decreasing if  $f'(x) < 0$

i.e. if  $6(x^2 - x - 2) < 0$

i.e. if  $x^2 - x - 2 < 0$

i.e. if  $x^2 - x < 2$

i.e. if  $x^2 - x + \frac{1}{4} < 2 + \frac{1}{4}$

i.e. if  $\left(x - \frac{1}{2}\right)^2 < \frac{9}{4}$

i.e. if  $-\frac{3}{2} < x - \frac{1}{2} < \frac{3}{2}$

i.e. if  $-\frac{3}{2} + \frac{1}{2} < x - \frac{1}{2} + \frac{1}{2} < \frac{3}{2} + \frac{1}{2}$

i.e. if  $-1 < x < 2$

$\therefore$  f is strictly decreasing if  $-1 < x < 2$ .

(ii)  $f(x) = x + \frac{25}{x}$

Solution:

$$f(x) = x + \frac{25}{x}, x \neq 0$$

$$\therefore f'(x) = \frac{d}{dx} \left(x + \frac{25}{x}\right)$$

$$= 1 + 25(-1)x^{-2}$$

$$= 1 - \frac{25}{x^2}$$

f is strictly decreasing if  $f'(x) < 0$

i.e. if  $1 - \frac{25}{x^2} < 0$

i.e. if  $1 < \frac{25}{x^2}$

i.e. if  $x^2 < 25$

i.e. if  $-5 < x < 5, x \neq 0$

i.e. if  $-5 < x < 5, x \neq 0$

i.e. if  $x \in (-5, 5) - \{0\}$

$\therefore f$  is strictly decreasing if  $x \in (-5, 5) - \{0\}$ .

(iii)  $f(x) = x^3 - 9x^2 + 24x + 12$

Solution:

$$f(x) = x^3 - 9x^2 + 24x + 12$$

$$\therefore f'(x) = \frac{d}{dx} (x^3 - 9x^2 + 24x + 12)$$

$$= 3x^2 - 9 \times 2x + 24 \times 1 + 0$$

$$= 3x^2 - 18x + 24$$

$$= 3(x^2 - 6x + 8)$$

$f$  is strictly decreasing if  $f'(x) < 0$

i.e. if  $3(x^2 - 6x + 8) < 0$

i.e. if  $x^2 - 6x + 8 < 0$

i.e. if  $x^2 - 6x < -8$

i.e. if  $x^2 - 6x + 9 < -8 + 9$

i.e. if  $(x - 3)^2 < 1$

i.e. if  $-1 < x - 3 < 1$

i.e. if  $-1 + 3 < x - 3 + 3 < 1 + 3$

i.e. if  $2 < x < 4$

i.e., if  $x \in (2, 4)$

$\therefore f$  is strictly decreasing if  $x \in (2, 4)$

Question 4.

Find the values of  $x$  for which the function  $f(x) = x^3 - 12x^2 - 144x + 13$

---

- (a) increasing  
(b) decreasing.

Solution:

$$f(x) = x^3 - 12x^2 - 144x + 13$$

$$\therefore f'(x) = \frac{d}{dx} (x^3 - 12x^2 - 144x + 13)$$

$$= 3x^2 - 12 \times 2x - 144 \times 1 + 0$$

$$= 3x^2 - 24x - 144$$

$$= 3(x^2 - 8x - 48)$$

(a)  $f$  is increasing if  $f'(x) \geq 0$

i.e. if  $3(x^2 - 8x - 48) \geq 0$

i.e. if  $x^2 - 8x - 48 \geq 0$

i.e. if  $x^2 - 8x \geq 48$

i.e. if  $x^2 - 8x + 16 \geq 48 + 16$

i.e. if  $(x - 4)^2 \geq 64$

i.e. if  $x - 4 \geq 8$  or  $x - 4 \leq -8$

i.e. if  $x > 12$  or  $x \leq -4$

$\therefore f$  is increasing if  $x \leq -4$  or  $x \geq 12$ ,

i.e.  $x \in (-\infty, -4] \cup [12, \infty)$ .

(b)  $f$  is decreasing if  $f'(x) \leq 0$

i.e. if  $3(x^2 - 8x - 48) \leq 0$

i.e. if  $x^2 - 8x - 48 \leq 0$

i.e. if  $x^2 - 8x \leq 48$

i.e. if  $x^2 - 8x + 16 \leq 48 + 16$

i.e. if  $(x - 4)^2 \leq 64$

i.e. if  $(x - 4)^2 \leq 64$

i.e. if  $-8 \leq x - 4 \leq 8$

i.e. if  $-4 \leq x \leq 12$

$\therefore f$  is decreasing if  $-4 \leq x \leq 12$ , i.e.  $x \in [-4, 12]$ .

Question 5.

Find the values of  $x$  for which  $f(x) = 2x^3 - 15x^2 - 144x - 7$  is

(a) strictly increasing

(b) strictly decreasing.

Solution:

$$f(x) = 2x^3 - 15x^2 - 144x - 7$$

$$f'(x) = \frac{d}{dx} (2x^3 - 15x^2 - 144x - 7)$$

$$= 2 \times 3x^2 - 15 \times 2x - 144 \times 1 - 0$$

$$= 6x^2 - 30x - 144$$

$$= 6(x^2 - 5x - 24)$$

(a)  $f$  is strictly increasing if  $f'(x) > 0$

i.e. if  $6(x^2 - 5x - 24) > 0$

i.e. if  $x^2 - 5x - 24 > 0$

i.e. if  $x^2 - 5x > 24$

i.e. if  $x^2 - 5x + \frac{25}{4} > 24 + \frac{25}{4}$

i.e. if  $\left(x - \frac{5}{2}\right)^2 > \frac{121}{4}$

i.e. if  $x - \frac{5}{2} > \frac{11}{2}$  or  $x - \frac{5}{2} < -\frac{11}{2}$  i.e. if  $x > 8$  or  $x < -3$

$\therefore f$  is strictly increasing, if  $x < -3$  or  $x > 8$ .

(b)  $f$  is strictly decreasing if  $f'(x) < 0$

i.e. if  $6(x^2 - 5x - 24) < 0$

i.e. if  $x^2 - 5x - 24 < 0$

---

i.e. if  $x^2 - 5x < 24$

i.e. if  $x^2 - 5x + \frac{25}{4} < 24 + \frac{25}{4}$

i.e. if  $\left(x - \frac{5}{2}\right)^2 < \frac{121}{4}$

i.e. if  $-\frac{11}{2} < x - \frac{5}{2} < \frac{11}{2}$

i.e. if  $-\frac{11}{2} + \frac{5}{2} < x - \frac{5}{2} + \frac{5}{2} < \frac{11}{2} + \frac{5}{2}$

i.e. if  $-3 < x < 8$

$\therefore f$  is strictly decreasing, if  $-3 < x < 8$ .

Question 6.

Find the values of  $x$  for which  $f(x) = \frac{x}{x^2+1}$  is

(a) strictly increasing

(b) strictly decreasing.

Solution:

$$f(x) = \frac{x}{x^2+1}$$

$$\therefore f'(x) = \frac{d}{dx} \left( \frac{x}{x^2+1} \right)$$

$$= \frac{(x^2+1) \cdot \frac{d}{dx}(x) - x \frac{d}{dx}(x^2+1)}{(x^2+1)^2}$$

$$= \frac{(x^2+1)(1) - x(2x+0)}{(x^2+1)^2}$$

$$= \frac{x^2 + 1 - 2x^2}{(x^2 + 1)^2} = \frac{1 - x^2}{(x^2 + 1)^2}$$

(a) f is strictly increasing if  $f'(x) > 0$

i.e. if  $\frac{1-x^2}{(x^2+1)^2} > 0$

i.e. if  $1 - x^2 > 0$  ..... [ $\because (x^2 + 1)^2 > 0$ ]

i.e. if  $1 > x^2$

i.e. if  $x^2 < 1$

i.e. if  $-1 < x < 1$

$\therefore$  f is strictly increasing if  $-1 < x < 1$

(b) f is strictly decreasing if  $f'(x) < 0$

i.e. if  $\frac{1-x^2}{(x^2+1)^2} < 0$

i.e. if  $1 - x^2 < 0$  ..... [ $\because (x^2 + 1)^2 > 0$ ]

i.e. if  $1 < x^2$  i.e. if  $x^2 > 1$

i.e. if  $x > 1$  or  $x < -1$

$\therefore$  f is strictly decreasing if  $x < -1$  or  $x > 1$

i.e.  $x \in (-\infty, -1) \cup (1, \infty)$ .

Question 7.

Show that  $f(x) = 3x + \frac{1}{3x}$  is increasing in  $(\frac{1}{3}, 1)$  and decreasing in  $(\frac{1}{9}, \frac{1}{3})$

Solution:

$$f(x) = 3x + \frac{1}{3x}$$

$$\therefore f'(x) = 3 \frac{d}{dx} (x) + \frac{1}{3} \frac{d}{dx} (x^{-1})$$

$$= 3 \times 1 + \frac{1}{3} (-1) x^{-2} = 3 - \frac{1}{3x^2}$$

Now,  $f$  is increasing if  $f'(x) > 0$  and is decreasing if  $f'(x) < 0$ .

Let  $x \in \left(\frac{1}{3}, 1\right)$ . Then  $\frac{1}{3} < x < 1$

$$\therefore \frac{1}{9} < x^2 < 1$$

$$\therefore \frac{1}{3} < 3x^2 < 3$$

$$\therefore 3 > \frac{1}{3x^2} > \frac{1}{3}$$

$$\therefore -3 < -\frac{1}{3x^2} < -\frac{1}{3}$$

$$\therefore 3 - 3 < 3 - \frac{1}{3x^2} < 3 - \frac{1}{3}$$

$$\therefore 0 < f'(x) < \frac{8}{3}$$

$\therefore f'(x) > 0$  for all  $x \in \left(\frac{1}{3}, 1\right)$

$\therefore f$  is increasing in the interval  $\left(\frac{1}{3}, 1\right)$ .

Let  $x \in \left(\frac{1}{9}, \frac{1}{3}\right)$ . Then  $\frac{1}{9} < x < \frac{1}{3}$

$$\therefore \frac{1}{81} < x^2 < \frac{1}{9}$$

$$\therefore \frac{1}{27} < 3x^2 < \frac{1}{3} \quad \therefore 27 > \frac{1}{3x^2} > 3$$

$$\therefore -27 < -\frac{1}{3x^2} < -3$$

$$\therefore 3 - 27 < 3 - \frac{1}{3x^2} < 3 - 3 \quad \therefore -24 < f'(x) < 0$$

$$\therefore f'(x) < 0 \text{ for all } x \in \left(\frac{1}{9}, \frac{1}{3}\right)$$

$$\therefore f \text{ is decreasing in the interval } \left(\frac{1}{9}, \frac{1}{3}\right).$$

Question 8.

Show that  $f(x) = x - \cos x$  is increasing for all  $x$ .

Solution:

$$f(x) = x - \cos x$$

$$\therefore f'(x) = \frac{d}{dx} (x - \cos x)$$

$$= 1 - (-\sin x)$$

$$= 1 + \sin x$$

Now,  $-1 \leq \sin x \leq 1$  for all  $x \in \mathbb{R}$

$$\therefore -1 + 1 \leq 1 + \sin x \leq 1 + 1 \text{ for all } x \in \mathbb{R}$$

$$\therefore 0 \leq f'(x) \leq 2 \text{ for all } x \in \mathbb{R}$$

$$\therefore f'(x) \geq 0 \text{ for all } x \in \mathbb{R}$$

$\therefore f$  is increasing for all  $x$ .

Question 9.

Find the maximum and minimum of the following functions:

(i)  $y = 5x^3 + 2x^2 - 3x$

Solution:

$$y = 5x^3 + 2x^2 - 3x$$

$$\begin{aligned}\therefore \frac{dy}{dx} &= \frac{d}{dx}(5x^3 + 2x^2 - 3x) \\ &= 5 \times 3x^2 + 2 \times 2x - 3 \times 1 \\ &= 15x^2 + 4x - 3\end{aligned}$$

$$\begin{aligned}\text{and } \frac{d^2y}{dx^2} &= \frac{d}{dx}(15x^2 + 4x - 3) \\ &= 15 \times 2x + 4 \times 1 - 0 = 30x + 4\end{aligned}$$

$$\frac{dy}{dx} = 0 \text{ gives } 15x^2 + 4x - 3 = 0$$

$$\therefore 15x^2 + 9x - 5x - 3 = 0$$

$$\therefore 3x(5x + 3) - 1(5x + 3) = 0$$

$$\therefore (5x + 3)(3x - 1) = 0$$

$$\therefore x = -\frac{3}{5} \text{ or } x = \frac{1}{3}$$

$$\therefore \text{the roots of } \frac{dy}{dx} = 0 \text{ are } x_1 = -\frac{3}{5} \text{ and } x_2 = \frac{1}{3}.$$

**Method 1 (Second Derivative Test) :**

$$(a) \left( \frac{d^2y}{dx^2} \right)_{\text{at } x = -\frac{3}{5}} = 30 \left( -\frac{3}{5} \right) + 4 = -14 < 0$$

∴ by the second derivative test,  $y$  is maximum at

$$x = -\frac{3}{5} \text{ and maximum value of } y \text{ at } x = -\frac{3}{5}$$

$$= 5 \left( -\frac{3}{5} \right)^3 + 2 \left( -\frac{3}{5} \right)^2 - 3 \left( -\frac{3}{5} \right)$$

$$= \frac{-27}{25} + \frac{18}{25} + \frac{9}{5} = \frac{36}{25}$$

$$(b) \left( \frac{d^2y}{dx^2} \right)_{\text{at } x = \frac{1}{3}} = 30 \left( \frac{1}{3} \right) + 4 = 14 > 0$$

∴ by the second derivative test,  $y$  is minimum at

$$x = \frac{1}{3} \text{ and minimum value of } y \text{ at } x = \frac{1}{3}$$

$$= 5 \left( \frac{1}{3} \right)^3 + 2 \left( \frac{1}{3} \right)^2 - 3 \left( \frac{1}{3} \right)$$

$$= \frac{5}{27} + \frac{2}{9} - 1 = -\frac{16}{27}$$

Hence, the function has maximum value  $\frac{36}{25}$  at  $x = -\frac{3}{5}$

and minimum value  $-\frac{16}{27}$  at  $x = \frac{1}{3}$ .

**Method 2 (First Derivative Test) :**

$$(a) \frac{dy}{dx} = 15x^2 + 4x - 3 = (5x + 3)(3x - 1)$$

$$\text{Consider } x = -\frac{3}{5}$$

Let  $h$  be a small positive number. Then

$$\begin{aligned} \left(\frac{dy}{dx}\right)_{\text{at } x = -\frac{3}{5} - h} &= \left[5\left(-\frac{3}{5} - h\right) + 3\right] \left[3\left(-\frac{3}{5} - h\right) - 1\right] \\ &= (-3 - 5h + 3) \left(-\frac{9}{5} - 3h - 1\right) \\ &= -5h \left(-\frac{14}{5} - 3h\right) \\ &= 5h \left(\frac{14}{5} + 3h\right) > 0 \end{aligned}$$

$$\begin{aligned} \text{and } \left(\frac{dy}{dx}\right)_{\text{at } x = -\frac{3}{5} + h} &= \left[5\left(-\frac{3}{5} + h\right) + 3\right] \left[3\left(-\frac{3}{5} + h\right) - 1\right] \\ &= (-3 + 5h + 3) \left(-\frac{9}{5} + 3h - 1\right) \\ &= 5h \left(3h - \frac{14}{5}\right) < 0, \end{aligned}$$

as  $h$  is small positive number.

$\therefore$  by the first derivative test,  $y$  is maximum at

$$x = -\frac{3}{5} \text{ and maximum value of } y \text{ at } x = -\frac{3}{5}$$

$$= 5\left(-\frac{3}{5}\right)^3 + 2\left(-\frac{3}{5}\right)^2 - 3\left(-\frac{3}{5}\right)$$

$$= -\frac{27}{25} + \frac{18}{25} + \frac{9}{5} = \frac{36}{25}$$

$$(b) \frac{dy}{dx} = 15x^2 + 4x - 3 = (5x + 3)(3x - 1)$$

$$\text{Consider } x = \frac{1}{3}$$

Let  $h$  be a small positive number. Then

$$\left(\frac{dy}{dx}\right)_{\text{at } x = \frac{1}{3} - h} = \left[5\left(\frac{1}{3} - h\right) + 3\right] \left[3\left(\frac{1}{3} - h\right) - 1\right]$$

$$= \left(\frac{5}{3} - 5h + 3\right)(1 - 3h - 1)$$

$$= \left(\frac{14}{3} - 5h\right)(-3h) < 0, \text{ as } h \text{ is small}$$

positive number

$$\text{and } \left(\frac{dy}{dx}\right)_{\text{at } x = \frac{1}{3} + h} = \left[5\left(\frac{1}{3} + h\right) + 3\right] \left[3\left(\frac{1}{3} + h\right) - 1\right]$$

$$= \left(\frac{5}{3} + 5h + 3\right)(1 + 3h - 1)$$

$$= \left(\frac{14}{3} + 5h\right)(3h) > 0$$

∴ by the first derivative test,  $y$  is minimum at  $x = \frac{1}{3}$

and minimum value of  $y$  at  $x = \frac{1}{3}$

$$= 5\left(\frac{1}{3}\right)^3 + 2\left(\frac{1}{3}\right)^2 - 3\left(\frac{1}{3}\right)$$

$$= \frac{5}{27} + \frac{2}{9} - 1 = \frac{-16}{27}$$

Hence, the function has maximum value  $\frac{36}{25}$  at

$x = -\frac{3}{5}$  and minimum value  $-\frac{16}{27}$  at  $x = \frac{1}{3}$ .

(ii)  $f(x) = 2x^3 - 21x^2 + 36x - 20$

Solution:

$$f(x) = 2x^3 - 21x^2 + 36x - 20$$

$$\therefore f'(x) = \frac{d}{dx} (2x^3 - 21x^2 + 36x - 20)$$

$$= 2 \times 3x^2 - 21 \times 2x + 36 \times 1 - 0$$

$$= 6x^2 - 42x + 36$$

$$\text{and } f''(x) = \frac{d}{dx} (6x^2 - 42x + 36)$$

$$= 6 \times 2x - 42 \times 1 + 0$$

$$= 12x - 42$$

$$f'(x) = 0 \text{ gives } 6x^2 - 42x + 36 = 0$$

$$\therefore x^2 - 7x + 6 = 0$$

$$\therefore (x-1)(x-6) = 0$$

the roots of  $f'(x) = 0$  are  $x_1 = 1$  and  $x_2 = 6$ .

Method 1 (Second Derivative Test):

$$(a) f''(1) = 12(1) - 42 = -30 < 0$$

$\therefore$  by the second derivative test,  $f$  has maximum at  $x = 1$

and maximum value of  $f$  at  $x = 1$

$$f(1) = 2(1)^3 - 21(1)^2 + 36(1) - 20$$

$$= 2 - 21 + 36 - 20$$

$$= -3$$

$$(b) f''(6) = 12(6) - 42 = 30 > 0$$

$\therefore$  by the second derivative test,  $f$  has minimum at  $x = 6$

and minimum value of  $f$  at  $x = 6$

$$f(6) = 2(6)^3 - 21(6)^2 + 36(6) - 20$$

$$= 432 - 756 + 216 - 20$$

$$= -128.$$

Hence, the function  $f$  has maximum value  $-3$  at  $x = 1$  and minimum value  $-128$  at  $x = 6$ .

Method 2 (First Derivative Test):

$$(a) f'(x) = 6(x - 1)(x - 6)$$

Consider  $x = 1$

Let  $h$  be a small positive number. Then

$$f'(1 - h) = 6(1 - h - 1)(1 - h - 6)$$

$$= 6(-h)(-5 - h)$$

$$= 6h(5 + h) > 0$$

$$\text{and } f'(1 + h) = 6(1 + h - 1)(1 + h - 6)$$

$$= 6h(h - 5) < 0, \text{ as } h \text{ is small positive number.}$$

$= 6h(h - 5) < 0$ , as  $h$  is small positive number.

$\therefore$  by the first derivative test,  $f$  has maximum at  $x = 1$  and maximum value of  $f$  at  $x = 1$

$$f(1) = 2(1)^3 - 21(1)^2 + 36(1) - 20$$

$$= 2 - 21 + 36 - 20$$

$$= -3$$

(b)  $f'(x) = 6(x - 1)(x - 6)$

Consider  $x = 6$

Let  $h$  be a small positive number. Then

$$f'(6 - h) = 6(6 - h - 1)(6 - h - 6)$$

$$= 6(5 - h)(-h)$$

$= -6h(5 - h) < 0$ , as  $h$  is small positive number

$$\text{and } f'(6 + h) = 6(6 + h - 1)(6 + h - 6) = 6(5 + h)(h) > 0$$

$\therefore$  by the first derivative test,  $f$  has minimum at  $x = 6$

and minimum value of  $f$  at  $x = 6$

$$f(6) = 2(6)^3 - 21(6)^2 + 36(6) - 20$$

$$= 432 - 756 + 216 - 20$$

$$= -128$$

Hence, the function  $f$  has maximum value  $-3$  at  $x = 1$

and minimum value  $-128$  at  $x = 6$ .

(iii)  $f(x) = x^3 - 9x^2 + 24x$

Solution:

$$f(x) = x^3 - 9x^2 + 24x$$

$$\therefore f'(x) = \frac{d}{dx} (x^3 - 9x^2 + 24x)$$

$$\therefore f'(x) = \frac{d}{dx} (x^3 - 9x^2 + 24x)$$

$$= 3x^2 - 9 \times 2x + 24 \times 1$$

$$= 3x^2 - 18x + 24$$

$$\text{and } f''(x) = \frac{d}{dx} (3x^2 - 18x + 24)$$

$$= 3 \times 2x - 18 \times 1 + 0$$

$$= 6x - 18$$

$$f'(x) = 0 \text{ gives } 3x^2 - 18x + 24 = 0$$

$$\therefore x^2 - 6x + 8 = 0$$

$$\therefore (x - 2)(x - 4) = 0$$

$\therefore$  the roots of  $f'(x) = 0$  are  $x_1 = 2$  and  $x_2 = 4$ .

$$(a) f''(2) = 6(2) - 18 = -6 < 0$$

$\therefore$  by the second derivative test,  $f$  has maximum at  $x = 2$

and maximum value of  $f$  at  $x = 2$

$$f(2) = (2)^3 - 9(2)^2 + 24(2)$$

$$= 8 - 36 + 48$$

$$= 20$$

$$(b) f''(4) = 6(4) - 18 = 6 > 0$$

$\therefore$  by the second derivative test,  $f$  has minimum at  $x = 4$

and minimum value of  $f$  at  $x = 4$

$$f(4) = (4)^3 - 9(4)^2 + 24(4)$$

$$= 64 - 144 + 96$$

$$= 16$$

Hence, the function  $f$  has maximum value 20 at  $x = 2$  and minimum value 16 at  $x = 4$ .

$$(iv) f(x) = x^2 + \frac{16}{x^2}$$

Solution:

$$f(x) = x^2 + \frac{16}{x^2}$$

$$\begin{aligned}\therefore f'(x) &= \frac{d}{dx}(x^2) + 16 \frac{d}{dx}(x^{-2}) \\ &= 2x + 16(-2)x^{-3} = 2x - \frac{32}{x^3}\end{aligned}$$

$$\begin{aligned}\text{and } f''(x) &= \frac{d}{dx}(2x) - 32 \frac{d}{dx}(x^{-3}) \\ &= 2 \times 1 - 32(-3)x^{-4} = 2 + \frac{96}{x^4}\end{aligned}$$

$$f'(x) = 0 \text{ gives } 2x - \frac{32}{x^3} = 0$$

$$\therefore 2x^4 - 32 = 0 \quad \therefore x^4 = 16$$

$$\therefore x = \pm 2$$

$\therefore$  the roots of  $f'(x) = 0$  are  $x_1 = 2$  and  $x_2 = -2$ .

$$(a) f''(2) = 2 + \frac{96}{(2)^4} = 8 > 0$$

$\therefore$  by the second derivative test,  $f$  has minimum at  $x = 2$  and minimum value of  $f$  at  $x = 2$

$$= f(2) = (2)^2 + \frac{16}{(2)^2} = 4 + 4 = 8$$

$$(b) f''(-2) = 2 + \frac{96}{(-2)^4} = 8 > 0$$

$\therefore$  by the second derivative test,  $f$  has minimum at  $x = -2$  and minimum value of  $f$  at  $x = -2$

$$= f(-2) = (-2)^2 + \frac{16}{(-2)^2} = 4 + 4 = 8$$

Hence, the function  $f$  has minimum value 8 at  $x = \pm 2$ .

(v)  $f(x) = x \log x$

Solution:

$$f(x) = x \log x$$

$$\begin{aligned}\therefore f'(x) &= \frac{d}{dx}(x \log x) \\ &= x \cdot \frac{d}{dx}(\log x) + \log x \cdot \frac{d}{dx}(x) \\ &= x \times \frac{1}{x} + (\log x) \times 1 = 1 + \log x\end{aligned}$$

$$\text{and } f''(x) = \frac{d}{dx}(1 + \log x) = 0 + \frac{1}{x} = \frac{1}{x}$$

Now,  $f'(x) = 0$ , if  $1 + \log x = 0$

i.e. if  $\log x = -1 = -\log e$

i.e. if  $\log x = \log(e^{-1}) = \log \frac{1}{e}$

i.e. if  $x = \frac{1}{e}$

When  $x = \frac{1}{e}$ ,  $f''(x) = \frac{1}{(1/e)} = e > 0$

∴ by the second derivative test,  $f$  is minimum at

$$x = \frac{1}{e}$$

Minimum value of  $f$  at  $x = \frac{1}{e}$

$$= \frac{1}{e} \log\left(\frac{1}{e}\right) = \frac{1}{e} \log(e^{-1})$$

$$= \frac{1}{e} \cdot (-1) \log e$$

$$= -\frac{1}{e} \quad \dots [\because \log e = 1]$$

(vi)  $f(x) = \frac{\log x}{x}$

Solution:

$$f(x) = \frac{\log x}{x}$$

$$\therefore f'(x) = \frac{d}{dx} \left( \frac{\log x}{x} \right)$$

$$= \frac{x \frac{d}{dx}(\log x) - \log x \frac{d}{dx}(x)}{x^2}$$

$$= \frac{x\left(\frac{1}{x}\right) - (\log x)(1)}{x^2} = \frac{1 - \log x}{x^2}$$

$$\text{and } f''(x) = \frac{d}{dx} \left( \frac{1 - \log x}{x^2} \right)$$

$$= \frac{x^2 \frac{d}{dx} (1 - \log x) - (1 - \log x) \frac{d}{dx} (x^2)}{x^4}$$

$$= \frac{x^2 \left( 0 - \frac{1}{x} \right) - (1 - \log x) \times 2x}{x^4}$$

$$= \frac{-x - 2x + 2x \log x}{x^4} = \frac{x(2 \log x - 3)}{x^4}$$

$$\therefore f''(x) = \frac{2 \log x - 3}{x^3}$$

$$\text{Now, } f'(x) = 0, \text{ if } \frac{1 - \log x}{x^2} = 0$$

i.e. if  $1 - \log x = 0$ , i.e. if  $\log x = 1 = \log e$

i.e. if  $x = e$

$$\text{and } f''(e) = \frac{2 \log e - 3}{e^3} = \frac{-1}{e^3} < 0 \quad \dots [\because \log e = 1]$$

$\therefore$  by the second derivative test,  $f(x)$  is maximum at  $x = e$ .

Maximum value of  $f$  at  $x = e$

$$= \frac{\log e}{e} = \frac{1}{e} \quad \dots [\because \log e = 1]$$

Question 10.

Divide the number 30 into two parts such that their product is maximum.

Solution:

Let the first part of 30 be  $x$ .

Then the second part is  $30 - x$ .

$\therefore$  their product =  $x(30 - x) = 30x - x^2 = f(x) \dots\dots$ (Say)

$$\therefore f'(x) = \frac{d}{dx} (30x - x^2)$$

$$= 30 \times 1 - 2x$$

$$= 30 - 2x$$

$$\text{and } f''(x) = \frac{d}{dx} (30 - 2x)$$

$$= 0 - 2 \times 1$$

$$= -2$$

The root of the equation  $f'(x) = 0$ ,

i.e.  $30 - 2x = 0$  is  $x = 15$  and  $f''(15) = -2 < 0$

$\therefore$  by the second derivative test,  $f$  is maximum at  $x = 15$ .

Hence, the required parts of 30 are 15 and 15.

Question 11.

Divide the number 20 into two parts such that the sum of their squares is minimum.

Solution:

---

Let the first part of 20 be  $x$ .

Then the second part is  $20 - x$ .

$\therefore$  sum of their squares =  $x^2 + (20 - x)^2 = f(x)$  ..... (Say)

$$\therefore f'(x) = \frac{d}{dx} [x^2 + (20 - x)^2]$$

$$= 2x + 2(20 - x) \cdot \frac{d}{dx} (20 - x)$$

$$= 2x + 2(20 - x) \times (0 - 1)$$

$$= 2x - 40 + 2x$$

$$= 4x - 40$$

$$\text{and } f''(x) = \frac{d}{dx} (4x - 40)$$

$$= 4 \times 1 - 0$$

$$= 4$$

The root of the equation  $f'(x) = 0$ ,

i.e.  $4x - 40 = 0$  is  $x = 10$  and  $f''(10) = 4 > 0$

$\therefore$  by the second derivative test,  $f$  is minimum at  $x = 10$ .

Hence, the required parts of 20 are 10 and 10.

Question 12.

A wire of length 36 meters is bent in the form of a rectangle. Find its dimensions if the area of the rectangle is maximum.

Solution:

Let  $x$  metres and  $y$  metres be the length and breadth of the rectangle.

Then its perimeter is  $2(x + y) = 36$

$$x + y = 18$$

$$y = 18 - x$$

Area of the rectangle =  $xy = x(18 - x)$

$$\text{Let } f(x) = x(18 - x) = 18x - x^2$$

$$\therefore f'(x) = \frac{d}{dx} (18x - x^2) = 18 - 2x$$

$$\text{and } f''(x) = \frac{d}{dx} (18 - 2x) = 0 - 2 \times 1 = -2$$

Now,  $f'(x) = 0$ , if  $18 - 2x = 0$

i.e. if  $x = 9$

$$\text{and } f''(9) = -2 < 0$$

$\therefore$  by the second derivative test,  $f$  has maximum value at  $x = 9$ .

When  $x = 9$ ,  $y = 18 - 9 = 9$

$\therefore x = 9$  cm,  $y = 9$  cm

$\therefore$  the rectangle is a square of side 9 metres.

Question 13.

A ball is thrown in the air. Its height at any time  $t$  is given by  $h = 3 + 14t - 5t^2$ . Find the maximum height it can reach.

Solution:

The height  $h$  at any  $t$  is given by  $h = 3 + 14t - 5t^2$

$$\therefore \frac{dh}{dt} = \frac{d}{dt} (3 + 14t - 5t^2)$$

$$= 0 + 14 \times 1 - 5 \times 2t = 14 - 10t$$

$$\text{and } \frac{d^2h}{dt^2} = \frac{d}{dt} (14 - 10t) = 0 - 10 \times 1 = -10$$

The root of  $\frac{dh}{dt} = 0$ , i.e.  $14 - 10t = 0$  is  $t = \frac{14}{10} = \frac{7}{5}$

$$\text{and } \left( \frac{d^2h}{dt^2} \right)_{\text{at } t = \frac{7}{5}} = -10 < 0$$

∴ by the second derivative test,  $h$  is maximum at  $t = \frac{7}{5}$ .

∴ maximum height =  $(3 + 14t - 5t^2)_{\text{at } t = \frac{7}{5}}$

$$= 3 + 14\left(\frac{7}{5}\right) - 5\left(\frac{7}{5}\right)^2$$

$$= 3 + \frac{98}{5} - \frac{245}{5}$$

$$= \frac{75 + 490 - 245}{25}$$


$$= \frac{320}{25} = 12.8$$

Hence, the maximum height the ball can reach = 12.8 units.

Question 14.

Find the largest size of a rectangle that can be inscribed in a semicircle of radius 1 unit, so that two vertices lie on the diameter.

Solution:



Let ABCD be the rectangle inscribed in a semicircle of radius 1 unit such that the vertices A and B lie on the diameter.

Let  $AB = DC = x$  and  $BC = AD = y$ .

Let O be the centre of the semicircle.

Join OC and OD. Then  $OC = OD = \text{radius} = 1$ .

Also,  $AD = BC$  and  $m\angle A = m\angle B = 90^\circ$ .

$\therefore OA = OB$

$\therefore OB = \frac{1}{2} AB = \frac{x}{2}$

In right angled triangle OBC,

$$OB^2 + BC^2 = OC^2$$

$$\therefore \left(\frac{x}{2}\right)^2 + y^2 = 1^2$$

$$\therefore y^2 = 1 - \frac{x^2}{4} = \frac{1}{4}(4 - x^2)$$

$$\therefore y = \frac{1}{2}\sqrt{4 - x^2} \quad \dots [\because y > 0]$$

$$\text{Area of the rectangle} = xy = x \cdot \frac{1}{2}\sqrt{4 - x^2}$$

$$\text{Let } f(x) = \frac{1}{2}x\sqrt{4 - x^2} = \frac{1}{2}\sqrt{4x^2 - x^4}$$

$$\therefore f'(x) = \frac{1}{2} \frac{d}{dx} (\sqrt{4x^2 - x^4})$$

$$= \frac{1}{2} \times \frac{1}{2\sqrt{4x^2 - x^4}} \cdot \frac{d}{dx} (4x^2 - x^4)$$

$$= \frac{1}{4\sqrt{4x^2 - x^4}} \times (4 \times 2x - 4x^3)$$

$$= \frac{4x(2 - x^2)}{4x\sqrt{4 - x^2}} = \frac{2 - x^2}{\sqrt{4 - x^2}} \quad \dots [\because x \neq 0]$$

$$\text{and } f''(x) = \frac{d}{dx} \left( \frac{2 - x^2}{\sqrt{4 - x^2}} \right) = \frac{d}{dx} \left[ \frac{(4 - x^2) - 2}{\sqrt{4 - x^2}} \right]$$

$$= \frac{d}{dx} \left[ \sqrt{4 - x^2} - \frac{2}{\sqrt{4 - x^2}} \right]$$

$$= \frac{d}{dx} (\sqrt{4 - x^2}) - 2 \frac{d}{dx} (4 - x^2)^{-\frac{1}{2}}$$

$$= \frac{1}{2\sqrt{4 - x^2}} \cdot \frac{d}{dx} (4 - x^2) - 2 \left( -\frac{1}{2} \right) (4 - x^2)^{-\frac{3}{2}} \cdot \frac{d}{dx} (4 - x^2)$$

$$= \frac{1}{2\sqrt{4 - x^2}} \times (0 - 2x) + \frac{1}{(4 - x^2)^{\frac{3}{2}}} \times (0 - 2x)$$

$$= \frac{-x}{\sqrt{4 - x^2}} - \frac{2x}{(4 - x^2)^{\frac{3}{2}}}$$

$$= \frac{-x(4 - x^2) - 2x}{(4 - x^2)^{\frac{3}{2}}}$$

$$= \frac{-4x + x^3 - 2x}{(4 - x^2)^{\frac{3}{2}}} = \frac{x^3 - 6x}{(4 - x^2)^{\frac{3}{2}}}$$

For maximum value of  $f(x)$ ,  $f'(x) = 0$

$$\therefore \frac{2-x^2}{\sqrt{4-x^2}} = 0 \quad \therefore 2-x^2 = 0$$

$$\therefore x^2 = 2 \quad \therefore x = \sqrt{2} \quad \dots [\because x > 0]$$

$$\text{Now, } f''(\sqrt{2}) = \frac{(\sqrt{2})^3 - 6\sqrt{2}}{[4 - (\sqrt{2})^2]^{\frac{3}{2}}} = \frac{-4\sqrt{2}}{2\sqrt{2}} = -2 < 0$$

$\therefore$  by the second derivative test,  $f$  is maximum when

$$x = \sqrt{2}$$

$$\text{When } x = \sqrt{2}, y = \frac{1}{2}\sqrt{4-x^2} = \frac{1}{2}\sqrt{4-2}$$

$$= \frac{1}{2} \times \sqrt{2} = \frac{1}{\sqrt{2}}$$

$$\therefore x = \sqrt{2} \text{ and } y = \frac{1}{\sqrt{2}}$$

Hence, the area of the rectangle is maximum (i.e. rectangle has the largest size) when its length is  $\sqrt{2}$  units and breadth is  $\frac{1}{\sqrt{2}}$  unit.

Question 15.

An open cylindrical tank whose base is a circle is to be constructed of metal sheet so as to contain a volume of  $na^3$  cu cm of water. Find the dimensions so that the quantity of the metal sheet required is minimum.

Solution:

Let  $x$  be the radius of the base,  $h$  be the height,  $V$  be the volume and  $S$  be the total surface area of the cylindrical tank.

---

Then  $V = \pi a^3 \dots$  (Given)

$$\therefore \pi x^2 h = \pi a^3$$

$$\therefore h = \frac{a^3}{x^2} \dots\dots\dots(1)$$

$$\text{Now, } S = 2\pi x h + \pi x^2$$

$$= 2\pi x \left( \frac{a^3}{x^2} \right) + \pi x^2 \quad \dots \text{ [By (1)]}$$

$$= \frac{2\pi a^3}{x} + \pi x^2$$

$$\therefore \frac{dS}{dx} = \frac{d}{dx} \left( \frac{2\pi a^3}{x} + \pi x^2 \right)$$

$$= 2\pi a^3 (-1)x^{-2} + \pi \times 2x = \frac{-2\pi a^3}{x^2} + 2\pi x$$

$$\text{and } \frac{d^2S}{dx^2} = \frac{d}{dx} \left( \frac{-2\pi a^3}{x^2} + 2\pi x \right)$$

$$= -2\pi a^3 (-2)x^{-3} + 2\pi \times 1$$

$$= \frac{4\pi a^3}{x^3} + 2\pi$$

$$\text{Now, } \frac{dS}{dx} = 0 \text{ gives } \frac{-2\pi a^3}{x^2} + 2\pi x = 0$$

$$\therefore -2\pi a^3 + 2\pi x^3 = 0$$

$$\therefore 2\pi x^3 = 2\pi a^3 \quad \therefore x = a$$

$$\text{and } \left( \frac{d^2S}{dx^2} \right)_{\text{at } x=a} = \frac{4\pi a^3}{a^3} + 2\pi = 6\pi > 0$$

$\therefore$  by the second derivative test,  $S$  is minimum when  $x = a$

When  $x = a$ , from (1)


$$h = \frac{a^3}{a^2} = a$$

Hence, the quantity of metal sheet is minimum when radius height = a cm.

Question 16.

The perimeter of a triangle is 10 cm. If one of the sides is 4 cm. What are the other two sides of the triangle for its maximum area?

Solution:



Let ABC be the triangle such that the side  $BC = a = 4$  cm.

Also, the perimeter of the triangle is 10 cm.

$$\text{i.e. } a + b + c = 10$$

$$\therefore 2s = 10$$

$$\therefore s = 5$$

$$\text{Also, } 4 + b + c = 10$$

$$\therefore b + c = 6$$

$$\therefore b = 6 - c$$

Let  $\Delta$  be the area of the triangle.

$$\text{Then } \Delta = \sqrt{s(s-a)(s-b)(s-c)}$$

$$= \sqrt{5(5-4)(5-6+c)(5-c)}$$

$$= \sqrt{5(c-1)(5-c)}$$

$$\therefore \Delta^2 = 5(c-1)(5-c) = 5(5c - c^2 - 5 + c)$$

$$\therefore \Delta^2 = 5(-c^2 + 6c - 5)$$

Differentiating both sides w.r.t.  $c$ , we get

$$2\Delta \frac{d\Delta}{dc} = 5 \frac{d}{dc} (-c^2 + 6c - 5)$$

$$= 5(-2c + 6 \times 1 - 0) = 5(-2c + 6)$$

$$\therefore \frac{d\Delta}{dc} = \frac{5(-c+3)}{\Delta}$$

$$\text{and } \frac{d^2\Delta}{dc^2} = 5 \frac{d}{dc} \left( \frac{-c+3}{\Delta} \right)$$

$$= 5 \cdot \frac{\Delta \frac{d}{dc}(-c+3) - (-c+3) \frac{d\Delta}{dc}}{\Delta^2}$$

$$= 5 \cdot \frac{\Delta(-1+0) - (c+3) \frac{d\Delta}{dc}}{\Delta^2}$$

$$= \frac{5}{\Delta^2} \left( -\Delta - (c+3) \frac{d\Delta}{dc} \right)$$

$$= \frac{-5}{\Delta^2} \left[ \Delta + (c+3) \frac{d\Delta}{dc} \right]$$

For maximum  $\Delta$ ,  $\frac{d\Delta}{dc} = 0$

$$\therefore \frac{5(-c+3)}{\Delta} = 0$$

$$\therefore -c+3=0 \quad \dots [\because \Delta \neq 0]$$

$$\therefore c=3$$

$$\text{If } c=3, \Delta = \sqrt{5(3-1)(5-3)} = 2\sqrt{5}$$

$$\begin{aligned} \therefore \left( \frac{d^2\Delta}{dc^2} \right)_{\text{at } c=3} &= \frac{-5}{4 \times 5} [2\sqrt{5} + (3+3)(0)] \\ &= -\frac{\sqrt{5}}{2} < 0 \end{aligned}$$

$\therefore$  by the second derivative test,  $\Delta$  is maximum when  $c = 3$ .

When  $c = 3$ ,  $b = 6 - c = 6 - 3 = 3$

Hence, the area of the triangle is maximum when the other two sides are 3 cm and 3 cm.

Question 17.

A box with a square base is to have an open top. The surface area of the box is 192 sq cm. What should be its dimensions in order that the volume is largest?

Solution:

Let  $x$  cm be the side of square base and  $h$  cm be its height.

Then  $x^2 + 4xh = 192$

$$\therefore h = \frac{192 - x^2}{4x} \dots\dots (1)$$

Let  $V$  be the volume of the box.

$$\text{Then } V = x^2h = x^2 \left( \frac{192 - x^2}{4x} \right) \quad \dots \text{ [By (1)]}$$

$$\therefore V = \frac{1}{4}(192x - x^3)$$

$$\begin{aligned} \therefore \frac{dV}{dx} &= \frac{1}{4} \frac{d}{dx}(192x - x^3) \\ &= \frac{1}{4}(192 \times 1 - 3x^2) = \frac{3}{4}(64 - x^2) \end{aligned}$$

$$\begin{aligned} \text{and } \frac{d^2V}{dx^2} &= \frac{3}{4} \frac{d}{dx}(64 - x^2) \\ &= \frac{3}{4}(0 - 2x) = -\frac{3}{2}x \end{aligned}$$

$$\text{For maximum } V, \quad \frac{dV}{dx} = 0$$

$$\therefore \frac{3}{4}(64 - x^2) = 0$$

$$\therefore x^2 = 64 \quad \therefore x = 8 \quad \dots [\because x > 0]$$

$$\text{and } \left( \frac{d^2V}{dx^2} \right)_{\text{at } x=8} = -\frac{3}{2} \times 8 = -12 < 0$$

$\therefore$  by the second derivative test,  $V$  is maximum at  $x = 8$ .

$$\text{If } x = 8, h = \frac{192 - 64}{4(8)} = \frac{128}{32} = 4$$

Hence, the volume of the box is largest, when the side of square base is 8 cm and its height is 4 cm.

Question 18.

The profit function  $P(x)$  of a firm, selling  $x$  items per day is given by  $P(x) = (150 - x)x - 1625$ . Find the number of items the firm should manufacture to get maximum profit. Find the maximum profit.

Solution:

Profit function  $P(x)$  is given by

$$P(x) = (150 - x)x - 1625 = 150x - x^2 - 1625$$

$$\therefore P'(x) = \frac{d}{dx} (150x - x^2 - 1625)$$

$$= 150 \times 1 - 2x - 0$$

$$= 150 - 2x$$

$$\text{and } P''(x) = \frac{d}{dx} (150 - 2x)$$

$$= 0 - 2 \times 1$$

$$= -2$$

Now,  $P'(x) = 0$  gives,  $150 - 2x = 0$

$$\therefore x = 75$$

$$\text{and } P''(75) = -2 < 0$$

$\therefore$  by the second derivative test,  $P(x)$  is maximum when  $x = 75$

Maximum profit =  $P(75)$

$$= (150 - 75)75 - 1625$$

$$= 75 \times 75 - 1625$$

$$= 4000$$

Hence, the profit will be maximum, if the manufacturer manufactures 75 items and the maximum profit is 4000.

Question 19.

Find two numbers whose sum is 15 and when the square of one multiplied by the cube of the

other is maximum.

Solution:

Let the two numbers be  $x$  and  $y$ .

Then  $x + y = 15$

$\therefore y = 15 - x$

Let  $P$  is the product of square of  $y$  and cube of  $x$ .

Then  $P = x^3y^2$

$= x^3(15 - x)^2$

$= x^3(225 - 30x + x^2)$

$= x^5 - 30x^4 + 225x^3$

$\therefore \frac{dP}{dx} = \frac{d}{dx} (x^5 - 30x^4 + 225x^3)$

$= 5x^4 - 30 \times 4x^3 + 225 \times 3x^2$

$= 5x^4 - 120x^3 + 675x^2$

and  $\frac{d^2P}{dx^2} = \frac{d}{dx} (5x^4 - 120x^3 + 675x^2)$

$= 5 \times 4x^3 - 120 \times 3x^2 + 675 \times 2x$

$= 20x^3 - 360x^2 + 1350x$

$= 10x(2x^2 - 36x + 135)$

Now,  $\frac{dP}{dx} = 0$  gives  $5x^4 - 120x^3 + 675x^2 = 0$

$\therefore 5x^2(x^2 - 24x + 135) = 0$

$\therefore 5x^2(x^2 - 15x - 9x + 135) = 0$

$\therefore 5x^2[x(x - 15) - 9(x - 35)] = 0$

$\therefore 5x^2(x - 15)(x - 9) = 0$

$\therefore$  the roots of  $\frac{dP}{dx} = 0$  are  $x_1 = 0$ ,  $x_2 = 15$  and  $x_3 = 9$

If  $x = 0$ , then  $y = 15 - 0 = 15$

If  $x = 15$ , then  $y = 15 - 15 = 0$

In both cases, product  $x^3y^2$  is zero, which is not maximum.

$\therefore x \neq 0$  and  $x \neq 15$

$\therefore x = 6$

$$\text{Now, } \left( \frac{d^2P}{dx^2} \right)_{\text{at } x=6} = 10(6)[2(6)^2 - 36 \times 6 + 135]$$

$$= 60[72 - 216 + 135]$$

$$= 60(-9)$$

$$= -540 < 0$$

$\therefore P$  is maximum when  $x = 6$

If  $x = 6$ , then  $y = 15 - 6 = 9$

Hence, the required numbers are 6 and 9.

Question 20.

Show that among rectangles of given area, the square has least perimeter.

Solution:

Let  $x$  be the length and  $y$  be the breadth of the rectangle whose area is  $A$  sq units (which is given as constant).

Then  $xy = A$

$$\therefore y = \frac{A}{x} \dots\dots\dots(1)$$

Let  $P$  be the perimeter of the rectangle.

$$\text{Then } P = 2(x + y) = 2\left(x + \frac{A}{x}\right) \quad \dots \text{ [By (1)]}$$

$$\therefore \frac{dP}{dx} = 2 \cdot \frac{d}{dx}\left(x + \frac{A}{x}\right) = 2[1 + A(-1)x^{-2}]$$

$$= 2 \left( 1 - \frac{A}{x^2} \right)$$

$$\text{and } \frac{d^2P}{dx^2} = 2 \frac{d}{dx} \left( 1 - \frac{A}{x^2} \right) = 2 [0 - A(-2)x^{-3}]$$

$$= \frac{4A}{x^3}$$

$$\text{Now, } \frac{dP}{dx} = 0, \text{ gives } 2 \left( 1 - \frac{A}{x^2} \right) = 0$$

$$\therefore x^2 - A = 0 \quad \therefore x^2 = A$$

$$\therefore x = \sqrt{A} \quad \dots [\because x > 0]$$

$$\text{and } \left( \frac{d^2P}{dx^2} \right)_{\text{at } x = \sqrt{A}} = \frac{4A}{(\sqrt{A})^3} > 0$$

$$\therefore P \text{ is minimum when } x = \sqrt{A}$$

$$\text{If } x = \sqrt{A}, \text{ then } y = \frac{A}{x} = \frac{A}{\sqrt{A}} = \sqrt{A}$$

$$x = y$$

$\therefore$  rectangle is a square.

Hence, among rectangles of given area, the square has least perimeter.

Question 21.

Show that the height of a closed right circular cylinder of given volume and least surface area is equal to its diameter.

Solution:

Let  $x$  be the radius of base,  $h$  be the height and  $S$  be the surface area of the closed right circular

cylinder whose volume is  $V$  which is given to be constant.

$$\text{Then } \pi r^2 h = V$$

$$\therefore h = \frac{V}{\pi r^2} = \frac{A}{x^2} \dots\dots(1)$$

where  $A = \frac{V}{\pi}$ , which is constant.

$$\text{Now, } S = 2\pi xh + 2\pi x^2$$

$$= 2\pi x \left( \frac{A}{x^2} \right) + 2\pi x^2 \quad \dots \text{ [By (1)]}$$

$$= \frac{2\pi A}{x} + 2\pi x^2$$

$$\therefore \frac{dS}{dx} = \frac{d}{dx} \left( \frac{2\pi A}{x} + 2\pi x^2 \right)$$

$$= 2\pi A(-1)x^{-2} + 2\pi \times 2x$$

$$= \frac{-2\pi A}{x^2} + 4\pi x$$

$$\text{and } \frac{d^2S}{dx^2} = \frac{d}{dx} \left( \frac{-2\pi A}{x^2} + 4\pi x \right)$$

$$= -2\pi A(-2)x^{-3} + 4\pi \times 1$$

$$= \frac{4\pi A}{x^3} + 4\pi$$

$$\text{Now, } \frac{dS}{dx} = 0 \text{ gives } \frac{-2\pi A}{x^2} + 4\pi x = 0$$

$$\therefore -2\pi A + 4\pi x^3 = 0$$

$$\therefore 4\pi x^3 = 2\pi A$$

$$\therefore 4\pi x^3 = 2\pi A$$

$$\therefore x^3 = \frac{A}{2}$$

$$\therefore x = \left(\frac{A}{2}\right)^{\frac{1}{3}}$$

$$\text{and } \left(\frac{d^2S}{dx^2}\right)_{\text{at } x = \left(\frac{A}{2}\right)^{\frac{1}{3}}} = \frac{4\pi A}{\left(\frac{A}{2}\right)^{\frac{1}{3}}} + 4\pi = 12\pi > 0$$

$\therefore$  by the second derivative test,  $S$  is minimum when

$$x = \left(\frac{A}{2}\right)^{\frac{1}{3}}$$

When  $x = \left(\frac{A}{2}\right)^{\frac{1}{3}}$ , from (1),

$$h = \frac{A}{\left(\frac{A}{2}\right)^{\frac{2}{3}}} = 2^{\frac{2}{3}} \cdot A^{\frac{1}{3}} = 2 \cdot \left(\frac{A}{2}\right)^{\frac{1}{3}}$$

$$\therefore h = 2x$$

Hence, the surface area is least when height of the closed right circular cylinder is equal to its diameter.

Question 22.

Find the volume of the largest cylinder that can be inscribed in a sphere of radius 'r' cm.

Solution:

Let  $R$  be the radius and  $h$  be the height of the cylinder which is inscribed in a sphere of radius  $r$


cm.

Then from the figure,

$$R^2 + \left(\frac{h}{2}\right)^2 = r^2$$

$$\therefore R^2 = r^2 - \frac{h^2}{4} \quad \dots (1)$$

Let  $V$  be the volume of the cylinder.



Then  $V = \pi R^2 h$

$$= \pi \left( r^2 - \frac{h^2}{4} \right) h \quad \dots \text{ [By (1)]}$$

$$= \pi \left( r^2 h - \frac{h^3}{4} \right)$$

$$\therefore \frac{dV}{dh} = \pi \frac{d}{dh} \left( r^2 h - \frac{h^3}{4} \right)$$

$$= \pi \left( r^2 \times 1 - \frac{1}{4} \times 3h^2 \right) = \pi \left( r^2 - \frac{3}{4} h^2 \right)$$

$$\text{and } \frac{d^2V}{dh^2} = \pi \frac{d}{dh} \left( r^2 - \frac{3}{4} h^2 \right)$$

$$= \pi \left( 0 - \frac{3}{4} \times 2h \right) = -\frac{3}{2} \pi h$$

$$\text{Now, } \frac{dV}{dh} = 0 \text{ gives, } \pi \left( r^2 - \frac{3}{4} h^2 \right) = 0$$

$$\therefore r^2 - \frac{3}{4}h^2 = 0$$

$$\therefore \frac{3}{4}h^2 = r^2 \quad \therefore h^2 = \frac{4r^2}{3}$$

$$\therefore h = \frac{2r}{\sqrt{3}} \quad \dots [\because h > 0]$$

$$\text{and } \left( \frac{d^2V}{dh^2} \right)_{\text{at } h = \frac{2r}{\sqrt{3}}} = -\frac{3}{2}\pi \times \frac{2r}{\sqrt{3}} < 0$$

$$\therefore V \text{ is maximum at } h = \frac{2r}{\sqrt{3}}$$

If  $h = \frac{2r}{\sqrt{3}}$ , then from (1)

$$R^2 = r^2 - \frac{1}{4} \times \frac{4r^2}{3} = \frac{2r^2}{3}$$

$\therefore$  volume of the largest cylinder

$$= \pi \times \frac{2r^2}{3} \times \frac{2r}{\sqrt{3}} = \frac{4\pi r^3}{3\sqrt{3}} \text{ cu cm.}$$

Hence, the volume of the largest cylinder inscribed in a sphere of radius 'r' cm =  $\frac{4\pi r^3}{3\sqrt{3}}$  cu cm.

Question 23.

Show that  $y = \log(1+x) - \frac{2x}{2+x}$ ,  $x > -1$  is an increasing function on its domain.

Solution:

$$y = \log(1+x) - \frac{2x}{2+x}, x > -1$$

$$\therefore \frac{dy}{dx} = \frac{d}{dx} \left[ \log(1+x) - \frac{2x}{2+x} \right]$$

$$= \frac{1}{1+x} \cdot \frac{d}{dx}(1+x) - \frac{(2+x) \cdot \frac{d}{dx}(2x) - 2x \cdot \frac{d}{dx}(2+x)}{(2+x)^2}$$

$$= \frac{1}{1+x} \times (0+1) - \frac{(2+x) \times 2 - 2x(0+1)}{(2+x)^2}$$

$$= \frac{1}{1+x} - \frac{4+2x-2x}{(2+x)^2}$$

$$= \frac{1}{1+x} - \frac{4}{(2+x)^2}$$

$$= \frac{(2+x)^2 - 4(1+x)}{(1+x)(2+x)^2}$$

$$= \frac{4+4x+x^2-4-4x}{(1+x)(2+x)^2}$$

$$= \frac{x^2}{(1+x)(2+x)^2} > 0 \text{ for all } x > -1$$

Hence, the given function is increasing function on its domain.

Question 24.

Prove that  $y = \frac{4 \sin \theta}{2 + \cos \theta} - \theta$  is an increasing function if  $\theta \in [0, \frac{\pi}{2}]$

Solution:

---

$$\begin{aligned}y &= \frac{4 \sin \theta}{2 + \cos \theta} - \theta \\ \therefore \frac{dy}{d\theta} &= \frac{d}{d\theta} \left[ \frac{4 \sin \theta}{2 + \cos \theta} - \theta \right] \\ &= \frac{d}{d\theta} \left( \frac{4 \sin \theta}{2 + \cos \theta} \right) - \frac{d}{d\theta} (\theta) \\ &= \frac{(2 + \cos \theta) \cdot \frac{d}{d\theta} (4 \sin \theta) - 4 \sin \theta \cdot \frac{d}{d\theta} (2 + \cos \theta)}{(2 + \cos \theta)^2} - 1 \\ &= \frac{(2 + \cos \theta) (4 \cos \theta) - (4 \sin \theta) (0 - \sin \theta)}{(2 + \cos \theta)^2} - 1 \\ &= \frac{8 \cos \theta + 4 \cos^2 \theta + 4 \sin^2 \theta}{(2 + \cos \theta)^2} - 1 \\ &= \frac{8 \cos \theta + 4(\cos^2 \theta + \sin^2 \theta)}{(2 + \cos \theta)^2} - 1 \\ &= \frac{8 \cos \theta + 4}{(2 + \cos \theta)^2} - 1 \\ &= \frac{(8 \cos \theta + 4) - (2 + \cos \theta)^2}{(2 + \cos \theta)^2} \\ &= \frac{8 \cos \theta + 4 - 4 - 4 \cos \theta - \cos^2 \theta}{(2 + \cos \theta)^2} \\ &= \frac{4 \cos \theta - \cos^2 \theta}{(2 + \cos \theta)^2} = \cos \theta (4 - \cos \theta)\end{aligned}$$

---

$$= \frac{4 \cos \theta - \cos^2 \theta}{(2 + \cos \theta)^2} = \frac{\cos \theta (4 - \cos \theta)}{(2 + \cos \theta)^2}$$

Since,  $\theta \in \left[ 0, \frac{\pi}{2} \right]$ ,  $\cos \theta \geq 0$ , Also,  $\cos \theta < 4$

$$\therefore 4 - \cos \theta > 0$$

$$\therefore \cos \theta (4 - \cos \theta) \geq 0$$

$$\therefore \frac{\cos \theta (4 - \cos \theta)}{(2 + \cos \theta)^2} \geq 0$$

$$\therefore \frac{dy}{d\theta} \geq 0 \text{ for all } \theta \in \left[ 0, \frac{\pi}{2} \right]$$

Hence,  $y$  is an increasing function if  $\theta \in \left[ 0, \frac{\pi}{2} \right]$ .



# Maharashtra Board Solutions

## Class 12 Arts & Science Maths

### (Part 2)

- Chapter 1- Differentiation
- Chapter 2- Applications of Derivatives
- Chapter 3- Indefinite Integration
- Chapter 4- Definite Integration
- Chapter 5- Application of Definite Integration
- Chapter 6- Differential Equations
- Chapter 7- Probability Distributions
- Chapter 8- Binomial Distribution

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>

## About About Maharashtra State Board (MSBSHSE)

The Maharashtra State Board of Secondary and Higher Secondary Education or MSBSHSE (Marathi: महाराष्ट्र राज्य माध्यमिक आणि उच्च माध्यमिक शिक्षण मंडळ), is an **autonomous and statutory body established in 1965**. The board was amended in the year 1977 under the provisions of the Maharashtra Act No. 41 of 1965.

The Maharashtra State Board of Secondary & Higher Secondary Education (MSBSHSE), Pune is an independent body of the Maharashtra Government. There are more than 1.4 million students that appear in the examination every year. The Maha State Board conducts the board examination twice a year. This board conducts the examination for SSC and HSC.

The Maharashtra government established the Maharashtra State Bureau of Textbook Production and Curriculum Research, also commonly referred to as Ebalbharati, in 1967 to take up the responsibility of providing quality textbooks to students from all classes studying under the Maharashtra State Board. MSBHSE prepares and updates the curriculum to provide holistic development for students. It is designed to tackle the difficulty in understanding the concepts with simple language with simple illustrations. Every year around 10 lakh students are enrolled in schools that are affiliated with the Maharashtra State Board.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>

## FAQs

Where do I get the Maharashtra State Board Books PDF For free download?

You can download the Maharashtra State Board Books from the eBalbharti official website, i.e. [cart.ebalbharati.in](http://cart.ebalbharati.in) or from this article.

Add image

How to Download Maharashtra State Board Books?

Students can get the Maharashtra Books for primary, secondary, and senior secondary classes from here. You can view or download the Maharashtra State Board Books from this page or from the official website for free of cost. Students can follow the detailed steps below to visit the official website and download the e-books for all subjects or a specific subject in different mediums.

Step 1: Visit the official website [ebalbharati.in](http://ebalbharati.in)

Step 2: On the top of the screen, select "Download PDF textbooks"

Step 3: From the "Classes" section, select your class.

Step 4: From "Medium", select the medium suitable to you.

Step 5: All Maharashtra board books for class 11th will now be displayed on the right side.

Step 6: Click on the "Download" option to download the PDF book.

Who developed the Maharashtra State board books?

As of now, the MSCERT and Balbharti are responsible for the syllabus and textbooks of Classes 1 to 8, while Classes 9 and 10 are under the Maharashtra State Board of Secondary and Higher Secondary Education (MSBSHSE).

How many state boards are there in Maharashtra?

The Maharashtra State Board of Secondary & Higher Secondary Education, conducts the HSC and SSC Examinations in the state of Maharashtra through its nine Divisional Boards located at Pune, Mumbai, Aurangabad, Nasik, Kolhapur, Amravati, Latur, Nagpur and Ratnagiri.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-part-2-chapter-2-applications-of-derivatives/>

## About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to [info@indcareer.com](mailto:info@indcareer.com)

### Postal Address

IndCareer.com  
52, Shilpa Nagar,  
Somalwada  
Nagpur - 440015  
Maharashtra, India

**WhatsApp:** +91 9561 204 888

**Website:** <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-arts-science-maths-p-art-2-chapter-2-applications-of-derivatives/>