


Maharashtra Board Solutions Class 12

Chemistry: Chapter 10- Halogen Derivatives

Class 12 - Chapter 10 Halogen Derivatives



For any clarifications or questions you can write to info@indcareer.com

Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888, **Website:** <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



Maharashtra Board Solutions Class 12

Chemistry: Chapter 10- Halogen Derivatives

Class 12: Chemistry Chapter 10 solutions. Complete Class 12 Chemistry Chapter 10 Notes.


Maharashtra Board Solutions Class 12 Chemistry: Chapter 10- Halogen Derivatives

Maharashtra Board 12th Chemistry Chapter 10, Class 12 Chemistry Chapter 10 solutions

1. Choose the most correct option.

Question i.

The correct order of increasing reactivity of C-X bond towards nucleophile in the following compounds is



a. I < II < III < IV

b. II < I < III < IV

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>


c. $\text{III} < \text{IV} < \text{II} < \text{I}$

d. $\text{IV} < \text{III} < \text{I} < \text{II}$

Answer:

(d) $\text{IV} < \text{III} < \text{I} < \text{II}$

Question ii.



The major product of the above reaction is,

a. $\text{I-CH}_2\text{-CH=CH}_2$ b. $\text{CH}_3\text{-CH}_2\text{-CH}_2\text{I}$

c. $\text{CH}_3\text{-}\underset{\text{I}}{\text{CH}}\text{-CH}_3$ d. $\text{CH}_3\text{-}\underset{\text{I}}{\text{CH}}\text{-}\underset{\text{OH}}{\text{CH}_2}$





Answer:

(c)

Question iii.

Which of the following is likely to undergo racemization during alkaline hydrolysis?

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



- a. Only I b. Only II
c. II and IV d. Only IV

Answer:

(a) Only I

Question iv.

The best method for preparation of alkyl fluorides is

- a. Finkelstein reaction
b. Swartz reaction
c. Free radical fluorination
d. Sandmeyer's reaction

Answer:

b. Swartz reaction

Question v.

Identify the chiral molecule from the following.

- a. 1-Bromobutane

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

- b. 1,1- Dibromobutane
- c. 2,3- Dibromobutane
- d. 2-Bromobutane

Answer:

(d) 2-Bromobutane

Question vi.

An alkyl chloride on Wurtz reaction gives 2,2,5,5-tetramethylhexane. The same alkyl chloride on reduction with zinc-copper couple in alcohol give hydrocarbon with molecular formula C_5H_{12} . What is the structure of alkyl chloride

- a. $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3-\text{C}-\text{CH}_2\text{Cl} \\ | \\ \text{CH}_3 \end{array}$ b. $\begin{array}{c} \text{CH}_3 \\ | \\ \text{CH}_3-\text{C}-\text{CH}_2\text{CH}_3 \\ | \\ \text{Cl} \end{array}$
- c. $\begin{array}{c} \text{CH}_3-\text{CH}_2-\text{CH}-\text{Cl} \\ | \\ \text{CH}_3 \end{array}$ d. $\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}-\text{CHCl} \\ | \quad | \\ \text{CH}_3 \quad \text{CH}_3 \end{array}$

Answer:

(a)

Question vii.

Butanenitrile may be prepared by heating

- a. propanol with KCN
- b. butanol with KCN
- c. n-butyl chloride with KCN
- d. n-propyl chloride with KCN

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Answer:

(d) n-propyl chloride with KCN

Question viii.


Choose the compound from the following that will react fastest by S_N1 mechanism.

- a. 1-iodobutane
- b. 1-iodopropane
- c. 2-iodo-2 methylbutane
- d. 2-iodo-3-methylbutane




Answer:

(c) 2-iodo-2 methylbutane

Question ix.



The product 'B' in the above reaction sequence is,

- a.  b. 
- c.  d. 

Answer:

(d)

Question x.

Which of the following is used as source of dichlorocarbene

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

- a. tetrachloromethane
- b. chloroform
- c. iodoform
- d. DDT





Answer:

(b) chloroform

2. Do as directed.


Question i.

Write IUPAC name of the following compounds




Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Compound	Ans. IUPAC names
(1) $\text{CH}_3 - \text{CH} = \underset{\text{CH}_3}{\text{C}} - \underset{\text{CH}_3}{\text{CH}} - \text{Br}$	4-Bromo-3, 4 dimethyl but-2-ene
(2) $\text{CH}_3 - \underset{\text{Cl}}{\text{CH}} - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_2 - \text{CH}_3$	2-Chloro-3-methyl pentane
(3) 	1-Chloro-4-ethyl cyclohexane
(4) 	1,4-Dichloro-2-methyl benzene

Question ii.






Write structure and IUPAC name of the major product in each of the following reaction.



Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Structure and IUPAC name



Question iii.


Identify chiral molecule/s from the following.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

- a. $\text{CH}_3-\underset{\text{OH}}{\text{CH}}-\text{CH}_2-\text{CH}_3$
- b. $\text{CH}_3-\text{CH}_2-\underset{\text{Br}}{\text{CH}}-\text{CH}_2-\text{CH}_3$
- c. $\text{CH}_3-\text{CH}_2-\text{CH}_2-\text{CH}_2\text{Br}$
- d. $\text{CH}_3-\underset{\text{CH}_3}{\text{CH}}-\text{CH}_3-\text{CH}_3$



Answer:

Chiral molecule




Question iv.

Which one compound from the following pairs would undergo $\text{S}_\text{N}2$ faster from the?




Answer:

(1) Sincey



<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>





is a primary halide it undergoes S_N2 reaction faster than



(2) Since iodine is a better leaving group than chloride, 1-iodo propane ($\text{CH}_3\text{CH}_2\text{CH}_2\text{I}$) undergoes S_N2 reaction faster than 1-chloropropane ($\text{CH}_3\text{CH}_2\text{CH}_2\text{Cl}$).






Question v.

Complete the following reactions giving major product.




Answer:




<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>





Answer:



Answer:



Answer:



Question vi.

Name the reagent used to bring about the following conversions.

- Bromoethane to ethoxyethane
- 1-Chloropropane to 1 nitropropane
- Ethyl bromide to ethyl isocyanide

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

d. Chlorobenzene to biphenyl

Answer:

Reactant	Ans.	Reagent	Product
(1) Bromoethane		$\xrightarrow[\text{Sodium alkoxide}]{\Delta}$	ethoxy ethane
(2) 1-Chloropropane		$\xrightarrow[\text{alcohol}]{\text{Silver nitrite}}$	1-nitropropane
(3) Ethyl bromide		$\xrightarrow[\text{AgCN}]{\text{Alcohol}}$	Ethyl isocyanide
(4) Chlorobenzene		$\xrightarrow[\text{dry ether}]{\text{Na}}$	Biphenyl
(5) Ethyl iodide		$\xrightarrow[\text{dry ether}]{\text{Mg}}$	Ethyl magnesium iodide
(6) Ethyl chloride		$\xrightarrow[\text{acetone}]{\text{NaI}}$	Ethyl iodide
(7) Toluene + Br ₂		$\xrightarrow[\text{dark}]{\text{Fe}}$	ortho and para bromo toluene

Question vii.

Arrange the following in the increase order of boiling points

- 1-Bromopropane
- 2- Bromopropane
- 1- Bromobutane
- 1-Bromo-2-methylpropane




Answer:

1-Bromo-2-methylpropane, 2-Bromopropane, 1-Bromopropane, 1-Bromo butane

Question viii.

Match the pairs.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Column I**Column II**




i. vinyl halide

ii. alkyl halide

iii. allyl halide

iv. benzyl halide


v. aryl halide

Answer:**3. Give reasons****Question i.****Haloarenes are less reactive than haloalkanes.****Answer:**

Haloarenes (Aryl halides) are less reactive than (alkyl halides) haloalkanes due to the following reasons :

(1) Resonance effect : In haloarenes, the electron pairs on halogen atom are in conjugation with π -electrons of the benzene ring. The delocalization of these electrons C-Cl bond acquires partial double bond character.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



Due to partial double bond character of C-Cl bond in aryl halides, the bond cleavage in haloarene is difficult and are less reactive. On the other hand, in alkyl halides, carbon is attached to chlorine by a single bond and it can be easily broken.

(2) Aryl halides are stabilized by resonance but alkyl halides are not. Hence, the energy of activation for the displacement of halogen from aryl halides is much greater than that of alkyl halides.

(3) Different hybridization state of carbon atom in C-X bond :

(i) In alkyl halides, the carbon of C-X bond is sp^3 -hybridized with less s-character and greater bond length of 178 pm, which requires less energy to break the C-X bond.

(ii) In aryl halides, the carbon of C-X bond is sp^2 -hybridized with more s-character and shorter bond length which requires more energy to break C-X bond. Therefore, aryl halides are less reactive than alkyl halides.


(iii) Polarity of the C-X bond : In aryl halide C-X bond is less polar than in alkyl halides. Because sp^2 -hybrid carbon of C-X bond has less tendency to release electrons to the halogen than a sp^3 -hybrid carbon in alkyl halides. Thus halogen atom in aryl halides cannot be easily displaced by nucleophile.

(2) Aryl halides are extremely less reactive towards nucleophilic substitution reactions.

Answer:

Aryl halides are extremely less reactive towards nucleophilic substitution reaction due to the following reasons : (1) Resonance effect : In haloarenes, the electron pairs on halogen atom are in conjugation with π -electrons of the benzene ring. The delocalization of these electrons C-Cl bond acquires partial double bond character.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>




Due to partial double bond character of C-Cl bond in aryl halides, the bond cleavage in haloarene is difficult and are less reactive towards nucleophilic substitution.

(2) sp^2 hybrid state of C : Different hybridization state of carbon atom in C-X bond : In aryl halides, the carbon of C-X bond is sp^2 -hybridized with more s-character and shorter bond length of 169 pm which requires more energy to break C-X bond. It is difficult to break a shorter bond than a longer bond, in alkyl chloride (bond length 178 pm) therefore, aryl halides are less reactive towards nucleophilic substitution reaction.

(3) Instability of phenyl cation : In aryl halides, the phenyl cation formed due to self ionisation will not be stabilized by resonance which rules out possibility of S_N1 mechanism. Also backside attack of nucleophile is blocked by the aromatic ring which rules out S_N2 mechanism. Thus cations are not formed and hence aryl halides do not undergo nucleophilic substitution reaction easily.

(4) As aryl halides are electron rich molecules due to the presence of re-bond, they repel electron rich nucleophilic, attack. Hence, aryl halides are less reactive towards nucleophilic substitution reactions. However, the presence of electron withdrawing groups at o/p position activates the halogen of aryl halides towards substitution.



(3) Aryl halides undergo electrophilic substitution reactions slowly.


Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>


Aryl halides undergo electrophilic substitution reactions slowly and it can be explained as follows :

(1) Inductive effect : The strongly electronegative halogen atom withdraws the electrons from carbon, atom of the ring, hence aryl halides show reactivity towards electrophilic attack.

(2) Resonance effect : The resonating structures of aryl halides show increase in electron density at ortho and para position, hence it is o, p directing.



The inductive effect and resonance effect compete with each other. The inductive effect is stronger than resonance effect. The reactivity of aryl halides is controlled by stronger inductive effect and o, p orientation is controlled by weaker resonating effect.



The attack of electrophile (Y) on haloarenes at ortho and para positions are more stable due to formation of chloronium ion. The chloronium ion formed is comparatively more stable than other hybrid structures of carbonium ion.

(4) Reactions involving Grignard reagent must be carried out under anhydrous condition.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

(1) Grignard reagent ($R\ Mg\ X$) is an organometallic compound. The carbon-magnesium bond is highly polar and magnesium halogen bond is ionic in nature. Grignard reagent is highly reactive.

(2) The reactions of Grignard reagent are carried out in dry conditions because traces of moisture may spoil the reaction and Grignard reagent reacts with water to produce alkane. Hence, reactions involving Grignard reagent must be carried out under anhydrous condition.

(5) Alkyl halides are generally not prepared by free radical halogenation of alkane.

Answer:

(1) Free radical halogenation of alkane gives a mixture of all different possible Monohaloalkanes as well as polyhalogen alkanes.

(2) In this method, by changing the quantity of halogen the desired product can be made to predominate over the other

products. Hence, alkyl halides are generally not prepared by free radical halogenation of alkane.

Question ii.

Alkyl halides though polar are immiscible with water.

Answer:

(1) In alkyl halide, the halogen atom is more electronegative than carbon atom, the $C - X$ bond is polar.

(2) Though alkyl halide is polar, it is insoluble in water because alkyl halide is not able to form hydrogen bonds with water. Attraction between alkyl halide molecule is stronger than attraction between alkyl halide and water.

(2) $C-F$ bond in CH_3F is the strongest bond and $C-I$ bond in CH_3I is the weakest bond. Explain.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

(1) Methyl fluoride (CH_3F) is highly polar molecule and has the shortest C-F bond length (139 pm) and the strongest C-F bond due to greater overlap of orbitals of the same principal quantum number i.e., overlap of 2sp^3 orbital of carbon with 2p_z orbital of fluorine.

(2) Methyl iodide (CH_3I) is much less polar and has the longest (C-I) bond length (214 pm) and the weakest C-I bond due to poor overlap of 2sp^3 orbital carbon with 5p_z orbital of iodine i.e., 2sp^3 orbital of carbon cannot penetrate into larger p-orbitals.

(3) The boiling point of alkyl iodide is higher than that of alkyl fluoride.

Answer:


For a given alkyl group, the boiling point increases with increasing atomic mass of the halogen, because magnitude of van der Waals force increases with increase in size and mass of halogen. Therefore, boiling point of alkyl iodide is higher than that of alkyl fluoride.

(4) The boiling point of isopropyl bromide is lower than that of n-propyl bromide.

Answer:

For isomeric alkyl halides (isopropyl bromide and n-propyl bromide), the boiling point decreases as the branching increases, surface area decreases on branching and van der Waals forces decrease, therefore, the boiling point of isopropyl bromide is lower than that of n-propyl bromide.

(5) p-Dichlorobenzene



has mp. higher than those of o-and m-isomers.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

p-Dichlorobenzene has higher melting point than those of o-and m-isomers. This is because of its symmetrical structure which can easily fits in crystal lattice. As a result intermolecular forces of attraction are stronger and therefore greater energy is required to overcome its lattice energy.

Question iii.

Reactions involving Grignard reagent must be carried out under anhydrous conditions.

Question iv.

Alkyl halides are generally not prepared by free radical halogenation of alkanes.

Answer:

(1) Direct fluorination of alkanes is highly exothermic, explosive and invariably leads to polyfluorination and decomposition of the alkanes. It is difficult to control the reaction.

(2) Direct iodination of alkanes is highly reversible and difficult to carry out.

(3) In direct chlorination and bromination, the reaction is not selective. It can lead to different isomeric monohalogenated alkanes (alkyl halides) as well as polyhalogenated alkanes.

Hence, halogenation of alkanes is not a good method of preparation of alkyl halides.

4. Distinguish between – S_N1 and S_N2 mechanism of substitution reaction ?

Answer:


<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Factor	S _N 1	S _N 2
(1) Number of steps	Two steps MaharashtraBoardSolutions.Guru	One step
(2) Molecularity/Order	Unimolecular/1st order	Bimolecular/2nd order
(3) Reaction rate	Depends upon concentration of one reacting species	Depends upon concentration of two reacting species
(4) Attack of a nucleophile	Back side attack and front side attack on a substrate with equal probability	Only back side attack on a substrate
(5) Transition state	Two steps, two transition states	One step, one transition state
(6) Type of substrate	Mainly tertiary (3°) substrate	Mainly primary (1°) substrate
(7) Stereochemistry	50% inversion and 50% retention of configuration	100% inversion of configuration
(8) Enantiomer	Forms racemic mixture	Forms opposite enantiomer
(9) Solvent	Polar solvent favourable	Nonpolar solvent favourable
(10) Energy of activation	Two values of energies of activation	One value of energy activation
(11) Intermediate	Carbocation intermediates	No intermediate
(12) Nucleophile	Weak nucleophile favourable	Strong nucleophile favourable
(13) Order of reactivity in alkyl halides	Tertiary > Secondary > Primary	Primary > Secondary > Tertiary

5. Explain – Optical isomerism in 2-chlorobutane.


Answer:

(1) 2-Chlorobutane contains an asymmetric.



carbon atom (the starred carbon atom) which is attached to four different groups, i.e., ethyl (-CH₂ – CH₃), methyl (CH₃), chloro (Cl) and hydrogen (H) groups.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



**Fig. Enantiomers of 2-Chlorobutane
(with structural and Fischer's projection formula)**

(2) Two different arrangements of these groups around the carbon atom are possible as shown in the figure. Hence, it exists as a pair of enantiomers. The two enantiomers are mirror images of each other and are not superimposable.

(3) One of the enantiomers will rotate the plane of plane-polarized light to the left hand side and is called the laevorotatory isomer (*l*-isomer). The other enantiomer will rotate the plane of plane-polarized light to the right hand side and is called the dextrorotatory isomer (*d*-isomer).

(4) Equimolar mixture of the *d*- and the *l*-isomers is optically inactive and is called the racemic mixture or the racemate (*dl*-mixture). The optical inactivity of the racemic mixture is due to external compensation.


6. Convert the following.

Question i.

Propene to propan-1-ol


Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



Propene

n-propyl bromide




Propan-1-ol

Question ii.

Benzyl alcohol to benzyl cyanide

Answer:



Question iii.

Ethanol to propane nitrile


Answer:



Ethanol

(48%)

Ethyl bromide



Propane nitrile

Question iv.


But-1-ene to n-butyl iodide

Answer:




But-1-ene

Tributyl borane



Butan-1-ol




n-butyl iodide

Question v.


2-Chloropropane to propan-1-ol

Answer:



Propene

tripropyl borane




Propan-1-ol

Question vi.


tert-Butyl bromide to isobutyl bromide

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



tert. butyl bromide



Question vii.

Aniline to chlorobenzene


Answer:



Question viii.

Propene to 1-nitropropane

Answer:




7. Answer the following

Question i.



HCl is added to a hydrocarbon 'A' (C_4H_8) to give a compound 'B' which on hydrolysis with aqueous alkali forms tertiary alcohol 'C' ($C_4H_{10}O$). Identify 'A', 'B' and 'C'.

Answer:



2-methyl propene


tert.butyl chloride











Question ii.

Complete the following reaction sequences by writing the structural formulae of the organic compounds 'A', 'B' and 'C'.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>




Answer:



Question iii.

Observe the following and answer the questions given below.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>



- Name the type of halogen derivative
- Comment on the bond length of C-X bond in it
- Can react by $\text{S}_{\text{N}}1$ mechanism? Justify your answer.

Answer:

- Vinyl halide
- C – X bond length shorter in vinyl halide than alkyl halide. Vinyl halide has partial double bond character due to resonance.

In vinyl halide, carbon is sp hybridised. The bond is shorter and stronger and the molecule is more stable.

- Yes, It reacts by $\text{S}_{\text{N}}1$ mechanism. $\text{S}_{\text{N}}1$ mechanism involves formation of carbocation intermediate. The vinylic carbocation intermediate formed is resonance stabilized, hence $\text{S}_{\text{N}}1$ mechanism is favoured.

Activity :


- Collect detailed information about Freons and their uses.
- Collect information about DDT as a persistent pesticide.

Reference books

- Organic chemistry by Morrison, Boyd, Bhattacharjee, 7th edition, Pearson
- Organic chemistry by Finar, Vol 1, 6th edition, Pearson


Use your brain power..... (Textbook page 212)

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>





Question 1.

Write IUPAC names of the following:




Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

- (i) $\text{CH}_3 - \underset{\text{Br}}{\text{CH}} - \text{CH}_3$ 2-Bromopropane
- (ii) $\text{CH}_3 - \underset{\text{CH}_3}{\text{CH}} - \text{CH}_2\text{I}$ 1-Iodo-2-methyl propane
- (iii) $\text{CH}_3 - \text{CH} = \text{CH} - \text{CH}_2 - \text{Cl}$ 1-Chloro-but-2-ene
- (iv) $\text{CH}_3 - \text{C} \equiv \text{C} - \text{CH}_2 - \text{Br}$ 1-Bromo-but-2-yne
- (v)  Bromobenzene
- (vi)  1, 4-Dibromobenzene (*p*-dibromobenzene)

Question 10.1 : (Textbook page 213)


How will you obtain 1.bromo.1-methylcyclohexane from alkene? Write possible structures of alkene and the reaction involved.




Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Rewrite the following reaction by filling the blanks:



(major) (minor)






(major) (minor)



(major) (minor)

Answer:



Question 10.2 : (Textbook page 216)

Arrange the following compounds in order of increasing boiling points :
bromoform, chloromethane, dibromomethane, bromomethane.

Answer:

The comparative boiling points of halogen derivatives are mainly related with van der Waals forces of attraction which depend upon the molecular size. In the present case all the compounds contain only one carbon. Thus the molecular size depends upon the size of halogen and number of halogen atoms present.

Thus increasing order of boiling point is, $\text{CH}_3\text{Cl} < \text{CH}_3\text{Br} < \text{CH}_2\text{Br}_2 < \text{CHBr}_3$

Try this (Textbook page 2016)

Question 1.

- (1) Make a three-dimensional model of 2-chlorobutane.
- (2) Make another model which is a mirror image of the first model.
- (3) Try to superimpose the two models on each other.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

(4) Do they superimpose on each other exactly ?

(5) Comment on whether the two models are identical or not.

Answer:

(1) (2) and (3)




Fig : Nonsuperimposable mirror images

(4) Two models are non-superimposable mirror images of each other called enantiomers.


(5) Two enantiomers are identical. They have the same physical properties (such as melting points, boiling points, densities, refractive index). They also have identical chemical properties. The magnitude of their optical rotation is equal but the sign of optical rotation is opposite.

Try this (Textbook page 219)

Question 1.

1. Draw structures of enantiomers of lactic acid

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>





using Fischer projection formulae.


2. Draw structures of enantiomers of 2-bromobutane using wedge formula.

Answer:

(1)



(2) Wedge formula : 2-bromobutane



Can you tell? (Textbook page 220)


Question 1.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Alkyl halides, when treated with alcoholic solution of silver nitrite, give nitroalkanes whereas with sodium nitrite they give alkyl nitrites. Explain.


Answer:

Nitrite ion is an ambident nucleophile, which can attack through 'O' or 'N'.



Both nitrogen and oxygen are capable of donating electron pair. C – N bond, being stronger than N – O bond, attack occurs through C atom from alkyl halide forming nitroalkane.


However, sodium nitrite (NaNO_2) is an ionic compound and oxygen is free to donate pair of electrons. Hence, attack occurs through oxygen resulting in the formation of alkyl nitrite.



Use your brain power! (Textbook page 222)


Question 1.

Draw the Fischer projection formulae of two products obtained when compound (A) reacts with OH^- by $\text{S}_{\text{N}}1$ mechanism.




Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>




Question 2.

Draw the Fischer projection formula of the product formed when compound (B) reacts with $\text{OH}^\ominus$ by $\text{S}_\text{N}2$ mechanism.



Answer:




Question 10.4 : (Textbook page 223)

Allylic and benzylic halides show high reactivity towards the $\text{S}_\text{N}1$ mechanism than other primary alkyl halides. Explain.


Answer:

In allylic and benzylic halide, the carbocation formed undergoes stabilization through the resonance. Hence, allylic and benzylic halides show high reactivity towards the $\text{S}_\text{N}1$ reaction. The resonating structures are

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>




Resonance stabilization of allylic carbocation



Resonance stabilization of benzylic carbocation

Question 10.5 : (Textbook page 224)

Which of the following two compounds would react faster by $\text{S}_{\text{N}}2$ mechanism and Why?



Answer :

In $\text{S}_{\text{N}}2$ mechanism, a pentacoordinate T.S. is involved. The order of reactivity of alkyl halides towards $\text{S}_{\text{N}}2$ mechanism is.

Primary > Secondary > Tertiary, (due to increasing crowding in T.S. from primary to tertiary halides.


1- Chlorobutane being primary halide will react faster by $\text{S}_{\text{N}}2$ mechanism, than the secondary halide 2- chlorobutane.)

Can you tell? (Textbook page 227)

Question 1.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

Conversion of chlorobenzene to phenol by aqueous sodium hydroxide requires high temperature of about 623K and high pressure. Explain.



Answer:

Due to the partial double bond character in chlorobenzene, the bond cleavage in chlorobenzene is difficult and is less reactive. Hence, during the conversion of chlorobenzene to phenol by a question NaOH requires high temperature & high pressure.



Maharashtra Board Solutions

Class 12 Chemistry

- Chapter 1- Solid State
- Chapter 2- Solutions
- Chapter 3- Ionic Equilibria
- Chapter 4- Chemical Thermodynamics
- Chapter 5- Electrochemistry
- Chapter 6- Chemical Kinetics
- Chapter 7- Elements of Groups 16, 17 and 18
- Chapter 8- Transition and Inner Transition Elements
- Chapter 9- Coordination Compounds
- Chapter 10- Halogen Derivatives
- Chapter 11- Alcohols, Phenols and Ethers
- Chapter 12- Aldehydes, Ketones and Carboxylic Acids
- Chapter 13- Amines
- Chapter 14- Biomolecules
- Chapter 15- Introduction to Polymer Chemistry
- Chapter 16- Green Chemistry and Nanochemistry

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

About About Maharashtra State Board (MSBSHSE)

The Maharashtra State Board of Secondary and Higher Secondary Education or MSBSHSE (Marathi: महाराष्ट्र राज्य माध्यमिक आणि उच्च माध्यमिक शिक्षण मंडळ), is an **autonomous and statutory body established in 1965**. The board was amended in the year 1977 under the provisions of the Maharashtra Act No. 41 of 1965.

The Maharashtra State Board of Secondary & Higher Secondary Education (MSBSHSE), Pune is an independent body of the Maharashtra Government. There are more than 1.4 million students that appear in the examination every year. The Maha State Board conducts the board examination twice a year. This board conducts the examination for SSC and HSC.

The Maharashtra government established the Maharashtra State Bureau of Textbook Production and Curriculum Research, also commonly referred to as Ebalbharati, in 1967 to take up the responsibility of providing quality textbooks to students from all classes studying under the Maharashtra State Board. MSBHSE prepares and updates the curriculum to provide holistic development for students. It is designed to tackle the difficulty in understanding the concepts with simple language with simple illustrations. Every year around 10 lakh students are enrolled in schools that are affiliated with the Maharashtra State Board.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

FAQs

Where do I get the Maharashtra State Board Books PDF For free download?

You can download the Maharashtra State Board Books from the eBalbharti official website, i.e. cart.ebalbharati.in or from this article.

Add image

How to Download Maharashtra State Board Books?

Students can get the Maharashtra Books for primary, secondary, and senior secondary classes from here. You can view or download the Maharashtra State Board Books from this page or from the official website for free of cost. Students can follow the detailed steps below to visit the official website and download the e-books for all subjects or a specific subject in different mediums.

Step 1: Visit the official website ebalbharati.in

Step 2: On the top of the screen, select "Download PDF textbooks"

Step 3: From the "Classes" section, select your class.

Step 4: From "Medium", select the medium suitable to you.

Step 5: All Maharashtra board books for class 11th will now be displayed on the right side.

Step 6: Click on the "Download" option to download the PDF book.

Who developed the Maharashtra State board books?

As of now, the MSCERT and Balbharti are responsible for the syllabus and textbooks of Classes 1 to 8, while Classes 9 and 10 are under the Maharashtra State Board of Secondary and Higher Secondary Education (MSBSHSE).

How many state boards are there in Maharashtra?

The Maharashtra State Board of Secondary & Higher Secondary Education, conducts the HSC and SSC Examinations in the state of Maharashtra through its nine Divisional Boards located at Pune, Mumbai, Aurangabad, Nasik, Kolhapur, Amravati, Latur, Nagpur and Ratnagiri.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>

About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to **info@indcareer.com**

Postal Address

IndCareer.com
52, Shilpa Nagar,
Somalwada
Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888

Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-10-halogen-derivatives/>