


Maharashtra Board Solutions Class 12

Chemistry: Chapter 3- Ionic Equilibria

Class 12 - Chapter 3 Ionic Equilibria



For any clarifications or questions you can write to info@indcareer.com

Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888, **Website:** <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>



Maharashtra Board Solutions Class 12

Chemistry: Chapter 3- Ionic Equilibria

Class 12: Chemistry Chapter 3 solutions. Complete Class 12 Chemistry Chapter 3 Notes.

Maharashtra Board Solutions Class 12 Chemistry: Chapter 3- Ionic Equilibria

Maharashtra Board 12th Chemistry Chapter 3, Class 12 Chemistry Chapter 3 solutions

1. Choose the most correct answer :

Question i.

The pH of 10^{-8} M of HCl is

- (a) 8
- (b) 7
- (c) less than 7
- (d) greater than 7

Answer:

- (c) less than 7

Question ii.

Which of the following solution will have pH value equal to 1.0?

- (a) 50 mL of 0.1M HCl + 50mL of 0.1 M NaOH

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

(b) 60 mL of 0.1M HCl + 40mL of 0.1 M NaOH

(c) 20 mL of 0.1M HCl + 80mL of 0.1 M NaOH

(d) 75 mL of 0.2M HCl + 25mL of 0.2 M NaOH

Answer:

(d) 75 mL of 0.2M HCl + 25mL of 0.2 M NaOH

Question iii.

Which of the following is a buffer solution ?

(a) $\text{CH}_3\text{COONa} + \text{NaCl}$ in water

(b) $\text{CH}_3\text{COOH} + \text{HCl}$ in water

(c) $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$ in water

(d) $\text{HCl} + \text{NH}_4\text{Cl}$ in water

Answer:

(c) $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$ in water

Question iv.

The solubility product of a sparingly soluble salt AX is 5.2×10^{-13} . Its solubility in mol dm^{-3} is

(a) 7.2×10^{-7}

(b) 1.35×10^{-4}

(c) 7.2×10^{-8}

(d) 13.5×10^{-8}

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

(a) 7.2×10^{-7}

Question v.

Blood in human body is highly buffered at pH of

(a) 7.4

(b) 7.0

(c) 6.9

(d) 8.1

Answer:

(a) 7.4

Question vi.

The conjugate base of $[\text{Zn}(\text{H}_2\text{O})_4]^{2+}$ is

(a) $[\text{Zn}(\text{H}_2\text{O})_4]^{2+} \text{NH}_3$

(b) $[\text{Zn}(\text{H}_2\text{O})_3]^{2+}$

(c) $[\text{Zn}(\text{H}_2\text{O})_3\text{OH}]^+$

(d) $[\text{Zn}(\text{H}_2\text{O})\text{H}]^{3+}$

Answer:

(c) $[\text{Zn}(\text{H}_2\text{O})_3\text{OH}]^+$

Question vii.

For $\text{pH} > 7$ the hydronium ion concentration would be

(a) 10^{-7} M

(b) $< 10^{-7} \text{ M}$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

(c) $> 10^{-7} \text{ M}$

(d) $\geq 10^{-7} \text{ M}$

Answer:

(b) $< 10^{-7} \text{ M}$

2. Answer the following in one sentence :

Question i.

Why cations are Lewis acids ?

Answer:

Since cations are deficient of electrons they accept a pair of electrons, hence they are Lewis acids.

Question ii.

Why is KCl solution neutral to litmus?

Answer:

1. Since KCl is a salt of strong base KOH and strong acid HCl, it does not undergo hydrolysis in its aqueous solution.
2. Due to strong acid and strong base, concentrations $[\text{H}_3\text{O}^+] = [\text{OH}^-]$ and the solution is neutral.

Question iii.

How are basic buffer solutions prepared?

Answer:

1. Basic buffer solution is prepared by mixing aqueous solutions of a weak base like NH_4OH and its salt of a strong acid like NH_4Cl .
2. A weak base is selected according to the required pH or pOH of the solution and dissociation constant of the weak base.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Question iv.

Dissociation constant of acetic acid is 1.8×10^{-5} . Calculate percent dissociation of acetic acid in 0.01 M solution.

Answer:

Given : $K_a = 1.8 \times 10^{-5}$; $C = 0.01$ M

Percent dissociation = ?

$$\begin{aligned}K_a &= \frac{C\alpha^2}{1-\alpha} \approx C\alpha^2 \\ \therefore \alpha &= \sqrt{\frac{K_a}{C}} \\ &= \left(\frac{1.8 \times 10^{-5}}{0.01} \right)^{\frac{1}{2}} \\ &= 4.242 \times 10^{-2}\end{aligned}$$

$\therefore$ Percent dissociation = $\alpha \times 100$

$$= 4.242 \times 10^{-2} \times 10^2$$

$$= 4.242\%$$

Percent dissociation = 4.242%

Question v.

Write one property of a buffer solution.

Answer:

Properties (or advantages) of a buffer solution :

- The pH of a buffer solution is maintained appreciably constant.
- By addition of a small amount of an acid or a base pH does not change.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

- On dilution with water, pH of the solution doesn't change.

Question vi.


The pH of a solution is 6.06. Calculate its H^+ ion concentration.

Question vii.

Calculate the pH of 0.01 M sulphuric acid.

Answer:

Given : $C = 0.01 \text{ M } H_2SO_4$, $pH = ?$



$$\therefore [H_3O^+] = 2 \times 0.01 = 0.02 \text{ M}$$

$$pH = -\log_{10} [H_3O^+]$$

$$= -\log_{10} 0.02$$

$$= -(2 - 0.3010)$$

$$= 2 - 0.3010$$

$$= 1.6990$$

$$pH = 1.6990.$$


Question viii.

The dissociation of H_2S is suppressed in the presence of HCl . Name the phenomenon.


Answer:

The weak dibasic acid H_2S is dissociated as follows :

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>



When HCl is added, it increases the concentration of common ion H_3O^+ .




Hence by Le Chatelier's principle, the equilibrium is shifted from right to left, suppressing the dissociation of weak electrolyte H_2S .

Question ix.


Why is it necessary to add H_2SO_4 while preparing the solution of CuSO_4 ?

Answer:

CuSO_4 is a salt of strong acid H_2SO_4 and weak base $\text{Cu}(\text{OH})_2$. CuSO_4 in aqueous solution undergoes hydrolysis and forms a precipitate of $\text{Cu}(\text{OH})_2$ and solution becomes turbid.



OR



When H_2SO_4 is added, the hydrolysis equilibrium is shifted to left hand side and $\text{Cu}(\text{OH})_2$ dissolves giving clear solution.

Question x.

Classify the following buffers into different types :

a. $\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$

b. $\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$

c. Sodium benzoate + benzoic acid

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

d. $\text{Cu}(\text{OH})_2 + \text{CuCl}_2$

Answer:

- (a) Acidic buffer ($\text{CH}_3\text{COOH} + \text{CH}_3\text{COONa}$)
- (b) Basic buffer ($\text{NH}_4\text{OH} + \text{NH}_4\text{Cl}$)
- (c) Acidic buffer (Sodium benzoate + benzoic acid)
- (d) Basic buffer ($\text{Cu}(\text{OH})_2 + \text{CuCl}_2$)

[Note : $\text{Cu}(\text{OH})_2$ being insoluble is not used to prepare a buffer solution.]

3. Answer the following in brief :


Question i.

What are acids and bases according to Arrhenius theory ?


Answer:

According to Arrhenius theory :

Acid : It is a substance which contains hydrogen and on dissolving in water produces hydrogen ions (H^+) E.g. HCl



Base : It is a substance which contains OH group and on dissolving in water produces hydroxyl ions (OH^-). E.g. NaOH




Question ii.

What is meant by conjugate acid-base pair?

Answer:



<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Conjugate acid-base pair : A pair of an acid and a base differing by a proton is called a conjugate acid-base pair.




Question iii.

Label the conjugate acid-base pair in the following reactions




Answer:



Question iv.

Write a reaction in which water acts as a base.

Answer:



Since water accepts a proton, it acts as a base.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Question v.

Ammonia serves as a Lewis base whereas AlCl_3 is Lewis acid. Explain.

Answer:

- Since ammonia molecule, NH_3 has a lone pair of electrons to donate it acts as a Lewis base.
- AlCl_3 is a molecule with incomplete octet hence it is electron deficient and acts as a Lewis acid.

Question vi.

Acetic acid is 5% ionised in its decimolar solution. Calculate the dissociation constant of acid.

Answer:

Given : $C = 0.1 \text{ M}$; Dissociation = 5%, $K_a = 2$ Percent dissociation

$$\begin{aligned}\alpha &= \frac{\text{Percent dissociation}}{100} \\ &= \frac{5}{100} = 0.05 \\ K_a &= \frac{C\alpha^2}{1 - \alpha} = \frac{0.1 \times (0.05)^2}{1 - 0.05} \\ &= \frac{0.1 \times (0.05)^2}{0.95} \\ &= 2.63 \times 10^{-4}\end{aligned}$$

Dissociation constant of acid = $K_a = 2.63 \times 10^{-4}$

Question vii.

Derive the relation $\text{pH} + \text{pOH} = 14$.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

The ionic product of water, K_w is given by,

$$K_w = [H_3O^+] \times [OH^-]$$

$$\text{At } 298 \text{ K, } K_w = 1 \times 10^{-14}$$

$$\therefore pK_w = -\log_{10} K_w = \log_{10} 1 \times 10^{-14} = 14$$

$$\therefore [H_3O^+] \times [OH^-] = 1 \times 10^{-14}$$

Taking logarithm to base 10 of both sides,

$$\log_{10} [H_3O^+] + \log_{10} [OH^-] = \log_{10} 1 \times 10^{-14}$$

Multiplying both the sides by -1,

$$-\log_{10} [H_3O^+] - \log_{10} [OH^-] = -\log_{10} 1 \times 10^{-14}$$

$$\therefore pH = -\log_{10} [H_3O^+]; pOH = -\log_{10} [OH^-];$$

$$pK_w = -\log_{10} K_w$$

$$\therefore pH + pOH = pK_w$$

$$\text{OR } pH + pOH = 14$$


Question viii.

Aqueous solution of sodium carbonate is alkaline whereas aqueous solution of ammonium chloride is acidic. Explain.

Answer:

(A) (i) Sodium carbonate is a salt of weak acid and strong base.


(ii) In aqueous solution it undergoes hydrolysis.



(iii) Strong base dissociates completely while weak acid dissociates partially since $[\text{OH}^-] > [\text{H}_3\text{O}^+]$, the solution is basic.

(B) (i) Ammonium chloride is a salt of strong acid and weak base.

(ii) In aqueous solution it undergoes hydrolysis



(iii) Since $[\text{H}^+]$ or $[\text{H}_3\text{O}^+] > [\text{OH}^-]$ the solution is acidic.

Question ix.

pH of a weak monobasic acid is 3.2 in its 0.02 M solution. Calculate its dissociation constant.

Answer:

Given : pH = 3.2; C = 0.02 M; K_a = ?

$$\text{pH} = -\log_{10} [\text{H}^+]$$

$$\therefore [\text{H}^+] = \text{Antilog} - \text{pH}$$

$$= \text{Antilog} - 3.2$$

$$= \text{Antilog } 4^{\text{---}}.8$$

$$= 6.31 \times 10^{-4}\text{M}$$

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

$$\begin{array}{ccccc} \text{HA}_{(\text{aq})} & \rightleftharpoons & \text{H}^+ & + & \text{A}^- \\ c(1-\alpha) & & c\alpha & & c\alpha \\ \therefore [\text{H}^+] & = & c\alpha & & \\ \alpha = \frac{[\text{H}^+]}{c} & = & \frac{6.31 \times 10^{-4}}{0.02} & = & 0.0315 \end{array}$$

$$\begin{aligned} K_a &= c\alpha^2 \\ &= 0.02 \times (0.0315)^2 \\ &= 1.984 \times 10^{-5} \end{aligned}$$

$$\text{Dissociation constant} = K_a = 1.984 \times 10^{-5}$$

Question x.

In NaOH solution $[\text{OH}^-]$ is 2.87×10^{-4} . Calculate the pH of solution.

Answer:

$$\text{Given : } [\text{OH}^-] = 2.87 \times 10^{-4} \text{ M, pH} = ?$$

$$\begin{aligned} \text{pOH} &= -\log_{10} [\text{OH}^-] \\ &= -\log_{10} 2.87 \times 10^{-4} \\ &= -(4 - 0.4579) \\ &= (4 - 0.4579) \\ &= 3.5421 \end{aligned}$$

$$\therefore \text{pH} + \text{pOH} = 14$$

$$\therefore \text{pH} = 14 - \text{pOH} = 14 - 3.5421 = 10.4579$$

$$\text{pH} = 10.4579.$$

4. Answer the following :

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Question i.

Define degree of dissociation. Derive Ostwald's dilution law for the CH_3COOH .

Answer:

(A) Degree of dissociation :

It is defined as a fraction of total number of moles of an electrolyte that dissociate into its ions at equilibrium.


It is denoted by α and represented by,

$$\alpha = \frac{\text{number of moles dissociated}}{\text{total number of moles of an electrolyte}}$$

$$\text{Or } \alpha = \frac{\text{Per cent dissociation}}{100}$$

$$\therefore \text{Per cent dissociation} = \alpha \times 100$$

(B) Consider $V \text{ dm}^3$ of a solution containing one mole of CH_3COOH . Then the concentration of acid is, $C = \frac{1}{V} \text{ mol dm}^3$. Let α be the degree of dissociation



If K_a is dissociation constant, then

$$K_a = \frac{[\text{CH}_3\text{COO}^-] \times [\text{H}^+]}{[\text{CH}_3\text{COOH}]}$$

$$= \frac{\frac{\alpha}{V} \times \frac{\alpha}{V}}{\frac{(1 - \alpha)}{V}} = \frac{\alpha^2}{(1 - \alpha)V}$$

$$\therefore C = \frac{1}{V} \text{ mol dm}^3$$

$$\therefore K_a = \frac{C\alpha^2}{(1 - \alpha)}$$

This is Ostwald's dilution law.

Question ii.

Define pH and pOH. Derive relationship between pH and pOH.

Answer:

(1) pH : The negative logarithm, to the base 10, of the molar concentration of hydrogen ions, H^+ is known as the pH of a solution.

$$\text{pH} = -\log_{10} [\text{H}^+]$$

(2) pOH : The negative logarithm, to the base 10, of the molar concentration of hydroxyl ions, OH^- is known as the pOH of a solution.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

$$\text{pOH} = -\log_{10} [\text{OH}^-]$$

Relationship between pH and pOH:

The ionic product of water, K_w is given by,

$$K_w = [\text{H}_3\text{O}^+] \times [\text{OH}^-]$$

$$\text{At } 298 \text{ K, } K_w = 1 \times 10^{-14}$$

$$\therefore \text{p}K_w = -\log_{10} K_w = \log_{10} 1 \times 10^{-14} = 14$$

$$\therefore [\text{H}_3\text{O}^+] \times [\text{OH}^-] = 1 \times 10^{-14}$$

Taking logarithm to base 10 of both sides,

$$\log_{10} [\text{H}_3\text{O}^+] + \log_{10} [\text{OH}^-] = \log_{10} 1 \times 10^{-14}$$

Multiplying both the sides by -1,

$$-\log_{10} [\text{H}_3\text{O}^+] - \log_{10} [\text{OH}^-] = -\log_{10} 1 \times 10^{-14}$$

$$\therefore \text{pH} = -\log_{10} [\text{H}_3\text{O}^+]; \text{pOH} = -\log_{10} [\text{OH}^-];$$

$$\text{p}K_w = -\log_{10} K_w$$

$$\therefore \text{pH} + \text{pOH} = \text{p}K_w$$

$$\text{OR } \text{pH} + \text{pOH} = 14$$

Question iii.

What is meant by hydrolysis ? A solution of $\text{CH}_3\text{COONH}_4$ is neutral. why ?


Answer:

Hydrolysis : A reaction in which the cations or anions or both the ions of a salt react with water to produce acidity or basicity or sometimes neutrality is called hydrolysis.

A salt of weak acid and weak base for which $K_a = K_b$:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Consider hydrolysis of $\text{CH}_3\text{COONH}_4$.




Since $K_a = K_b$, the weak acid CH_3COOH and weak base NH_4OH dissociate to the same extent, hence, $[\text{H}_3\text{O}^+] = [\text{OH}^-]$ and the solution reacts neutral after hydrolysis.

Question iv.

Dissociation of HCN is suppressed by the addition of HCl. Explain.

Answer:

The weak acid HCN is dissociated as follows :



The dissociation constant K_a is represented as,

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{CN}^-]}{[\text{HCN}]}$$

When HCl is added, it increases the concentration of H_3O^+ , hence in order to keep the ratio constant, then by Le Chatelier's principle, the equilibrium is shifted from right to left, suppressing the dissociation of HCN.

Question v.

Derive the relationship between degree of dissociation and dissociation constant in weak electrolytes.

Answer:

Expression of Ostwald's dilution law in the case of a weak electrolyte : Consider the dissociation of a weak electrolyte BA. Let $V \text{ dm}^3$ of a solution contain one mole of the electrolyte. Then the concentration of a solution is, $C = 1/V \text{ mol dm}^{-3}$. Let α be the degree of dissociation of the electrolyte.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Maharashtra Board Solutions Class 12 Chemistry: Chapter 3- Ionic Equilibria
Question 4.v

Applying the law of mass action to this dissociation equilibrium, we have,

$$K = \frac{[B^+][A^-]}{[BA]}$$
$$\therefore K = \frac{\frac{\alpha}{V} \times \frac{\alpha}{V}}{\frac{(1-\alpha)}{V}} = \frac{\alpha^2}{(1-\alpha)V}$$

As the electrolyte is weak, α is very small as compared to unity, $\therefore (1 - \alpha) \approx 1$.

$$\therefore K = \frac{\alpha^2}{V} \quad \therefore \alpha = \sqrt{KV} \quad \therefore \alpha \propto \sqrt{V}$$

$$\frac{1}{V} = C, \text{ where } C = \text{concentration in mol dm}^{-3}$$

$$\therefore K = \alpha^2 C \quad \therefore \alpha = \sqrt{\frac{K}{C}}$$

$$\therefore \alpha = \sqrt{K \times V}$$

$$\therefore C = \frac{1}{V} \text{ or } V = \frac{1}{C}$$

$$\alpha = \sqrt{\frac{K}{C}}$$

This is the expression of Ostwald's dilution law. Thus, the degree of dissociation of a weak electrolyte is directly proportional to the square root of the volume of the solution containing 1 mole of an electrolyte.


Question vi.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>


Sulfides of cation of group II are precipitated in acidic solution ($\text{H}_2\text{S} + \text{HCl}$) whereas sulfides of cations of group IIIB are precipitated in ammoniacal solution of H_2S . Comment on the relative values of solubility product of sulfides of these.

Answer:

- (1) In qualitative analysis, the cations of group II are precipitated as sulphides, namely HgS , CuS , PbS , etc., while cations of group IIIB are precipitated as sulphides, namely, CoS , NiS , ZnS .
- (2) The sulphides of group II have extremely low solubility product (K_{sp}) about 10^{-29} to 10^{-53} while the sulphides of group IIIB have slightly higher K_{sp} values about 10^{-20} to 10^{-23} .
- (3) In group II, sulphides are precipitated by adding H_2S in acidic solution while in IIIB group they are precipitated in a basic solution like ammoniacal solution.
- (4) In acidic medium due to common ion H^+ , H_2S is dissociated to very less extent but gives sufficient S^{2-} ion to exceed solubility product of group II sulphides of cations and precipitate them.



- (5) In basic medium, H^+ from H_2S are removed by OH^- in solution, or by NH_4OH , increasing the dissociation of H_2S and concentration of S^{2-} , so that $\text{IP} > K_{\text{sp}}$.



- (6) Therefore group II cations are precipitated in an acidic medium while cations of group IIIB are precipitated in ammoniacal solution.


Question vii.

Solubility of a sparingly soluble salt get affected in presence of a soluble salt having one common ion. Explain.

Answer:

Consider the solubility equilibrium of a sparingly soluble salt, AgCl .


<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>



The solubility product, K_{sp} is given by,

$$K_{sp} = [\text{Ag}^+] \times [\text{Cl}^-]$$

Consider addition of a strong electrolyte AgNO_3 with a common ion Ag^+ .



The concentration Ag^+ in the solution is increased, hence by Le Chatelier's principle the equilibrium of AgCl is shifted to left hand side since the value of K_{sp} is constant.

Thus in the presence of a common ion, the solubility of a sparingly soluble salt is suppressed.

Question viii.

The pH of rain water collected in a certain region of Maharashtra on particular day was 5.1. Calculate the H_3O^+ ion concentration of the rain water and its percent dissociation.

Answer:

Given : $\text{pH} = 5.1$, $[\text{H}_3\text{O}^+] = ?$

$$\text{pH} = -\log_{10} [\text{H}_3\text{O}^+]$$

$$\therefore \log_{10} [\text{H}_3\text{O}^+] = -\text{pH}$$

$$\therefore [\text{H}_3\text{O}^+] = \text{Antilog} - \text{pH}$$

$$= \text{Antilog} - 5.1$$

$$= \text{Antilog } 6^{\text{---}}.9$$

$$= 7.943 \times 10^{-6} \text{ M}$$

(H_3O^+ in rainwater is due to dissolved gases, CO_2 , SO_2 , etc. forming acids which dissociate giving H_3O^+ and acidity to rainwater.)

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

$$[\text{H}_3\text{O}^+] = 7.943 \times 10^{-4} \text{ M}$$

Question ix.

Explain the relation between ionic product and solubility product to predict whether a precipitate will form when two solutions are mixed?

Answer:

If ionic product and solubility product are indicated by IP and K_{sp} respectively then,

(I) When $\text{IP} = K_{sp}$, the solution is saturated.

(II) When $\text{IP} > K_{sp}$, the solution is supersaturated and hence precipitation will occur, when two solutions are mixed.

(III) When $\text{IP} < K_{sp}$, the solution is unsaturated and precipitation will not occur, when two solutions are mixed.

Use your brain power (Textbook Page No. 47)

Question 1.

Which of the following is a strong electrolyte ?

HF, AgCl, CuSO_4 , $\text{CH}_3\text{COONH}_4$, H_3PO_4 .

Answer:

$\text{CH}_3\text{COONH}_4$ is a strong electrolyte since in aqueous solution it dissociates completely. Sparingly soluble salts AgCl, CuSO_4 are also strong electrolytes.

Use your brain power (Textbook Page No. 49)


Question 1.

All Bronsted bases are also Lewis bases, but all Bronsted acids are not Lewis acids. Explain.


Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

NH_3 is a Bronsted base since it can accept a proton while it is also a Lewis base since it has a lone pair of electrons to donate.




Bronsted base



Lewis base

(2) HCl is a Bronsted acid since it can donate a proton but it is not a Lewis acid since it can't accept a pair of electrons.



Bronsted acid

Use your brain power (Textbook Page No. 53)

Question 1.

Suppose that pH of monobasic and dibasic acid is the same. Does this mean that the molar concentrations of both acids are identical ?

Answer:

Even if monobasic acid and dibasic acid give same pH, their molar concentrations are different. One mole of monobasic acid like HCl gives 1 mol of H^+ while one mole of dibasic acid gives 2 mol of H^+ in solution. Hence the concentration of dibasic acid will be half of the concentration of monobasic acid. For example, for same pH. $[\text{Monobasic acid}] = [\text{Dibasic acid}]/2$

Question 2.

How does pH of pure water vary with temperature ? Explain.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Since the increase in temperature, increases the dissociation of water, its pH decreases.

Can you tell ? (Textbook Page No. 54)

Question 1.


Why (i) an aqueous solution of NH_4Cl is acidic.

(ii) while that of HCOOK basic ?

Answer:


(i) (i) Ammonium chloride is a salt of strong acid and weak base.

(ii) In aqueous solution it undergoes hydrolysis



(iii) Since $[\text{H}^+]$ or $[\text{H}_3\text{O}^+] > [\text{OH}^-]$ the solution is acidic.

(ii) HCOOK is a salt of weak acid HCOOH and strong base KOH . In aqueous solution it undergoes hydrolysis giving weak acid and strong base KOH which dissociates completely,



$\therefore [\text{OH}^-] > [\text{H}_3\text{O}^+]$, and the solution reacts basic.

Can you think ? (Textbook Page No. 56)

Question 1.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

Home made jams and jellies without any added chemical preservative additives spoil in a few days whereas commercial jams and jellies have a long shelf life. Explain. What role does added sodium benzoate play ?

Answer:

Sodium benzoate added to jams and jellies in commercial products maintains the pH constant and acts as a preservative. Hence jams and jellies are not spoiled for a very long time unlike homemade products.

Can you tell ? (Textbook Page No. 56)

Question 1.

It is enough to add a few mL of a buffer solution to maintain its pH. Which property of buffer is used here ?

Answer:

The important property of reserve acidity and reserve basicity of a buffer solution is used to maintain constant pH. Weak acid or weak base along with ions (cations or anions) from salt react with excess of added acid (H^+) or base [OH^-] and maintain pH constant.


Use your brain power (Textbook Page No. 59)

Question 1.

What is the relationship between molar solubility and solubility product for salts given below : (i) Ag_2CrO_4 (ii) $Ca_3(PO_4)_2$ (iii) $Cr(OH)_3$.

Answer:

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>




$$\therefore K_{sp} = [\text{Ag}^+]^2 \times [\text{CrO}_4^{2-}]$$

$$\therefore K_{sp} = x^2 \times y \times (S)$$

$$= 2^2 \times 1 \times S^3$$

$$= 4S^3$$

$$\therefore S = \left(\frac{K_{sp}}{4}\right)^{\frac{1}{3}} \text{ mol dm}^3$$




$$\therefore K_{sp} = [\text{Ca}^{2+}]^3 \times [\text{PO}_4^{3-}]^2$$

$$\therefore K_{sp} = x^x \cdot y^y \cdot (S)^{x+y}$$

$$= 3^3 \cdot 2^2 \cdot (S)^{3+2}$$

$$= 108 S^5$$

$$\therefore S = \left(\frac{K_{sp}}{108}\right)^{\frac{1}{5}} \text{ mol dm}^{-3}$$



$$K_{sp} = x^x \cdot y^y \cdot (S)^{x+y}$$

$$= 1^1 \cdot 3^3 \cdot (S)^{1+3}$$

$$= 9 S^4$$

$$\therefore S = \left(\frac{K_{sp}}{9}\right)^{\frac{1}{4}} \text{ mol dm}^{-3}$$

Can you tell ? (Textbook Page No. 60)


Question 1.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

How is the ionization of NH_4OH suppressed by addition of NH_4Cl to the solution of NH_4OH ?

Answer:


Ionisation of NH_4OH is represented as follows :



It has ionisation constant,

$$K_b = \frac{[\text{NH}_4^+] \times [\text{OH}^-]}{[\text{NH}_4\text{OH}]}$$

K_b has constant value at constant temperature. When strong electrolyte NH_4Cl is added to its solution, it dissociates completely.



Due to common ion NH_4^+ , by Le Chatelier's principle, the equilibrium is shifted from right to left, suppressing the ionisation of NH_4OH .



Maharashtra Board Solutions

Class 12 Chemistry

- Chapter 1- Solid State
- Chapter 2- Solutions
- Chapter 3- Ionic Equilibria
- Chapter 4- Chemical Thermodynamics
- Chapter 5- Electrochemistry
- Chapter 6- Chemical Kinetics
- Chapter 7- Elements of Groups 16, 17 and 18
- Chapter 8- Transition and Inner Transition Elements
- Chapter 9- Coordination Compounds
- Chapter 10- Halogen Derivatives
- Chapter 11- Alcohols, Phenols and Ethers
- Chapter 12- Aldehydes, Ketones and Carboxylic Acids
- Chapter 13- Amines
- Chapter 14- Biomolecules
- Chapter 15- Introduction to Polymer Chemistry
- Chapter 16- Green Chemistry and Nanochemistry

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

About About Maharashtra State Board (MSBSHSE)

The Maharashtra State Board of Secondary and Higher Secondary Education or MSBSHSE (Marathi: महाराष्ट्र राज्य माध्यमिक आणि उच्च माध्यमिक शिक्षण मंडळ), is an **autonomous and statutory body established in 1965**. The board was amended in the year 1977 under the provisions of the Maharashtra Act No. 41 of 1965.

The Maharashtra State Board of Secondary & Higher Secondary Education (MSBSHSE), Pune is an independent body of the Maharashtra Government. There are more than 1.4 million students that appear in the examination every year. The Maha State Board conducts the board examination twice a year. This board conducts the examination for SSC and HSC.

The Maharashtra government established the Maharashtra State Bureau of Textbook Production and Curriculum Research, also commonly referred to as Ebalbharati, in 1967 to take up the responsibility of providing quality textbooks to students from all classes studying under the Maharashtra State Board. MSBHSE prepares and updates the curriculum to provide holistic development for students. It is designed to tackle the difficulty in understanding the concepts with simple language with simple illustrations. Every year around 10 lakh students are enrolled in schools that are affiliated with the Maharashtra State Board.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

FAQs

Where do I get the Maharashtra State Board Books PDF For free download?

You can download the Maharashtra State Board Books from the eBalbharti official website, i.e. cart.ebalbharati.in or from this article.

Add image

How to Download Maharashtra State Board Books?

Students can get the Maharashtra Books for primary, secondary, and senior secondary classes from here. You can view or download the Maharashtra State Board Books from this page or from the official website for free of cost. Students can follow the detailed steps below to visit the official website and download the e-books for all subjects or a specific subject in different mediums.

Step 1: Visit the official website ebalbharati.in

Step 2: On the top of the screen, select "Download PDF textbooks"

Step 3: From the "Classes" section, select your class.

Step 4: From "Medium", select the medium suitable to you.

Step 5: All Maharashtra board books for class 11th will now be displayed on the right side.

Step 6: Click on the "Download" option to download the PDF book.

Who developed the Maharashtra State board books?

As of now, the MSCERT and Balbharti are responsible for the syllabus and textbooks of Classes 1 to 8, while Classes 9 and 10 are under the Maharashtra State Board of Secondary and Higher Secondary Education (MSBSHSE).

How many state boards are there in Maharashtra?

The Maharashtra State Board of Secondary & Higher Secondary Education, conducts the HSC and SSC Examinations in the state of Maharashtra through its nine Divisional Boards located at Pune, Mumbai, Aurangabad, Nasik, Kolhapur, Amravati, Latur, Nagpur and Ratnagiri.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>

About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to **info@indcareer.com**

Postal Address

IndCareer.com
52, Shilpa Nagar,
Somalwada
Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888

Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-12-chemistry-chapter-3-ionic-equilibria/>