


Maharashtra Board Solutions Class 11-Physics: Chapter 2- Mathematical Methods

Class 11 - Chapter 2 Mathematical Methods



For any clarifications or questions you can write to info@indcareer.com

Postal Address

IndCareer.com, 52, Shilpa Nagar, Somalwada Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888, **Website:** <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>



Maharashtra Board Solutions Class 11-Physics: Chapter 2- Mathematical Methods

Class 11: Physics Chapter 2 solutions. Complete Class 11 Physics Chapter 2 Notes.

Maharashtra Board Solutions Class 11-Physics: Chapter 2- Mathematical Methods

Class 11: Physics Chapter 2 solutions. Complete Class 11 Physics Chapter 2 Notes.

1. Choose the correct option.

Question 1.

The resultant of two forces 10 N and 15 N acting along + x and – x-axes respectively, is

- (A) 25 N along + x-axis
- (B) 25 N along – x-axis
- (C) 5 N along + x-axis
- (D) 5 N along – x-axis

Answer:

- (D) 5 N along – x-axis

Question 2.

For two vectors to be equal, they should have the

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>

- (A) same magnitude
- (B) same direction
- (C) same magnitude and direction
- (D) same magnitude but opposite direction

Answer:

- (C) same magnitude and direction

Question 3.

The magnitude of scalar product of two unit vectors perpendicular to each other is

- (A) zero
- (B) 1
- (C) -1
- (D) 2

Answer:

- (A) zero

Question 4.

The magnitude of vector product of two unit vectors making an angle of 60° with each other is

- (A) 1

- (C) $\frac{3}{2}$
(D) $\frac{\sqrt{3}}{2}$

Answer:

- (D) $\frac{\sqrt{3}}{2}$

Question 5.

If \vec{A} , \vec{B} , and \vec{C} are three vectors, then which of the following is not correct?

- (A) $\vec{A} \cdot (\vec{B} + \vec{C}) = \vec{A} \cdot \vec{B} + \vec{A} \cdot \vec{C}$
(B) $\vec{A} \cdot \vec{B} = \vec{B} \cdot \vec{A}$
(C) $\vec{A} \times \vec{B} = \vec{B} \times \vec{A}$
(D) $\vec{A} \times (\vec{B} \times \vec{C}) = \vec{A} \times \vec{B} + \vec{B} \times \vec{C}$

Answer:

$$(c) \vec{A} \times \vec{B} = \vec{B} \times \vec{A}$$

2. Answer the following questions.

Question 1.

Show that $\vec{A} = \frac{\hat{i} - \hat{j}}{\sqrt{2}}$ is a unit vector.

Solution:

Let \hat{a} be unit vector of \vec{a} .

$$\therefore \hat{a} = \frac{\vec{a}}{|\vec{a}|}$$

$$\text{Now, } |\vec{a}| = \sqrt{a_x^2 + a_y^2} = \sqrt{\left(\frac{1}{\sqrt{2}}\right)^2 + \left(\frac{-1}{\sqrt{2}}\right)^2} = 1$$

$$\therefore \hat{a} = \frac{\vec{a}}{1} \Rightarrow \vec{a} \text{ itself is a unit vector.}$$

Question 2.

If $\vec{v}_1 = 3\hat{i} + 4\hat{j} + \hat{k}$ and $\vec{v}_2 = \hat{i} - \hat{j} - \hat{k}$, determine the magnitude of $\vec{v}_1 + \vec{v}_2$.

Solution:

$$\begin{aligned}\vec{v}_1 + \vec{v}_2 &= (3\hat{i} + 4\hat{j} + \hat{k}) + (\hat{i} - \hat{j} - \hat{k}) \\ &= 3\hat{i} + 3\hat{i} + 4\hat{j} - \hat{j} + \hat{k} - \hat{k} \\ &= 4\hat{i} + 3\hat{j}\end{aligned}$$

\therefore Magnitude of $(\vec{v}_1 + \vec{v}_2)$,

$$|\vec{v}_1 + \vec{v}_2| = \sqrt{4^2 + 3^2} = \sqrt{25} = 5 \text{ units.}$$

Answer:

Magnitude of $\vec{v}_1 + \vec{v}_2 = 5$ units.

Question 3.

For $\vec{v}_1 = 2\hat{i} - 3\hat{j}$ and $\vec{v}_2 = -6\hat{i} + 5\hat{j}$, determine the magnitude and direction of $\vec{v}_1 + \vec{v}_2$.

Answer:

$$\vec{v}_1 + \vec{v}_2 = (2\hat{i} - 3\hat{j}) + (-6\hat{i} + 5\hat{j})$$

$$= (2\hat{i} - 6\hat{i}) + (-3\hat{j} + 5\hat{j})$$

$$= -4\hat{i} + 2\hat{j}$$

$$\therefore |\vec{v}_1 + \vec{v}_2| = \sqrt{(-4)^2 + 2^2} = \sqrt{20} = \sqrt{4 \times 5} = 2\sqrt{5}$$

Comparing $\vec{v}_1 + \vec{v}_2$, with $\vec{R} = R_x\hat{i} + R_y\hat{j}$

$$\Rightarrow R_x = -4 \text{ and } R_y = 2$$

Taking θ to be angle made by \vec{R} with X-axis,

$$\therefore \theta = \tan^{-1} \left(\frac{R_y}{R_x} \right) = \tan^{-1} \left(\frac{2}{-4} \right)$$

$$= \tan^{-1} \left(-\frac{1}{2} \right) \text{ with X - axis}$$

Answer:

Magnitude and direction of $\vec{v}_1 + \vec{v}_2$, is

respectively $2\sqrt{5}$ and $\tan^{-1} \left(-\frac{1}{2} \right)$ with X - axis.

Question 4.

Find a vector which is parallel to $\vec{v} = \hat{i} - 2\hat{j}$ and has a magnitude 10.

Answer:

$$\text{Let the vector be } \vec{w} = w_x\hat{i} + w_y\hat{j}$$

$$|\vec{w}| = \sqrt{w_x^2 + w_y^2} = 10 \quad \dots \text{(Given)}$$

$$\therefore w_x^2 + w_y^2 = 100 \quad \dots \text{(i)}$$

$$\text{Also, } \vec{v} \cdot \vec{w} = vw$$

$$\dots (\because |\vec{v}| \text{ and } |\vec{w}| \text{ are parallel vectors})$$

$$\Rightarrow (\hat{i} - 2\hat{j}) \cdot (w_x \hat{i} + w_y \hat{j}) = \sqrt{(1)^2 + (-2)^2} \times 10$$

$$\dots \left(\because |\vec{v}| = \sqrt{(1)^2 + (-2)^2} \right)$$

$$\therefore w_x - 2w_y = 10\sqrt{5} \quad \dots \text{(ii)}$$

Substituting for w_x in (i) using equation (ii),

$$(10\sqrt{5} + 2w_y)^2 + w_y^2 = 100$$

$$\therefore 500 + 40\sqrt{5} w_y + 4w_y^2 - 100 = 0$$

$$\therefore 5w_y^2 + 40\sqrt{5} w_y + 400 = 0$$

$$\therefore w_y^2 + 8\sqrt{5} w_y + 80 = 0$$

Using factorisation formula,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$w_y = \frac{-8\sqrt{5} \pm \sqrt{(8\sqrt{5})^2 - 4 \times 1 \times 80}}{2 \times 1}$$

$$= w_y = \frac{-8\sqrt{5} \pm 0}{2} = -4\sqrt{5} = \frac{-20}{\sqrt{5}}$$

Using equation (ii),

$$\begin{aligned} w_x &= 10\sqrt{5} + 2\left(\frac{-20}{\sqrt{5}}\right) \\ &= 10\sqrt{5} - \frac{40}{\sqrt{5}} \\ &= \frac{(10\sqrt{5} \times \sqrt{5}) - 40}{\sqrt{5}} = \frac{50 - 40}{\sqrt{5}} = \frac{10}{\sqrt{5}} \end{aligned}$$

$$\therefore \vec{w} = w_x \hat{i} + w_y \hat{j} = \frac{10}{\sqrt{5}} \hat{i} - \frac{20}{\sqrt{5}} \hat{j}$$

Answer:

Required vector is $\frac{10}{\sqrt{5}} \hat{i} - \frac{20}{\sqrt{5}} \hat{j}$

Alternate method:

When two vectors are parallel, one vector is scalar multiple of another,

i.e., if \vec{v} and \vec{w} are parallel then, $\vec{w} = n \vec{v}$ where, n is scalar.

This means, $\vec{w} = n v_x \hat{i} + n v_y \hat{j}$

$$= n\hat{i} - 2n\hat{j} \quad \dots (\because v_x = 1, v_y = 2)$$

$$\therefore |\vec{w}| = \sqrt{(n)^2 + (-2n)^2} = \sqrt{5} n$$

$$\text{Given: } |\vec{w}| = 10$$

$$\therefore n = \frac{10}{\sqrt{5}} = 2\sqrt{5}$$

$$\begin{aligned}\therefore \vec{w} &= 2\sqrt{5}\hat{i} - 2(2\sqrt{5})\hat{j} \\ &= 2\sqrt{5}\hat{i} - 4\sqrt{5}\hat{j} \\ &= \frac{2\sqrt{5} \times \sqrt{5}}{\sqrt{5}}\hat{i} - \frac{4\sqrt{5} \times \sqrt{5}}{\sqrt{5}}\hat{j} \\ \therefore \vec{w} &= \frac{10}{\sqrt{5}}\hat{i} - \frac{20}{\sqrt{5}}\hat{j}\end{aligned}$$

Question 5.

Show that vectors $\vec{a} = 2\hat{i} + 5\hat{j} - 6\hat{k}$ and $\vec{b} = \hat{i} + \frac{5}{2}\hat{j} - 3\hat{k}$ are parallel.

Answer:

Let angle between two vectors be θ .

$$\begin{aligned}\therefore \cos \theta &= \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} \\ &= \frac{(2\hat{i} + 5\hat{j} - 6\hat{k}) \cdot (\hat{i} + \frac{5}{2}\hat{j} - 3\hat{k})}{\sqrt{2^2 + 5^2 + (-6)^2} \times \sqrt{1^2 + (\frac{5}{2})^2 + (-3)^2}} \\ &= \frac{2 + \frac{25}{2} + 18}{\sqrt{65} \times \sqrt{65/4}}\end{aligned}$$

$$= \frac{65/2}{65/2} = 1$$

$$\Rightarrow \theta = \cos^{-1}(1) = 0^\circ$$

\Rightarrow Two vectors are parallel.

Alternate method:

$$\vec{a} = 2(\hat{i} + \frac{5}{2}\hat{j} + \hat{k}) = 2\vec{b}$$

Since \vec{a} is a scalar multiple of \vec{b} , the vectors are parallel.

3. Solve the following problems.

Question 1.

Determine $\vec{a} \times \vec{b}$, given $\vec{a} = 2\hat{i} + 3\hat{j}$ and $\vec{b} = 3\hat{i} + 5\hat{j}$.

Answer:

Using determinant for vectors in two dimensions,

$$\begin{aligned}\vec{a} \times \vec{b} &= \begin{vmatrix} \hat{i} & \hat{j} \\ a_x & a_y \\ b_x & b_y \end{vmatrix} = \begin{vmatrix} \hat{i} & \hat{j} \\ 2 & 3 \\ 3 & 5 \end{vmatrix} \\ &= [(2 \times 5) - (3 \times 3)]\hat{k} = (10 - 9)\hat{k} = \hat{k}\end{aligned}$$

Answer:

$\vec{a} \times \vec{b}$ gives \hat{k}

Question 2.

Show that vectors $\vec{a} = 2\hat{i} + 3\hat{j} + 6\hat{k}$, $\vec{b} = 3\hat{i} - 6\hat{j} + 2\hat{k}$ and $\vec{c} = 6\hat{i} + 2\hat{j} - 3\hat{k}$ are mutually perpendicular.

Solution:

As dot product of two perpendicular vectors is zero. Taking dot product of \vec{a} and \vec{b}

$$\begin{aligned}\vec{a} \cdot \vec{b} &= (2\hat{i} + 3\hat{j} + 6\hat{k}) \cdot (3\hat{i} - 6\hat{j} + 2\hat{k}) \\ &= (2\hat{i} \times 3\hat{i}) + (3\hat{j} \times -6\hat{j}) + (6\hat{k} \times 2\hat{k}) \\ &\quad \dots (\because \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0) \\ &= 6 - 18 + 12 \quad \dots (\because \hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1) \\ &= 0\end{aligned}$$

$$\begin{aligned}\text{Similarly, } \vec{b} \cdot \vec{c} &= (3\hat{i} - 6\hat{j} + 2\hat{k}) \cdot (6\hat{i} + 2\hat{j} - 3\hat{k}) \\ &= (3\hat{i} \times 6\hat{i}) + (-6\hat{j} \times 2\hat{j}) + (2\hat{k} \times -3\hat{k}) \\ &\quad \dots (\because \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0) \\ &= 18 - 12 - 6 \quad \dots (\because \hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1) \\ &= 0\end{aligned}$$

Combining two results, we can say that given three vectors \vec{a} , \vec{b} , and \vec{c} are mutually perpendicular to each other.

Question 3.

Determine the vector product of $\vec{v}_1 = 2\hat{i} + 3\hat{j} - \hat{k}$ and $\vec{v}_2 = \hat{i} + 2\hat{j} - 3\hat{k}$ are perpendicular to each other, determine the value of a.

Solution:

$$\text{As } \vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ A_x & A_y & A_z \\ B_x & B_y & B_z \end{vmatrix}$$

Using determinant to find vector product,

$$\begin{aligned} \therefore \vec{v}_1 \times \vec{v}_2 &= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & 3 & -1 \\ 1 & 2 & -3 \end{vmatrix} \\ &= [(3 \times -3) - (-1 \times 2)]\hat{i} + [(-1 \times 1) - (2 \times -3)]\hat{j} \\ &\quad + [(2 \times 2) - (3 \times 1)]\hat{k} \\ &= [-9 + 2]\hat{i} + [-1 + 6]\hat{j} + [4 - 3]\hat{k} \\ &= -7\hat{i} + 5\hat{j} + \hat{k} \end{aligned}$$

Answer:

Required vector product is $-7\hat{i} + 5\hat{j} + \hat{k}$

Question 4.

Given $\vec{v}_1 = 5\hat{i} + 2\hat{j}$ and $\vec{v}_2 = a\hat{i} - 6\hat{j}$ are perpendicular to each other, determine the value of a.

Solution:

As \vec{v}_1 and \vec{v}_2 are perpendicular to each other, $\theta = 90^\circ$

$$\vec{v}_1 \cdot \vec{v}_2 = 0$$

$$\begin{aligned}\therefore (5\hat{i} + 2\hat{j}) \cdot (a\hat{i} - 6\hat{j}) &= 0 \\ \therefore (5\hat{i} \cdot a\hat{i}) + (2\hat{j} \cdot -6\hat{j}) &= 0 \quad \dots (\because \hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{i} = 0) \\ \therefore 5a + (-12) &= 0 \quad \dots (\because \hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = 1) \\ \therefore 5a &= 12 \\ \therefore a &= \frac{12}{5}\end{aligned}$$

Answer:

Value of a is $\frac{12}{5}$.

Question 5.

Obtain derivatives of the following functions:

(i) $x \sin x$

(ii) $x^4 + \cos x$

(iii) $x/\sin x$

Answer:

(i) $x \sin x$

Solution:

$$\frac{d}{dx} [f_1(x) \times f_2(x)] = f_1(x) \frac{df_2(x)}{dx} + \frac{df_1(x)}{dx} f_2(x)$$

For $f_1(x) = x$ and $f_2(x) = \sin x$

$$\frac{d}{dx} (x \sin x) = x \frac{d(\sin x)}{dx} + \frac{d(x)}{dx} \sin x$$

$$= x \cos x + 1 \times \sin x$$

$$= \sin x + x \cos x$$

(ii) $x^4 + \cos x$

Solution:

$$\text{Using } \frac{d}{dx} [f_1(x) + f_2(x)] = \frac{df_1(x)}{dx} + \frac{df_2(x)}{dx}$$

For $f_1(x) = x^4$ and $f_2(x) = \cos x$

$$\frac{d}{dx} (x^4 + \cos x) = \frac{d(x^4)}{dx} + \frac{d(\cos x)}{dx}$$

$$= 4x^3 - \sin x$$

(iii) $\frac{x}{\sin x}$

Solution:

$$\text{Using } \frac{d}{dx} \left[\frac{f_1(x)}{f_2(x)} \right] = \frac{1}{f_2(x)} \frac{df_1(x)}{dx} - \frac{f_1(x)}{f_2^2(x)} \frac{df_2(x)}{dx}$$

For $f_1(x) = x$ and $f_2(x) = \sin x$

For $f_1(x) = x$ and $f_2(x) = \sin x$

$$\begin{aligned}\therefore \frac{d}{dx} \left(\frac{x}{\sin x} \right) &= \frac{1}{\sin x} \times \frac{d(x)}{dx} - \frac{x}{\sin^2 x} \times \frac{d(\sin x)}{dx} \\ &= \frac{1}{\sin x} \times 1 - \frac{x}{\sin^2 x} \times \cos x \\ &\quad \dots \left[\because \frac{d}{dx}(\sin x) = \cos x \right] \\ &= \frac{1}{\sin x} - \frac{x \cos x}{\sin^2 x}\end{aligned}$$

[Note: As derivative of $(\sin x)$ is $\cos x$, negative sign that occurs in rule for differentiation for quotient of two functions gets retained in final answer]

Question 6.

Using the rule for differentiation for quotient of two functions, prove that $\frac{d}{dx} \left(\frac{\sin x}{\cos x} \right) = \sec^2 x$

Solution:

Using,

$$\frac{d}{dx} \left[\frac{f_1(x)}{f_2(x)} \right] = \frac{1}{f_2(x)} \frac{df_1(x)}{dx} - \frac{f_1(x)}{f_2^2(x)} \frac{df_2(x)}{dx}$$

For $f_1(x) = \sin x$ and $f_2(x) = \cos x$

$$\begin{aligned}\frac{d}{dx} \left(\frac{\sin x}{\cos x} \right) &= \frac{1}{\cos x} \times \frac{d(\sin x)}{dx} - \frac{\sin x}{\cos^2 x} \times \frac{d(\cos x)}{dx} \\ &= \frac{1}{\cos x} \times \cos x - \frac{\sin x}{\cos^2 x} \times (-\sin x) \\ &= 1 + \frac{\sin^2 x}{\cos^2 x} = \frac{\cos^2 x + \sin^2 x}{\cos^2 x}\end{aligned}$$

$$\begin{aligned}\therefore \frac{d}{dx} \left(\frac{\sin x}{\cos x} \right) &= \frac{1}{\cos^2 x} \quad \dots (\sin^2 x + \cos^2 x = 1) \\ &= \sec^2 x \quad \dots \left(\because \frac{1}{\cos x} = \sec x \right)\end{aligned}$$

Question 7.

Evaluate the following integral:

(i) $\int_0^{\pi/2} \sin x dx$

(ii) $\int_1^5 x dx$

Answer:

(i) $\int_0^{\pi/2} \sin x dx$

Solution:

$$\text{Using } \int_a^b f(x) dx = F(x) \Big|_a^b$$

$$\therefore \int_0^{\pi/2} \sin x dx = -\cos x \Big|_0^{\pi/2} = -\left[\cos\left(\frac{\pi}{2}\right) - \cos 0\right]$$

$$\text{Since, } \cos\left(\frac{\pi}{2}\right) = 0 \text{ and } \cos 0 = 1$$

$$\int_0^{\pi/2} \sin x dx = -(0 - 1) = 1$$

(ii) $\int_1^5 x dx$

Solution:

$$\text{Using, } \int_a^b f(x) dx = F(x) \Big|_a^b$$

$$\begin{aligned} \int_1^5 x dx &= \frac{x^2}{2} \Big|_1^5 \\ &= \frac{5^2}{2} - \frac{1^2}{2} \end{aligned}$$

$$\begin{aligned} &= \frac{25-1}{2} \\ &= \frac{24}{2} \\ &= 12 \end{aligned}$$

Can you recall? (Textbook Page No. 16)

Question 1.

Define scalars and vectors.

Answer:

1. Physical quantities which can be completely described by their magnitude (a number and unit) are called scalars.
2. Physical quantities which need magnitude, as well as direction for their complete description, are called vectors.

Question 2.

Which of the following are scalars or vectors?

Displacements, distance travelled, velocity, speed, force, work done, energy

Answer:

1. Scalars: Distance travelled, speed, work done, energy.
2. Vectors: Displacement, velocity, force.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>

Question 3.

What is the difference between a scalar and a vector?

Answer:

No.	Scalars	Vectors
i.	It has magnitude only	It has magnitude as well as direction.
ii.	Scalars can be added or subtracted according to the rules of the algebra.	Vectors are added or subtracted by the geometrical (graphical) method or vector algebra.
iii.	It has no specific representation.	It is represented by the symbol (\rightarrow) arrow.
iv.	The division of a scalar by another scalar is valid.	The division of a vector by another vector is not valid.
	Example: Length, mass, time, volume, etc.	Example: Displacement, velocity, acceleration, force, etc.

Internet my friend (Textbook page no. 28)

1. hyperphysics.phy-astr.gsu.edu/hbase/vect.html#veccon
2. hyperphysics.phy-astr.gsu.edu/hbase/hframe.html

Answer:

[Students can use links given above as a reference and collect information about mathematical

methods.



<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>

Maharashtra Board Solutions

Class 11 Physics

- Chapter 1- Units and Measurements
- Chapter 2- Mathematical Methods
- Chapter 3- Motion in a Plane
- Chapter 4- Laws of Motion
- Chapter 5- Gravitation
- Chapter 6- Mechanical Properties of Solids
- Chapter 7- Thermal Properties of Matter
- Chapter 8- Sound
- Chapter 9- Optics
- Chapter 10- Electrostatics
- Chapter 11- Electric Current Through Conductors
- Chapter 12- Magnetism
- Chapter 13- Electromagnetic Waves and Communication System
- Chapter 14- Semiconductors

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>

About About Maharashtra State Board (MSBSHSE)

The Maharashtra State Board of Secondary and Higher Secondary Education or MSBSHSE (Marathi: महाराष्ट्र राज्य माध्यमिक आणि उच्च माध्यमिक शिक्षण मंडळ), is an **autonomous and statutory body established in 1965**. The board was amended in the year 1977 under the provisions of the Maharashtra Act No. 41 of 1965.

The Maharashtra State Board of Secondary & Higher Secondary Education (MSBSHSE), Pune is an independent body of the Maharashtra Government. There are more than 1.4 million students that appear in the examination every year. The Maha State Board conducts the board examination twice a year. This board conducts the examination for SSC and HSC.

The Maharashtra government established the Maharashtra State Bureau of Textbook Production and Curriculum Research, also commonly referred to as Ebalbharati, in 1967 to take up the responsibility of providing quality textbooks to students from all classes studying under the Maharashtra State Board. MSBHSE prepares and updates the curriculum to provide holistic development for students. It is designed to tackle the difficulty in understanding the concepts with simple language with simple illustrations. Every year around 10 lakh students are enrolled in schools that are affiliated with the Maharashtra State Board.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>

FAQs

Where do I get the Maharashtra State Board Books PDF For free download?

You can download the Maharashtra State Board Books from the eBalbharti official website, i.e. cart.ebalbharati.in or from this article.

Add image

How to Download Maharashtra State Board Books?

Students can get the Maharashtra Books for primary, secondary, and senior secondary classes from here. You can view or download the Maharashtra State Board Books from this page or from the official website for free of cost. Students can follow the detailed steps below to visit the official website and download the e-books for all subjects or a specific subject in different mediums.

Step 1: Visit the official website ebalbharati.in

Step 2: On the top of the screen, select "Download PDF textbooks"

Step 3: From the "Classes" section, select your class.

Step 4: From "Medium", select the medium suitable to you.

Step 5: All Maharashtra board books for class 11th will now be displayed on the right side.

Step 6: Click on the "Download" option to download the PDF book.

Who developed the Maharashtra State board books?

As of now, the MSCERT and Balbharti are responsible for the syllabus and textbooks of Classes 1 to 8, while Classes 9 and 10 are under the Maharashtra State Board of Secondary and Higher Secondary Education (MSBSHSE).

How many state boards are there in Maharashtra?

The Maharashtra State Board of Secondary & Higher Secondary Education, conducts the HSC and SSC Examinations in the state of Maharashtra through its nine Divisional Boards located at Pune, Mumbai, Aurangabad, Nasik, Kolhapur, Amravati, Latur, Nagpur and Ratnagiri.

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>

About IndCareer

IndCareer.com is a leading developer of online career guidance resources for the Indian marketplace. Established in 2007, IndCareer.com is currently used by over thousands of institutions across India, including schools, employment agencies, libraries, colleges and universities.

IndCareer.com is designed to assist you in making the right career decision - a decision that meets your unique interests and personality.

For any clarifications or questions you can write to **info@indcareer.com**

Postal Address

IndCareer.com
52, Shilpa Nagar,
Somalwada
Nagpur - 440015
Maharashtra, India

WhatsApp: +91 9561 204 888

Website: <https://www.indcareer.com>

<https://www.indcareer.com/schools/maharashtra-board-solutions-class-11-physics-chapter-2-mathematical-methods/>